

1. Maxima and Minima

→ Assignment 1

2. Integration

→ Assignment 2

3. Polar equation

4. Mean Value Theorem

→ Rolle's Theorem

→ Lagrange's M.V.T.

5. Conic Section

→ Parabola

→ Ellipse

→ Hyperbola

Derivatives

Formulae:

i) Power Rule:

$$\frac{d(u^n)}{du} = nu^{n-1}$$

$$\text{e.g. } \frac{d u^8}{du} = 8u^7$$

ii) Constant Rule:

$$\frac{d(a)}{du} = 0, \text{ e.g. } \frac{d(8)}{du} = 0$$

iii) Addition or Subtraction Rule:

$$\begin{aligned} \frac{d(4u^3 + 3)}{du} &= \frac{d(4u^3)}{du} + \frac{d(3)}{du} \\ &= \frac{d(4u^3)}{du} + 0 \\ &= 12u^2 + 0 \\ &= 12u^2 \end{aligned} \quad \begin{aligned} \frac{d(au^2 + bu + c)}{du} &= \frac{d(au^2)}{du} + \frac{d(bu)}{du} + \frac{d(c)}{du} \\ &= 2au + b + 0 \\ &= 2au + b \end{aligned}$$

iv) General Power Rule:

$$\begin{aligned} \frac{d(au+b)^n}{du} &= \frac{d(au+b)}{du} \times \frac{d(au+b)^{n-1}}{du} \\ &= n(au+b)^{n-1} \cdot \frac{d(au+b)}{du} \end{aligned}$$

$$\begin{aligned} \text{e.g. } \frac{d(2u+3)^{1/2}}{du} &= \frac{d(2u+3)}{du} \times \frac{d(2u+3)^{1/2}}{du} \\ &= \frac{1}{2} (2u+3)^{-1/2} \times \frac{d(2u)}{du} + \frac{d(3)}{du} \\ &= \frac{1}{2} (2u+3)^{-1/2} \times 2 + 0 \\ &= \frac{1}{\sqrt{2u+3}} \end{aligned}$$

e.g.

$$\begin{aligned}
 \frac{d}{du} (\sqrt{au^2 + bu + c}) &= \frac{d}{du} (au^2 + bu + c)^{\frac{1}{2}} \\
 &= \frac{d}{du} (au^2 + bu + c)^{\frac{1}{2}} \times \frac{d}{du} (au^2 + bu + c) \\
 &= \frac{1}{2} (au^2 + bu + c)^{-\frac{1}{2}} \times \frac{d}{du} (au^2) + \frac{d(bu)}{du} + \frac{d(c)}{du} \\
 &= \frac{1}{2} (2au + b) \times \frac{d}{du} (au^2 + bu + c)
 \end{aligned}$$

v) Product Rule:

$$\frac{d}{du}(u \cdot v) = u \frac{dv}{du} + v \frac{du}{du}$$

$$\text{e.g. } \frac{d}{du} (4u^2 + 8) \cdot (3u^3 + 6u)$$

$$= (4u^2 + 8) \cdot \frac{d}{du} (3u^3 + 6u) + (3u^3 + 6u) \frac{d}{du} (4u^2 + 8)$$

$$= (4u^2 + 8)(9u^2 + 6) + (3u^3 + 6u)(8u)$$

$$= (4u^2 + 8)(9u^2 + 6) + (24u^3 + 48u^2)$$

vi) Quotient Rule:

$$\frac{d}{du}\left(\frac{u}{v}\right) = \frac{v \frac{du}{du} - u \frac{dv}{du}}{v^2}$$

$$\text{e.g. } \frac{d}{du}\left(\frac{3u^2 + 2}{4u + 7}\right) = (4u + 7) \frac{d}{du}(3u^2 + 2) - (3u^2 + 2) \frac{d}{du}(4u + 7)$$

$$= (4u + 7)(6u) - (3u^2 + 2) \cdot 4$$

$$= \frac{24u^2 + 42u - 3u^2 - 8}{(4u + 7)^2}$$

$$= \frac{12u^2 + 42u - 8}{(4u + 7)^2}$$

$$\begin{aligned}
 \text{e.g. } & \frac{d}{du} \left(\sqrt{\frac{1+u}{1-u}} \right) \\
 &= \frac{d \left(\frac{1+u}{1-u} \right)^{\frac{1}{2}}}{du} \times \frac{d \left(\frac{1+u}{1-u} \right)}{du} \\
 &= \frac{1}{2} \left(\frac{1+u}{1-u} \right)^{-\frac{1}{2}} \times (1-u) \frac{d \left(\frac{1+u}{1-u} \right)}{du} - (1+u) \frac{d \left(\frac{1-u}{1-u} \right)}{du} \\
 &= \frac{1}{2} \sqrt{\frac{1+u}{1-u}} \times \frac{(1-u) \cdot 1 - (1+u) \cdot (-1)}{(1-u)^2} \\
 &= \frac{1-u - (-1-u)}{2 \sqrt{\frac{1+u}{1-u}} \cdot (1-u)^2} \\
 &= \frac{1-u+1+u}{2 \sqrt{\frac{1+u}{1-u}} \cdot (1-u)^2} = \frac{2}{2 \sqrt{\frac{1+u}{1-u}} \cdot (1-u)^2} \\
 &= \frac{1}{\sqrt{\frac{1+u}{1-u}} \cdot (1-u)^2}
 \end{aligned}$$

vii) Trigonometric:

- a) $\frac{d(\sin u)}{du} = \cos u$
- b) $\frac{d(\cos u)}{du} = -\sin u$
- c) $\frac{d(\tan u)}{du} = \sec^2 u$
- d) $\frac{d(\sec u)}{du} = \sec u \cdot \tan u$
- e) $\frac{d(\csc u)}{du} = -\csc u \cdot \cot u$
- f) $\frac{d(\cot u)}{du} = -\operatorname{cosec}^2 u$

① e.g. $\frac{d}{du} (\sin^2 u + 8 \cos^2 u + 2)$

$$= \frac{d(\sin^2 u)}{d \sin u} \times \frac{d \sin u}{du} + 8 \left(\frac{d(\cos^2 u)}{d \cos u} \times \frac{d \cos u}{du} \right) + \frac{d(2)}{du}$$

$$= 2 \sin u \cdot \cos u + 8 \cdot 2 \cos u \cdot (-\sin u) + 0$$

$$= 2 \sin u \cdot \cos u - 16 \sin u \cdot \cos u$$

$$= -14 \sin u \cdot \cos u \quad [\because \sin 2u = 2 \sin u \cdot \cos u]$$

$$= -7 \sin 2u \quad \#.$$

② e.g. $\frac{d}{du} (\tan^2 u + \sec u)$

$$= \frac{d(\tan^2 u)}{d \tan u} \times \frac{d(\tan u)}{du} + \frac{d \sec u}{du}$$

$$= 2 \tan u \cdot \sec^2 u + \sec u \cdot \tan u$$

$$= \tan u \cdot \sec u (2 \sec u + 1) \quad \#$$

③ e.g. $\frac{d}{du} \sqrt{\sin^3 u + 2 \sin^2 u}$

$$= \frac{d}{du} (\sin^3 u + 2 \sin^2 u)^{\frac{1}{2}}$$

$$= \frac{d(\sin^3 u + 2 \sin^2 u)^{\frac{1}{2}}}{d(\sin^3 u + 2 \sin^2 u)} \times \frac{d(\sin^3 u + 2 \sin^2 u)}{du}$$

$$= \frac{1}{2} (\sin^3 u + 2 \sin^2 u)^{-\frac{1}{2}} \times \frac{d \sin^3 u}{ds \sin u} \times \frac{ds \sin u}{du} + 2 \left[\frac{d \sin^2 u}{ds \sin u} \times \frac{ds \sin u}{du} \right]$$

$$= \frac{1}{2 \sqrt{\sin^3 u + 2 \sin^2 u}} \times 3 \sin^2 u \cdot \cos u + 2 \cdot 2 \sin u \cdot \cos u$$

$$= \frac{\sin u \cdot \cos u (3 \sin u + 4)}{2 \sqrt{\sin^3 u + 2 \sin^2 u}}$$

$$= \frac{\sin u \cdot \cos u \cdot (3 \sin u + 4)}{2 \sqrt{\sin^3 u + 2 \sin^2 u}} \quad \#$$

$$\textcircled{2} \text{ e.g. } \frac{d}{du} \left(\frac{\sin 4u}{\tan 6u} \right)$$

$$= \tan 6u \frac{d(\sin 4u)}{du} - \sin 4u \frac{d(\tan 6u)}{du}$$

$$= \tan 6u \frac{d \sin 4u}{d \cancel{4u}} \times \cancel{4u} - \sin 4u \cdot \frac{d \tan 6u}{d 6u} \times \frac{d 6u}{\tan^2 6u}$$

$$= \tan 6u \cdot \cos 4u \cdot 4 - \sin 4u \cdot \sec^2 6u \cdot 6 / \tan^2 6u$$

$$= \frac{4 \tan 6u \cdot \cos 4u - 6 \sin 4u \cdot \sec^2 6u}{\tan^2 6u}$$

viii) Logarithmic function:

$$\frac{d(\log u)}{du} = \frac{1}{u}$$

$$\textcircled{1} \text{ e.g. } \frac{d(\log u^2)}{du^2} \times \frac{d(u^2)}{du} \quad \textcircled{2} \text{ e.g. } \frac{d(u + 4 \log 4u)}{du}$$

$$= \frac{1}{u^2} \cdot 2u$$

$$= \frac{2}{u} \#$$

$$= \frac{du}{du} + 4 \frac{d \log 4u}{d(4u)} \times \frac{d(4u)}{du}$$

$$= 1 + 4 \times \frac{1}{4u} \cdot 4$$

$$= 1 + \frac{4}{u} \#$$

$$\textcircled{3} \text{ e.g. } \frac{d(\sin u + \log(\tan u))}{du}$$

$$= \frac{d(\sin u)}{du} + \frac{d \log(\tan u)}{d \tan u} \times \frac{d \tan u}{du}$$

$$= \cos u + \frac{1}{\tan u} \times \sec^2 u$$

$$= \cos u + \frac{\sec^2 u}{\tan u} \#$$

④ e.g. $\frac{d(\tan u + \log \sqrt{\sin u})}{du}$

$$= \frac{d(\tan u)}{du} + \frac{d \log (\sin u)^{\frac{1}{2}}}{d(\sin u)^{\frac{1}{2}}} \times \frac{d(\sin u)^{\frac{1}{2}}}{d \sin u} \times \frac{d \sin u}{du}$$

$$= \sec^2 u + \frac{1}{\sqrt{\sin u}} \times \frac{1}{2} \sin^{-\frac{1}{2}} u \cdot \cos u$$

$$= \sec^2 u + \frac{1}{\sqrt{\sin u}} \cdot \frac{\cos u}{2 \sqrt{\sin u}}$$

$$= \sec^2 u + \frac{\cot u}{2} \#.$$

⑤ e.g. $\frac{d(\cot u + \log \sqrt{\cos u})}{du}$

$$= \frac{d(\cot u)}{du} + \frac{d \log \sqrt{\cos u}}{d \sqrt{\cos u}} \times \frac{d \sqrt{\cos u}}{d \cos u} \times \frac{d \cos u}{du}$$

$$= -\operatorname{cosec}^2 u + \frac{1}{\sqrt{\cos u}} \cdot \frac{1}{2} \cos^{-\frac{1}{2}} u \cdot -\sin u$$

$$= -\operatorname{cosec}^2 u + \frac{(-) \sin u}{2 \cos u}$$

$$= -\operatorname{cosec}^2 u - \frac{\tan u}{2} \#.$$

ix.) Exponential function :-

$$\frac{d(e^u)}{du} = e^u$$

e.g.

$$\textcircled{1} \frac{d(e^{4u})}{du} = \frac{d(e^{4u})}{d(4u)} \times \frac{d(4u)}{du} = e^{4u} \cdot 4 = 4e^{4u} \#.$$

$$\textcircled{2} \frac{d(e^{\sin u + \tan u})}{du} = \frac{d(e^{\sin u + \tan u})}{d(\sin u + \tan u)} \times \frac{d(\sin u + \tan u)}{du}$$

$$= e^{\sin u + \tan u} \cdot (\cos u + \sec^2 u) \#.$$

$$\textcircled{3} \frac{d(e^{\frac{1}{u}})}{du} = \frac{d(e^{\frac{1}{u}})}{d(\frac{1}{u})} \times \frac{d\frac{1}{u}}{du} = e^{\frac{1}{u}} (-1) u^{-2} = -e^{\frac{1}{u}} \frac{1}{u^2} \#.$$

$$\textcircled{4} \frac{d(e^{u^2+4u})}{du} = \frac{d(e^{u^2+4u})}{d(u^2+4u)} \times \frac{d(u^2+4u)}{du} \\ = e^{u^2+4u} \cdot (2u+4)$$

$$\textcircled{5} \frac{d(e^{\sqrt{\cos u}})}{du} = \frac{d(e^{\sqrt{\cos u}})}{d(\cos u)^{\frac{1}{2}}} \times \frac{d(\cos u)^{\frac{1}{2}}}{d \cos u} \times \frac{d \cos u}{du} \\ = e^{\sqrt{\cos u}} \cdot \frac{1}{2} \cos^{-\frac{1}{2}} u \cdot (-\sin u) \\ = -\frac{1}{2} e^{\sqrt{\cos u}} \cos^{-\frac{1}{2}} u (-\sin u) \\ = e^{\sqrt{\cos u}} \cdot \frac{1}{2} \cos^{-\frac{1}{2}} u (-\sin u) \#.$$

x) Implicit Differentiation :-

Find $\frac{dy}{du}$ in, ① eqⁿ of circle, $u^2+y^2=4$

② eqⁿ of ellipse, $\frac{u^2}{9} + \frac{y^2}{16} = 1$

③ eqⁿ of Parabola, ~~Parabola~~ $y^2=16u$

④ Eqⁿ of circle,

$$u^2+y^2=4$$

differentiating both sides w.r.t to u

$$\frac{d(u^2+y^2)}{du} = \frac{d(4)}{du}$$

$$\text{or, } \frac{d(u^2)}{du} + \frac{d(y^2)}{du} = 0$$

$$\text{or, } -2u = 2y \frac{dy}{du}$$

$$\text{or, } 2u + \frac{dy^2}{dy} \times \frac{dy}{du} = 0$$

$$\therefore \frac{dy}{du} = -\frac{u}{y} \#$$

$$\text{or, } 2u + 2y \times \frac{dy}{du} = 0$$

⑥ eqⁿ of ellipse,

$$\frac{u^2}{9} + \frac{y^2}{16} = 1$$

differentiating both sides w.r.t u.

$$\frac{d\left(\frac{u^2}{9} + \frac{y^2}{16}\right)}{du} = \frac{d(1)}{du}$$

$$\text{or, } \frac{d(9u^2 + 16y^2)}{du} = \frac{d(9 \times 16)}{du}$$

$$\text{or, } 16\frac{du^2}{du} + 9\frac{dy^2}{dy} \times \frac{dy}{du} = 0$$

$$\text{or, } 16 \cdot 2u + 9 \cdot 2y \frac{dy}{du} = 0$$

$$\text{or, } 18y \cdot \frac{dy}{du} = -32u$$

$$\therefore \frac{dy}{du} = -\frac{16u}{9y} \quad \#.$$

⑦ eqⁿ of Parabola,

~~$$y^2 = 16u$$~~

differentiating both sides w.r.t u.

$$\frac{d(y^2)}{du} = \frac{d(16u)}{du}$$



~~$$2y \cdot \frac{dy}{du} = 16 \cdot 1$$~~

$$\text{or, } \frac{dy^2}{dy} \times \frac{dy}{du} = 16 \frac{du}{du}$$

$$\text{or, } \frac{dy}{du} = \frac{16}{2y}$$

$$\therefore \frac{dy}{du} = \frac{8}{y} \quad \#.$$

Maxima And Minima :-



Rise is denoted by \nearrow and fall is denoted by \searrow

i) Stationary (critical point)

$f'(u) = 0$ and solve for u

$u = a, b, c$ (Stationary point)

ii) Point of inflection :

$f''(u) = 0$ and solve for u

check for maxima and minima

if $u=0$, $f''(a) > 0$ (minima)

$f''(a) < 0$ (maxima)

$f''(a) = 0$ (neither max nor min)

Graph Sketching :

Step I :- Find stationary / critical point and point of inflection.

Step II :- Sketch Rise (\nearrow) and fall (\searrow) pattern using $f'(u)$.

if +ve rise if -ve fall

For more notes visit <https://collegenote.pythonanywhere.com>

Step III: Sketch concavity up and down using $f''(u)$.
if +ve = concave up, if -ve, concave down

Step IV: Combine Step 2 and Step 3 (figure) gives general shape of $f(u)$

Step V: find point of $f(u)$

x			
y			

and plot in the graph.

Example 1:

Graph the function: $y = 4u^3 - u^4$

→ Solution,

$$f'(u) = 12u^2 - 4u^3$$

$$f''(u) = 24u - 12u^2$$

Step I: Stationary point, and Point of inflection

$$f'(u) = 0$$

$$12u^2 - 4u^3 = 0$$

$$4u^2(3-u) = 0$$

$$\text{either } u=3$$

$$u=0$$

$$f''(u) = 0$$

$$24u - 12u^2 = 0$$

$$12u(2-u) = 0$$

$$\text{either, } u=2$$

$$u=0$$

Step II: Rise and fall pattern using $f'(u)$

$12u^2 - 4u^3$	+	+	-
	$\xrightarrow{\text{rise}} 0 \xrightarrow{\text{rise}} 3 \xrightarrow{\text{fall}}$		

to check +ve, -ve. Put value in front of 0 in $f'(u)$

$$\text{sup. } f'(-1) = 12(-1)^2 - 4(-1)^3 = 12 + 4 = 16 \text{ (+ve), so +}$$

$$\text{sup. } f'(4) = 12(4)^2 - 4(4)^3 = 192 - 256 = -64 \therefore -\text{ve}$$


if +ve, rise \rightarrow

if -ve, fall \rightarrow


Step III: concavity using $f''(u)$ (same as Step II to check +ve, -ve)

$24u - 12u^2$	-	+	-
	Concave down	Concave up	Concave down

Step IV: Combine Step II and Step III.



general shape



Step V: find points

of $f(u)$ and plot. $y = 4u^3 - u^4$ when $u=1$

$$y = 4 \cdot 1^3 - 1^4 \\ = 3$$

$$u=2 \\ y=16$$

$$u=3 \\ y=27$$

$$u=-1 \\ y=-5$$

$$u=4 \\ y=0$$

x	1	2	3	-1	4
y	3	16	27	-5	0

(in graph)

example: 2.

$$y = 1 - (u+1)^3$$

→ solution,

$$f'(u) = -3(u+1)^2$$

$$f''(u) = -6(u+1)$$

Step I: Stationary point,

$$f'(u) = 0$$

$$-3(u+1)^2 = 0$$

$$(u+1) = 0$$

$$\therefore u = -1$$

Point of inflection,

$$f''(u) = 0$$

$$-6(u+1) = 0$$

$$u+1 = 0$$

$$\therefore u = -1$$

$$\frac{d(u)}{du} = \frac{d(u+1)^3}{du} \times \frac{d(u+1)}{du} \\ = 0 - 3(u+1)^2 \\ = -3(u+1)^2$$

$$-3 \frac{d(u+1)^2}{d(u+1)} \times \frac{d(u+1)}{du} \\ = -3 \cdot 2(u+1) \\ = -6(u+1)$$

Step 2: Rise and fall pattern:

Put 0 in $f'(u)$

$$-3(0+1)^2 = -3$$

$$-3(u+1)^2 \quad - \quad + \quad -$$

↓ ↓ ↓

Put -2 in $f'(u)$

$$= -3(-2+1)^2 = -3(-1)^2 = -3$$

Step 3: Concavity

$$-6(u+1) \quad + \quad -$$

↓ ↓ ↓

Concave up -1 Concave down

Put -2 in $f''(u)$

$$= -6(-2+1) = -6(-1) = +6$$

Step 4: Combine Step 2 and 3

Put 0 in $f''(u)$

$$= -6(0+1) = -6$$

rise/fall	fall	↓	fall	↓
up/down		↓		↓

general form

Step 5: Plot the in graph.

find point $f(u)$ $y = 1 - (u+1)^3$ when $u = -1$

$$y = 1 - (-1+1)^3$$

$$y = 1 - 0 = 1$$

when $u = 0$

$$y = 1 - (0+1)^3$$

$$y = 1 - 1 = 0$$

when $u = 1$

$$y = 1 - (1+1)^3$$

$$y = 1 - 8 = -7$$

when $u = 2$

$$y = 1 - (2+1)^3$$

$$y = 1 - 27 = -26$$

X	-1	0	1	2	-2	-3
Y	1	0	-1	-8	2	9

In graph

Example 33

$$y = u^4 + 2u^3$$

solution, $f'(u) = 4u^3 + 6u^2$
 $f''(u) = 12u^2 + 12u$

Step 1: Stationary point Point of inflection,

$$f'(u) = 0 \quad f''(u) = 0$$

$$4u^3 + 6u^2 = 0 \quad 12u^2 + 12u = 0$$

$$\therefore 2u^2(2u+3) = 0 \quad 12u(u+1) = 0$$

either $u = 0$ either $u = 0$

$$u = -\frac{3}{2} = -1.5 \quad u = -1$$

Step 2:

Rise and fall pattern from $f'(u)$

$$\begin{array}{c|ccccc} 4u^3 + 6u^2 & - & + & + & + \\ \hline & \rightarrow -1.5 & \rightarrow 0 & \rightarrow & \\ \end{array} = 4(-2)^3 + 6(-2)^2 \\ = 4(-8) + 6 \cdot 4 \\ = -32 + 24 = -8$$

Step 3:

Concavity using $f''(u)$

$$\begin{array}{c|ccccc} 12u^2 + 12u & + & - & + & + \\ \hline & up & -1 & down & 0 & up \\ \end{array} = 12(-2)(-2+1) \\ = -24 \times (-1) \\ = +24$$

Step 4: Combining Step 2 and 3

$$\begin{array}{c|ccccc} \text{rise/fall} & \rightarrow & \rightarrow & \rightarrow & \rightarrow \\ \hline & -1.5 & -1 & 0 & \\ \text{up/down} & \swarrow & \uparrow & \swarrow & \uparrow \\ \end{array} = -6 \times 0.5 \\ = -3$$

Step 5: Find the point $f(u)$ and plot the graph. $y = u^4 + 2u^3$

x	0	1	-1	-2	+2
y	0	3	-1	0	32

In graph

Assignment #1

Graph the function:

Ex. 1 ④ $y = -u^3 + 12u + 5$

→ Solution,

$f'(u) = -3u^2 + 12$, $f''(u) = -6u$

1. Stationary point

$f'(u) = 0$

$-3u^2 + 12 = 0$

$3(-u^2 + 4) = 0$

or, $-u^2 + 4 = 0$

or, $u^2 = 4$

$\therefore u = \pm 2$

Point of inflection

$f''(u) = 0$

$-6u = 0$

$\therefore u = 0$


2. Rise and fall pattern:

$-3u^2 + 12$	-	+	-
	-2		+2


3. Concavity

$-6u$	+	-
	up	down

4. Combine 2 and 3



general shape:

5. find point x and y in $f(u)$ and plot in graph

x	-1	0	-2	2	-3
y	-6	5	-11	21	-9

In graph copy

Ex. 5

① $y = u^3 - 3u + 3$
 $f'(u) = 3u^2 - 3$
 $f''(u) = 6u$

1. Stationary point.

$$f'(u) = 0$$

$$3u^2 - 3 = 0$$

$$3(u^2 - 1) = 0$$

$$u = \sqrt{1}$$

$$\therefore u = \pm 1$$

Point of inflection.

$$f''(u) = 0$$

$$6u = 0$$

$$\therefore u = 0$$

2. Rise and fall pattern.

$$3u^2 - 3$$

	+		-		+
	↗ -1		↘ +1		↗

3. concavity


$$6u$$

	-		+
	concave down	0	up

4. Combining 2 and 3

rise/fall	↗	↓	↗	↓	↗
up/down	↗ -1	↘ 0	↗ +1	↙	

general shape

5. find point x and y
and plot in graph.

x	0	1	-1	2	-2
y	3	1	5	5	1

In graph

Q. NO: 6 $y = u(6-2u^2)$

→ Solution,

$$f'(u) = 6u - 2u^3$$

$$f'(u) = 6 - 6u^2$$

$$f''(u) = -12u$$

Step 1: Stationary point is given by $f'(u) = 0$

$$6 - 6u^2 = 0$$

$$\therefore 6(1 - u^2) = 0$$

$$\therefore u^2 = 1$$

$$\therefore u = \pm 1$$

Point of inflection is given by $f''(u) = 0$

$$-12u = 0$$

$$\therefore u = 0$$

Step 2: Rise and fall pattern:


(Q.P.Q.P.Q.P)	(+)	(-)	(+)	(-)
	(-)	(+)	(-)	(+)

$6 - 6u^2$	-	+	-	-
	→	-1	→	1


Step 3: Concavity

$-12u$	+	-
	Conc. up	Conc. down

Step 4: Combine Step 2 and 3.



general shape



Step 5: find the value of x and y and plot in graph.

X	1	2	-1	-2	0
Y	4	-4	-4	+4	0

Q. NO: 7 $y = 1 - 9u - 6u^2 - u^3$

→ solution,

$$f'(u) = -9 - 12u - 3u^2$$

$$f''(u) = -12 - 6u$$

Step 1: Stationary point from $f'(u) = 0$ critical point from $f''(u) = 0$

$$f'(u) = 0$$

$$f''(u) = 0$$

$$-9 - 12u - 3u^2 = 0$$

$$-12 - 6u = 0$$

$$3u^2 + 12u = -9$$

$$6(-2 - u) = 0$$

$$u^2 + 4u = -3$$

$$u = -2$$

$$u^2 + 4u + 3 = 0$$

$$u + u + 3u + 3 = 0$$

$$-2(u+1) + 3(u+1) = 0$$

$$\text{either } u = -1 \\ u = -3$$


Step 2: Rise and fall pattern from $f'(u)$

$$\begin{array}{c|ccc} -9 - 12u - 3u^2 & -ve & +ve & +ve \\ \hline & \nearrow -3 & \nearrow -1 & \nearrow \end{array}$$

Step 3: Concavity from $f''(u)$

$$\begin{array}{c|cc} -12 - 6u & +ve & -ve \\ \hline & \downarrow & \\ \text{Conc. up} & -2 & \text{Conc. down} \end{array}$$

Step 4: combine Step 2 and 3.



Step 5: find X and Y → plot in graph

X	0	-1	-2	-3	-4	-1
Y	2	5	3	1	5	-15

$$8. y = 1 - (u+1)^3$$

$$f'(u) = -3(u+1)^2$$

$$f''(u) = -6(u+1)$$

Step 1: Point of stationary

$$f'(u) = 0$$

$$-3(u+1)^2 = 0$$

$$(u+1) = 0$$

$$u = -1$$

Point of inflection

$$f''(u) = 0$$

$$-6(u+1) = 0$$

$$\Rightarrow u+1 = 0$$

$$\therefore u = -1$$

Step 2: Rise and fall pattern


$$\begin{array}{c|ccc} -3(u+1)^2 & - & + & - \\ \hline & \rightarrow & -1 & \rightarrow \end{array}$$

Step 3: Concavity

$$\begin{array}{c|cc} -6(u+1) & + & - \\ \hline & \text{up} & -1 \text{ down} \end{array}$$

Step 4:

Combine Step 2 and 3



Step 5:

find x and y \rightarrow plot in graph $y = 1 - (u+1)^3$

x	1	0	-1	-2	-3
y	-7	0	1	2	3

$$y = (u-2)^3 + 1$$

→ Solution,

$$f'(u) = 3(u-2)^2$$

$$f''(u) = 6(u-2)$$

Step 1: Stationary Point

$$f'(u) = 0$$

$$3(u-2)^2 = 0$$

$$(u-2) = 0$$

$$\therefore u = 2$$

$$\frac{d(u-2)^3}{d(u-2)} \times \frac{d(u-2)}{d(u-2)}$$

$$= 3(u-2)^2 \times 1$$

$$= 3 \times 2 \frac{d(u-2)^2}{d(u-2)} + \frac{d(u-2)}{d(u-2)}$$

Point of inflection

$$f''(u) = 0$$

$$6(u-2) = 0$$

$$u = 2$$

$$6(u-2) \neq 0$$

Step 2: Rise and Fall pattern

$$3(u-2)^2 \quad + \quad \frac{1}{2} \quad +$$

Step 3: Concavity

$$6(u-2) \quad - \quad 1 \quad +$$

down

Step 4: Combining Step 2 and 3

$$\begin{array}{c|c|c|c} \text{rise/fall} & \nearrow & | & \nearrow \\ \text{up/down} & \curvearrowleft & 2 & \curvearrowright \end{array}$$

general form

Step 5: find x and y and plot in graph

x	1	2	0	3	4
y	0	1	-7	2	9

$$10. y = -u^4 + 6u^2 - 4$$

→ Solution,

$$f'(u) = -4u^3 + 12u$$

$$f''(u) = -12u^2 + 12$$

Step 1: Stationary point, $f'(u) = 0$

$$-4u^3 + 12u = 0$$

$$4u(-u^2 + 3) = 0$$

$$-u^2 + 3 = 0$$

$$u = \sqrt{3} \approx 1.73$$

Point of inflection,

$$-12u^2 + 12 = 0$$

$$12(-u^2 + 1) = 0$$

$$-u^2 + 1 = 0$$

$$u = \pm 1$$

Step 2: Rise and fall pattern,

$-4u^3 + 12u$	+	1	-
	→	1.73	→

Step 3: Concavity

$-12u^2 + 12$	+	1	-
	up	1	down

Step 4: Combine ② and ③

rise/fall	↑	1	↑
up/down	↓	1	↓

general form

Step 5: Find x and y and plot in graph

x	1	-1	0	2
y	1	1	-4	5

Now
08/15

New Link

Date _____

Page _____

Graph the function.

11. $y = u^{5/3} - 5u^{2/3}$

→ Solution,

$$f'(u) = \frac{5}{3}u^{5/3-1} - 5 \times \frac{2}{3}u^{2/3-1}$$

$$= \frac{5}{3}u^{2/3} - \frac{10}{3}u^{-1/3}$$

$$f''(u) = \frac{5}{3} \times \frac{2}{3}u^{2/3-1} - \frac{10}{3} \times \left(-\frac{1}{3}\right)u^{-1/3-1}$$

$$= \frac{10}{9}u^{-1/3} + \frac{10}{9}u^{-4/3}$$

Stationary Point is given by $f'(u)=0$

$$\frac{5}{3}u^{2/3} - \frac{10}{3}u^{-1/3} = 0$$

$$\frac{5}{3}u^{-1/3}(u-2) = 0$$

$$\therefore x = 0$$

$$\therefore x = +2$$

Point of inflection is given by $f''(u)=0$

$$\frac{10}{9}u^{-1/3} + \frac{10}{9}u^{-4/3} = 0$$

$$\frac{10}{9}u^{-4/3}(u+1) = 0$$

Either $u=0$ $u=-1$

Rise and fall pattern


$f'(u)$	+	1	-	1	+	1
	rise	0	fall	2	rise	



Step 3: sketch concave up and concave down using $f''(u)$

$f''(u)$	-	+	-	+
	down	-1	up	0 up

Step 4: combine Step 2 and 3.




general form,

plot in graph

x	x
0	1
-4	2
-4.7	3
-4.16	4
-2.5	5
0	6
3.3	7
7.3	8
12	

general shape



Integration

$$1. \int u^n du = \frac{u^{n+1}}{n+1} + C \quad (C = \text{constant of integration})$$

e.g. @ $\int (u^{\frac{2}{3}} + u) du$

$$= \int u^{\frac{2}{3}} du + \int u du$$

$$= \frac{u^{\frac{2}{3}+1}}{\frac{2}{3}+1} + \frac{u^2}{2} + C$$

$$= \frac{u^{\frac{5}{3}}}{\frac{5}{3}} + \frac{u^2}{2} + C \quad \#.$$

$$2. \int du = u + C$$

$$3. \int (2u+b)^n du = \frac{(2u+b)^{n+1}}{(n+1)2} + C$$

e.g. @ $\int (4u+3)^5 du = \frac{(4u+3)^6}{6 \times 4} + C \quad \#.$

$$\textcircled{b} \int (4-6u)^{\frac{2}{3}} du = \frac{(4-6u)^{\frac{2}{3}+1}}{(\frac{2}{3}+1) \times (-6)} + C = \frac{(4-6u)^{\frac{5}{3}}}{-30} + C$$

$$= -\frac{(4-6u)^{\frac{5}{3}}}{10} + C \quad \#.$$

$$4. \int \frac{1}{u} du = \log u + C$$

e.g. @ $\int (au^2 + bu + u^{-1}) du$ (log: ln) $\int \frac{1}{(1-u)} du$

$$= au^3 + bu^2 + \log u + C \quad - \log(1-u) + C$$

$$\textcircled{b} \quad \int \frac{1-u}{1+u} du$$

$$= \cancel{\int \frac{1-u}{1+u} du}$$

$$= \int \frac{(1+u)-2u}{(1+u)} du$$

$$= \int \frac{(1+u)}{(1+u)} du - \int \frac{2u}{(1+u)} du$$

$$= \int du - 2 \int \left(\frac{u+1-1}{u+1} \right) du$$

$$= u - 2 \left\{ \int \frac{u+1}{u+1} du - \int \frac{1-1}{u+1} du \right\}$$

$$= u - 2u - 2 \log(u+1) + C \quad \# \quad = -u - 2 \log(1+u) + C \quad \#$$

$$\textcircled{c} \quad \int \frac{(u+3)}{(u-3)} du$$

$$= \int \frac{(u-3)+6}{(u-3)} du$$

$$= \int \left(\frac{(u-3)}{(u-3)} + \frac{6}{(u-3)} \right) du = \int 1 du + 6 \int \frac{1}{(u-3)} du$$

$$= \cancel{6u} + 6 \log(u-3) + C$$

(b) \rightarrow OR

$$\int \frac{1}{1+u} - \int \frac{u+1-1}{1+u}$$

$$\log(u+1) - \frac{u+1}{u+1} + \frac{1}{1+u}$$

$$5) \int e^{au} du = \frac{e^{au}}{a} + C$$

E.g.

$$a) \int e^{5u} du = \frac{e^{5u}}{5} + C \#.$$

$$b) \int e^{-4/3u} du = \frac{e^{-4/3u}}{-4/3} + C \#.$$

$$c) \int (e^{-pu} + e^{qu}) du = \frac{e^{-pu}}{-p} + \frac{e^{qu}}{q} + C \#.$$

6. Trigonometric

$$i) \int \cos au du = \frac{\sin au}{a} + C \#$$

$$ii) \int \sin au du = -\frac{\cos au}{a} + C \#.$$

$$iii) \int \sec^2 au du = \frac{\tan au}{a} + C \#.$$

$$iv) \int \sec au \times \tan au du = \frac{\sec au}{a} + C \#$$

$$v) \int \cosec^2 au du = -\frac{\cot au}{a} + C \#.$$

$$vi) \int \cosec au \cdot \cot au du = -\frac{\cosec au}{a} + C \#.$$

Trigonometric Substitution:

$$i) a^2 - u^2,$$

$$\text{put } u = a \sin \theta$$

② $u^2 + a^2$,

put $u = a \tan \theta$

$$\begin{aligned} &= a^2 (1 + \tan^2 \theta) \\ &= a^2 \sec^2 \theta \end{aligned}$$

③ $u^2 - a^2$

put $u = a \sec \theta$

7. Definite Integrals:

$$\int_{u=a}^{u=b} u du = \left[\frac{u^2}{2} \right]_a^b$$

$$= \frac{b^2}{2} - \frac{a^2}{2} = \frac{b^2 - a^2}{2}$$

e.g. $u = \frac{\pi}{2}$

④ $\int_0^{\pi/2} 6 \sin u du = [-\cos u]_0^{\pi/2}$

$$= [-\cos \frac{\pi}{2} + \cos 0] = [0 + 1] = 1$$

⑤ $\int_0^3 \sqrt{9-u^2} du$

Put $u = 3 \sin \theta$ since $\therefore 2 = 3$

$u = 3 \sin \theta$

differentiating both sides by θ .

① Q.E.D.

$\frac{du}{d\theta} = 3 \cos \theta$

$du = 3 \cos \theta d\theta$

$x = 3 \sin \theta$ when $x = 3$

when $x = 0$

$0 = 3 \sin \theta$

$\sin \theta = 0$

$\therefore \theta = 0$

$$\theta = \frac{\pi}{2}$$

$$= \int \sqrt{g - g \sin^2 \theta} \cdot 3 \cos \theta d\theta \quad \cos^2 \theta = \frac{1 + \cos 2\theta}{2}$$

$$\theta = 0^\circ \quad \sqrt{g(1 - \sin^2 \theta)}$$

$$= \sqrt{g} \cos^2 \theta$$

$$= \int_0^{\frac{\pi}{2}} 3 \cos \theta \cdot \sqrt{g} \cos^2 \theta d\theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

$$\sin 2\theta = 2 \sin \theta \cdot \cos \theta$$

$$= \int_0^{\frac{\pi}{2}} 9 \cos^2 \theta d\theta$$

$$\omega s^2 \theta = \cos^2 \theta - (1 - \cos^2 \theta)$$

$$\cos 2\theta = \cos^2 \theta - 1 + \omega s^2 \theta$$

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$$

$$\sin 2\theta = \frac{1 - \cos 2\theta}{2}$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

$$= \omega s^2 \theta - 1 + \cos^2 \theta$$

$$\omega s^2 \theta = 2 \cos^2 \theta - 1$$

$$\text{Ans} = \frac{1 + \omega s^2 \theta}{2}$$

$$= \frac{9}{2} \int_0^{\frac{\pi}{2}} (1 + \omega s^2 \theta) d\theta$$

↓
Integrate

$$= \frac{9}{2} \left[\theta + \frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{2}}$$

$$= \frac{9}{2} \left[\frac{\pi}{2} + \sin 2 \cdot \frac{\pi}{2} \cdot \frac{1}{2} - 0 \right] \quad \sin \frac{\pi}{2} = 0$$

$$= \frac{9\pi}{4} \times \frac{1}{2}$$

$$= \frac{9\pi}{8}$$


Ans

How Assignment

2073/08/30, Thu.


New Link
Date _____
Page _____

8. Area




$$\text{Area } PQRS = \int_{x=a}^{x=b} y du$$

$$= \int_{x=a}^{x=b} f(u) du$$




$$\text{Area of } PQRS = \int_{x=a}^{x=b} y du = \int_{x=a}^{x=b} f(u) du$$



$$\text{Area of } PQRS = \int_{y=a}^{y=b} u dy = \int_{y=a}^{y=b} f(y) dy$$

Area between two curves




$$= \text{Area PQRS} - \text{Area TURS}$$

$$= \int_{u=0}^{u=b} y_2 du - \int_{u=0}^{u=b} y_1 du$$

$$= \int_{x=0}^{x=b} (y_2 - y_1) dx$$

Q. Find the area of $y^2 = 4ax$ from $u=0$ to $u=4$



$$u=4$$

$$= 2\sqrt{a} \int u^{1/2} du$$

$$u=0$$

$$= 2\sqrt{a} \left[\frac{u^{1/2+1}}{\frac{1}{2}+1} \right]_0^4$$

$$= 2\sqrt{a} \left[\frac{2}{3} u^{3/2} \right]_0^4$$

$$= 2\sqrt{a} \left[\frac{2}{3} \times (4)^{3/2} - 0 \right]$$

$$= 2\sqrt{a} \left[\frac{2}{3} \times (2)^2 \right]$$

$$= 2\sqrt{a} \frac{2 \times 8}{3}$$

$$= \frac{32\sqrt{a}}{3}$$

$$\therefore \text{Area of Parabola} = 2 \times \frac{32\sqrt{a}}{3}$$

$$= \frac{64\sqrt{a}}{3} \text{ Sq. units}$$

Area of circle :-

$$u^2 + y^2 = 16 \quad r=4$$

$$x=b$$


$$\Rightarrow \int y du$$

$$x=a$$

$$u=4$$

$$\Rightarrow \int \sqrt{16-u^2} du$$

$$u=0$$



Now,

Put $u = 4 \sin \alpha \dots \text{--- (1)}$ when $u=0$

$$4 \sin \alpha = 0$$

$$\therefore \sin \alpha = 0$$

$$\therefore \sin \alpha = \sin 0^\circ$$

$$\therefore \alpha = 0^\circ$$

diff. eqn (1) w.r.t. α

$$\frac{du}{d\alpha} = \frac{4 \sin \alpha}{d\alpha}$$

$$du = 4 \cos \alpha d\alpha$$

Again,

$$u = \cancel{4}$$

$$\Rightarrow \int \sqrt{16 - u^2} du$$

$$u = 0$$

$$0 \quad \frac{\pi}{2}$$

$$\Rightarrow \int_{0}^{\frac{\pi}{2}} \sqrt{16 - 16 \sin^2 \alpha} \cdot 4 \cos \alpha d\alpha$$

$$0^\circ \quad \sqrt{16(1 - \sin^2 \alpha)} = 4 \sqrt{\cos^2 \alpha} = 4 \cos \alpha$$

$$\Rightarrow \int_{0}^{\frac{\pi}{2}} 4 \cos \alpha \cdot 4 \cos \alpha d\alpha$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha$$

$$\cos 2\alpha = \cos^2 \alpha - (1 - \cos^2 \alpha)$$

$$\therefore \cos^2 \alpha = \frac{\cos 2\alpha + 1}{2}$$

$$\Rightarrow \int_{0}^{\frac{\pi}{2}} 16 \cos^2 \alpha d\alpha$$

$$\Rightarrow \int_{0}^{\frac{\pi}{2}} 16 \cdot \frac{\cos 2\alpha + 1}{2}$$

$$\begin{aligned}
 & \int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^{\cos 2\theta} \int_0^{\sqrt{16-4\theta^2}} d\theta d\phi d\alpha \\
 &= \frac{16}{2} \left\{ \left[\frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{2}} + \frac{\pi}{2} \right\} \\
 &= 8 \left\{ \left[\frac{\sin 2 \cdot \frac{\pi}{2}}{2} - 0 \right] + \frac{\pi}{2} \right\} \\
 &= 8 \left\{ \left[\sin \frac{\pi}{2} \right] + \frac{\pi}{2} \right\} \\
 &= 8 \left\{ 1 + \frac{\pi}{2} \right\} \\
 &= \frac{8\pi}{2} = 4\pi
 \end{aligned}$$

$$\frac{u^2}{16} + \frac{y^2}{9} = 1$$

$$\frac{y^2}{9} = 1 - \frac{u^2}{16}$$


$$\frac{y^2}{9} = \frac{16 - u^2}{16}$$

$$y = \sqrt{\frac{9}{16} (16 - u^2)}$$

$$= 16\pi \# \cdot \frac{3}{4} \sqrt{16 - u^2}$$

$$\therefore \text{Area of circle: } 4\pi \times 4 = 16\pi$$

Q. Find the area of the ellipse $\frac{u^2}{16} + \frac{y^2}{9} = 1$ u is ASW



→ Solution :

given equation of ellipse

$$\textcircled{1} \quad \frac{u^2}{16} + \frac{y^2}{9} = 1$$

$$\frac{y^2}{9} = 1 - \frac{u^2}{16}$$

$$\textcircled{2} \quad \frac{y^2}{9} = \frac{16 - u^2}{16}$$

$$\textcircled{3} \quad y = \sqrt{\frac{9}{16} (16 - u^2)}$$

$$\therefore y = \frac{3}{4} \sqrt{16 - u^2}$$

Now,

Let $x = a \sin \theta$

diff. w.r.t. $d\theta$

$$\frac{du}{d\theta} = a \cos \theta$$

when $u = 0$

$$0 = a \sin \theta$$

$$\therefore \sin \theta = \sin 0^\circ$$

$$\therefore 0 = 0^\circ$$

when $u = \frac{a}{2}$

$$\frac{a}{2} = a \sin \theta$$

$$\therefore \sin \theta = \sin \frac{\pi}{2}$$

$$\theta = \frac{\pi}{2}$$

so, $du = a \cos \theta d\theta$

we have,

$$= \int_0^{\frac{\pi}{2}} \frac{3}{4} \sqrt{16 - 16 \sin^2 \theta} a \cos \theta d\theta$$

$$= \frac{3}{4} \int_0^{\frac{\pi}{2}} \sqrt{16(1 - \sin^2 \theta)} d\theta$$

$$= \frac{3}{4} \int_0^{\frac{\pi}{2}} 4 \cos \theta \cdot 4 \cos \theta d\theta$$

$$= \frac{3}{4} \int_0^{\frac{\pi}{2}} 16 \cos^2 \theta d\theta$$

$$= \frac{3 \times 16}{4} \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta$$

$$= 12 \int_0^{\frac{\pi}{2}} \frac{\cos^2 \theta + 1}{2} d\theta$$

$$= 12 \int_0^{\frac{\pi}{2}} \frac{\cos^2 \theta}{2} d\theta + \int_0^{\frac{\pi}{2}} d\theta$$

$$= 12 \left\{ \left[\frac{\cos \theta \cdot \frac{\pi}{2}}{2} + 0 \right] + \left[\frac{\frac{\pi}{2} - 0}{2} \right] \right\}$$

$$= 12 \cdot 0 + 12 \cdot \frac{\pi}{2} \times \frac{1}{2}$$

$$= 6\pi \frac{1}{2} = 3\pi$$


 \therefore Area of ellipse (A) = $4 \times 3\pi$

$$= 12\pi \text{ Sq. units}$$


* Find the area of $f(u) = \sin u$ between $x=0$ and $x=2\pi$

→ Solution,

given function $f(u) = \sin u$



$$0 \leq u \leq 2\pi$$



$$\because A_1 = \int_{u=0}^{x=\pi} \sin u du$$

$$\therefore A_2 = - \int_{\pi}^{2\pi} \sin u du$$

$$\therefore A = A_1 + A_2$$

Now,

$$A_1 = \int_0^{\pi} \sin u du$$

$$= \left[-\cos u \right]_0^{\pi}$$

$$= (-\cos \pi + \cos 0^\circ)$$

$$= -(-1) + 1$$

$$= 1 + 1$$

$$= 2$$

$$A_2 = - \int_{\pi}^{2\pi} \sin u du$$

$$= -[-\cos u]_{\pi}^{2\pi}$$

$$= -[-\cos 2\pi + \cos \pi]$$

$$= -[-1 + (-1)]$$

$$= -(-1 - 1)$$

$$= 2$$

\therefore Total area of $\sin u = 2+2$

$$= 4$$

* Find the area enclosed by the curve $y^2 = 4u$ and the line $y = 2u$

→ Solution:

$$y^2 = 4u \quad \dots \dots (i)$$

$$y = 2u \quad \dots \dots (ii)$$

Solving eqn (i) and (ii)

$$(2u)^2 = 4u$$


$$4u^2 = 4u$$

$$4u^2 - 4u = 0$$

$$4u(u-1) = 0$$

$$u=0 \text{ and } u=1$$

$$y=0 \quad y=2$$



$$\text{Area enclosed by parabola } (A_1) = \int_{u=0}^{u=1} 2\sqrt{u} du$$

$$\text{Area enclosed by line } (A_2) = \int_{u=0}^{u=1} 2u du$$

$$\therefore \text{Area enclosed } (A) = A_1 - A_2.$$

here,

$$\text{Area enclosed by parabola } (A_1) = \int_{0}^{1} 2\sqrt{u} du$$

$$= 2 \int_{0}^{1} u^{1/2} du$$

$$= 2 \left[\frac{u^{3/2}}{3/2} \right]_0^1$$

$$= 2 \times \frac{2}{3} \left[u^{3/2} \right]_0^1$$

$$= \frac{4}{3} \left[1^{3/2} \right]$$

$$= \frac{4}{3}$$

again,

$$\text{area enclosed by line } (A_2) = \int_{0}^{1} 2u du$$

$$= 2 \left[\frac{u^2}{2} \right]_0^1$$

$$= 2 \cdot \left(\frac{1}{2} \right)$$

$$= 1$$

$$\therefore \text{Area enclosed } (A) = A_1 - A_2$$

$$= \frac{4}{3} - 1$$

$$= \frac{1}{3} \text{ Ans.}$$

* Find the area of the region between the x-axis and the graph $f(u) = u^3 - u^2 - 2u$ $-1 \leq u \leq 2$

→ Solution,

When, $u=0, y=0$

$$u=1 \quad y=1-1-2 = -2$$

$$u=-1 \quad y=-1+1+2 = 0$$

$$u=2 \quad y=0$$

Now,

$$\text{Area}(\Delta_1) = \int_{-1}^0 (u^3 - u^2 - 2u) du$$

$$= \left[\frac{u^4}{4} \right]_{-1}^0 - \left[\frac{u^3}{3} \right]_{-1}^0 - \left[\frac{2u^2}{2} \right]_{-1}^0$$

$$= \left(\frac{0}{4} - \frac{1}{4} \right) - \left(\frac{0}{3} - \frac{1}{3} \right) - (0^2 - 1) = \left(\frac{0}{4} \right) - \left(\frac{1}{4} \right) - \left(\frac{0}{3} \right) + \left(\frac{1}{3} \right) - 0 + 1$$

$$= -\left(\frac{1}{4} - \frac{1}{3} + 1 \right)$$

$$= -\left(\frac{1}{4} - \frac{4}{3} \right)$$

$$= -\left(\frac{3-16}{12} \right) = +\frac{13}{12}$$

$$\therefore \text{Area}(\Delta_1) = -\frac{13}{12} \times -1 = \frac{13}{12} = \frac{5}{12}$$

$$\text{Area}(A)_2 = - \int_0^2 (u^3 - u^2 - 2u) du$$

$$= - \left[\frac{u^4}{4} \right]_0^2 - \left[\frac{u^3}{3} \right]_0^2 - \left[2u \right]_0^2$$


$$= -\left(\frac{16}{4} - \frac{8}{3} - 4 \right)$$

$$= -\left(-\frac{8}{3} \right) = \frac{32}{12}$$

$$= \frac{8}{3}$$

$$\therefore \text{total area} = \left(\frac{13}{12} + \frac{8}{3} \right) = 3.87$$

$$= 1.083 + 2.667 = 3.750$$



* Find the area of region enclosed by parabola

$y = 2 - u^2$ and the line $y = -u$

→ Solution:

$$y = 2 - u^2 \dots (i)$$

$$y = -u \dots (ii)$$

we have,

$$2 - u^2 = -u$$

$$\text{or, } u^2 - u - 2 = 0$$

$$\text{or, } u^2 - 2u + u - 2 = 0$$

$$\text{or, } u(u-2) + 1(u-2) = 0$$

$$\text{or, } (u-2)(u+1) = 0$$

Either $u=2, u=-1$

$$y = -2, y = 1$$

For parabola,

$$y = 2 - u^2$$

$$\text{when } u=0, y=2$$

$$\text{when } u=\pm 1, y=1$$

$$\text{when } u=\pm 2, y=-2$$

Now,

$$\text{Area}(A) = \int_{-1}^0 (2 - u^2) du - \int_{-1}^0 -u du$$

$$= \left[2u - \frac{u^3}{3} \right]_{-1}^0 - \left[-\frac{u^2}{2} \right]_{-1}^0$$


$$= 0 - \left[2 \times (-1) - \frac{(-1)^3}{3} \right] + \left(\frac{0}{2} - \frac{(-1)^2}{2} \right)$$

$$= 2 - \frac{1}{3} - \frac{1}{2}$$

$$= \frac{5}{3} - \frac{1}{2}$$

$$= \frac{10-3}{6}$$

$$= \frac{7}{6} \text{ sq. unit}$$



In y-axis,

$$y = 2 - u^2$$

$$u^2 = 2 - y$$

$$\therefore u = \sqrt{2-y} - 1 \quad \text{--- (1)}$$

$$u = -y \quad \text{--- (2)}$$

Again,

$$\text{Area}(A_2) = \int_{-2}^2 \sqrt{2-y} dy - \int_{-2}^0 -y dy$$

$$= \int_{-2}^2 (2-y)^{\frac{1}{2}} dy + \int_{-2}^0 y dy$$

$$= \left[\frac{(2-y)^{\frac{1}{2}+1}}{\left(\frac{1}{2}+1\right)(-1)} \right]_{-2}^2 + \left[\frac{y^2}{2} \right]_{-2}^0 \quad \left[\because \int (2u+b)^n du = \frac{(2u+b)^{n+1}}{(n+1)2} \right]$$

$$= \left[\frac{(2-y)^{\frac{3}{2}}}{-\frac{3}{2}} \right]_{-2}^2 + \left[\frac{0}{2} - \frac{(-2)^2}{2} \right]$$

$$= \frac{(2-2)^{\frac{3}{2}}}{-\frac{3}{2}} - \frac{(2+2)^{\frac{3}{2}}}{-\frac{3}{2}} + \left(-\frac{4}{2} \right)$$

$$= 0 - \frac{2}{3} (4)^{\frac{3}{2}} + (-2)$$

$$= \frac{2}{3} \times 2^{\frac{3}{2}} - 2$$

$$= \frac{16}{3} - 2$$

$$= \frac{10}{3}$$

$$= \frac{10}{3} \text{ sq. unit.}$$

Hence the area enclosed by parabola $y=2-u^2$ and line $y=-u$ is $A_1 + A_2$

$$A = \left(\frac{7}{6} + \frac{10}{3} \right) \text{ sq. unit}$$

$$= \left(\frac{27}{6} \right) \text{ sq. unit}$$

$$= 4.5 \text{ sq. unit}$$

* Find the area of the region in the first quadrant that is bounded above by $y = \sqrt{u}$ and below by the x-axis and the line $y = u - 2$.

→ Solution.

$$y = \sqrt{u} \cdot \text{Squaring}$$

$$y^2 = u \quad \dots \text{(i)}$$

$$y = u - 2 \quad \dots \text{(ii)}$$

Solving

$$y = \sqrt{u} - 2$$

$$\text{or, } y^2 - y - 2 = 0$$

$$\text{or, } y^2 - 2y + y - 2 = 0$$

$$\text{or } y(y-2) + 1(y-2) = 0$$

$$\therefore y = +2, -1$$

$$u = 4, 1$$

$$\therefore u = 4, 1$$

$$y = 2, -1$$

$$\text{For } y^2 = u$$

$$\text{when } u = 0, y = 0$$

$$\text{when } u = 1, y = 1$$

$$\text{when } u = 4, y = 2$$

$$\begin{array}{ll} y = 0 & u = 0 \\ y = 1 & u = 1 \\ y = 2 & u = 4 \end{array}$$

Now,

$$\text{Area (A)} = \int_0^4 (\sqrt{u}) du - \int_2^4 (u - 2) du$$

$$= \left[\frac{u^{3/2}}{3/2} \right]_0^4 - \left\{ \left[\frac{u^2}{2} \right]_2^4 - \left[2u \right]_2^4 \right\}$$

$$= \frac{2}{3} \times 2^{3/2} - \left\{ \left(\frac{16}{2} - \frac{4}{2} \right) - (2 \times 4 - 2 \times 2) \right\}$$

$$= \frac{16}{3} - (8 - 2 - 8 + 4)$$

$$= \frac{16-2}{3} \text{ (sq. unit)}$$

$$= \frac{10}{3} = 3.33 \text{ sq. unit}$$

∴ Area enclosed = 3.33 sq. unit

→ Solution,

* Find the area of region enclosed by line $u=0$, $y=3$ and curve $u=2y^2$.

→ Solution,

$$u = 2y^2 \quad \dots \text{(i)}$$

$$\text{when } y=3$$

$$u = 3^2 \times 2 = 18$$

$$\text{when } y=0$$

$$u=0$$

$$\text{when } u=2y^2$$

$$u=0$$

$$y=0$$

$$u=2$$

$$y=1$$

$$u=8$$

$$y=2$$

$$u=27$$

$$y=3$$

Now,

$$\text{Area (A)} = \int_0^3 2y^2 \cdot dy$$

$$= 2 \cdot \left[\frac{y^3}{3} \right]_0^3$$

$$= 2 \cdot \left[\frac{3^3}{3} - \frac{0^3}{3} \right]$$

$$= 2 \times 9$$

$$= 18 \text{ sq. unit}$$

Hence area enclosed is 18 sq. unit

* Find the area bounded on right by line $x+y=2$ and left $y=u^2$ below by the x -axis.

→ Solution,

$$x+y=2 \quad \dots \text{(i)}$$

$$y=u^2 \quad \dots \text{(ii)}$$

Solving eqn ① & ②, we get

$$u+u^2=2$$

$$u^2+u-2=0$$

$$u^2+2u-u-2=0$$

$$\text{or } u(u+2)-1(u+2)=0$$

$$\text{or } (u+2)(u-1)=0$$

$$\therefore u = +1, -2$$

$$y = 1, 4$$

$$y = u^2$$

$$x = 1 \quad y = 1$$

$$x = 0 \quad y = 0$$

$$x = 2 \quad y = 4$$

$$x = 3 \quad y = 9$$

Now,

$$\text{Area (A)} = \int_1^2 u^2 du + \int (2-u) du$$

$$= \left[\frac{u^3}{3} \right]_0^1 + \left[2u \right]_1^2 - \left[\frac{u^2}{2} \right]_1^2$$

$$= \left(\frac{1}{3} - 0 \right) + (2 \times 2 - 2) - \left(\frac{1}{2} - \frac{1}{2} \right)$$

$$= \frac{1}{3} + 2 - \frac{3}{2}$$

$$= \frac{1}{3} + \frac{1}{2}$$

$$= \frac{5}{6} \text{ Sq. unit}$$

Hence the area bounded is $\frac{5}{6}$ sq. unit

* Find the area of region in first quadrant bounded by the line $y = u$, that line $u = 2$ and curve $y = \frac{1}{u^2}$, and x-axis.

→ Solution,

$$y = u \text{ and } y = \frac{1}{u^2}$$

$$y = u$$

$$\frac{1}{u^2} = u$$

$$\therefore u^3 = 1$$

$$\therefore x = 1$$

$$y = 1$$

for curve,

$$y = \frac{1}{u^2}$$

when

$$u = 1$$

$$y = 1$$

$$u = 2$$

$$y = 0.25$$

$$\text{Area (A)} = \int_0^1 u du + \int_1^2 u^{-2} du$$

$$= \left[\frac{u^2}{2} \right]_0^1 + \left[\frac{u^{-1}}{-1} \right]_1^2 = \left[\frac{1}{2} - \frac{0}{2} \right] + \left[\frac{1}{2} + \frac{1}{1} \right]$$

$$= \frac{1}{2} + \left(-\frac{1}{2} \right) + 1 = 1 \text{ Sq. unit}$$

Hence the area in 1st quadrant bounded by line $y=u$, line $x=2$, curve $y=\frac{1}{u^2}$ is 1 sq. unit.

* Find the area of triangular region in the first quadrant bounded by the Y-axis and the curves $y = \sin u$, $y = \cos u$.

→ Solution,

$$y = \sin u \quad \text{--- (i)} \quad y = \cos u \quad \text{--- (ii)}$$

$$\sin u = \cos u$$

$$\Rightarrow \tan u = 1$$

$$\Rightarrow \tan u = \tan \frac{\pi}{4}$$

$$\therefore u = \frac{\pi}{4}$$

$$\frac{\pi}{4}$$

$$\frac{\pi}{4}$$

$$\text{Required Area} = \int_0^{\frac{\pi}{4}} \cos u du - \int_0^{\frac{\pi}{4}} \sin u du$$

$$= [\sin u + \cos u]_0^{\frac{\pi}{4}}$$

$$= \sin \frac{\pi}{4} + \cos \frac{\pi}{4} - (\sin 0 + \cos 0)$$

$$= \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} - (0+1)$$

$$= \frac{2}{\sqrt{2}} - 1$$

$$= \frac{2\sqrt{2} - 2}{\sqrt{2}}$$

$$= \sqrt{2} - 1 \text{ sq. unit}$$

Hence the area of triangular region is $\sqrt{2} - 1$ sq. unit

* Find the area betⁿ $y=u$ and $y=u^3$ from

$x=-1$ to $x=1$

→ solution,

$$\text{when } u=1, u=-1$$

$$y=1 \quad y=-1$$

$$\text{for, } y=u^3$$

$$\text{when } u=0, y=0$$

$$u=1, y=1$$

$$u=2, y=8$$

Now,

$$\begin{aligned} \text{Area (A)} &= \int_0^1 u \, du - \int_0^1 u^3 \, du \\ &= \left[\frac{u^2}{2} \right]_0^1 - \left[\frac{u^4}{4} \right]_0^1 \\ &= \left(\frac{1}{2} - 0 \right) - \left(\frac{1}{4} - 0 \right) \\ &= \frac{1}{2} - \frac{1}{4} \\ &= \frac{2-1}{4} \\ &= \frac{1}{4} \end{aligned}$$

∴ Total area betⁿ $y=u$ and $y=u^3$ from

$$x=-1 \text{ to } x=1 \text{ is } = \frac{1}{4} \times 2$$

$$= \frac{1}{2} \text{ sq. unit}$$

* Find the area between two parabola

$$y^2=4ax \text{ and } u^2=4ay$$

→ solution,

$$y^2=4ax \quad \text{--- (i)}$$

$$u^2=4ay \quad \text{--- (ii)}$$

$$\text{or, } y = \frac{u^2}{4a}$$

Now,

$$\left(\frac{u^2}{4a}\right)^2 = 4au$$

~~$$\therefore u^4 = 16a^2 \cdot 4au$$~~

$$\therefore u^4 = 64a^3 u$$

~~$$\therefore u^3 = 4 \times 16a^3$$~~

$$\therefore u^4 - 64a^3 u = 0$$

~~$$\therefore u^3 = 64a^3$$~~

$$\therefore u(u^3 - 64a^3) = 0$$

either $u=0$

$$= u^3 - 64a^3$$

$$\therefore u^3 = 64a^3$$

put cube root on both sides

$$u = 4a$$

Now,

$$\text{Area (A)} = \int_0^{4a} au \, du - \int_0^{4a} \frac{u^2}{4a} \, du$$

$$= \int_0^{4a} 2\sqrt{a} u^{1/2} \, du - \int_0^{4a} \frac{1}{4a} u^2 \, du$$

$$= 2\sqrt{a} \left[\frac{u^{3/2}}{3/2} \right]_0^{4a} - \frac{1}{4a} \left[\frac{u^3}{3} \right]_0^{4a}$$

$$= 2\sqrt{a} \times \frac{2}{3} \left[4a \right]^{3/2} - \frac{1}{4a} \times \frac{1}{3} \left[4a \right]^3$$

$$= \frac{4\sqrt{a}}{3} 2^{2 \cdot \frac{3}{2}} a^{3/2} - \frac{1}{3} 64a^3$$

$$= \frac{4\sqrt{a}}{3} \times 8a^{3/2} - \frac{16a^2}{3}$$

$$= \frac{32a^{1/2 + 3/2}}{3} - \frac{16a^2}{3}$$

$$= \frac{32a^2 - 16a^2}{3}$$

$$= \frac{16a^2}{3} \text{ sq. unit}$$

\therefore Req. Area is $\frac{16a^2}{3}$ sq. unit.

* Find the area bound by x-axis and parabola $y = 4 - u^2$.

→ Solution,

$$\text{In x-axis } y=0$$

$$0 = 4 - u^2$$

$$\therefore u = \sqrt{4}$$

$$\therefore u = \pm 2$$

$$\text{when } u=0$$

$$y=4$$

Now,

$$\text{Area (A)} = \int_{-2}^2 (4 - u^2) du$$

$$= \int_0^2 4 du - \int_0^2 u^2 du$$

$$= [4u]_0^2 - \left[\frac{u^3}{3} \right]_0^2$$

$$= [4 \times 2 - 0] - \left[\frac{2^3}{3} - \frac{0^3}{3} \right]$$

$$= 8 - \frac{8}{3}$$

$$= \frac{24 - 8}{3}$$

$$= \frac{16}{3}$$

$$\therefore \text{Total area (A)} = \frac{16}{3} \times 2$$

$$= \frac{32}{3} \text{ Sq. unit}$$

Hence the area bound by x-axis and parabola $y = 4 - u^2$ is $\frac{32}{3}$ Sq. unit.

A function f is even if the graph of f is symmetric with respect to y -axis. Algebraically f is even if and only if $f(-u) = f(u)$ for all u in the domain of f .

A function f is odd if the graph of f is symmetric with respect to the origin.

Algebraically, f is odd if and only if $f(-u) = -f(u)$ for all u in the domain of f .

ii) One to One and Onto function:-

The function is one to one (injective) if every element of the codomain is mapped to by at most one element of domain.

The function is onto (surjective) if every element of the codomain is mapped to by at least one element of the domain.

i.e. the image of domain and codomain of function are equal.

iii) Piecewise define function: (hybrid function)


A piecewise-defined function or hybrid function is a function which is defined by multiple sub-functions, each sub-function applying to a certain interval of the main function's domain (a sub domain).

iv) Composite function:-

→ function whose values are found from two given functions by applying one function to an independent variable and then applying the second function to the

result and whose domain consists of those values of the independent variable for which the result yielded by the first function lies in the domain of second function is called composite function.

e.g.



$g(f(x))$ where


$$f(x) = y = 2x + 1$$

$$g(y) = z = \frac{y}{3}$$

Composite function


i) Even function:

e.g.




even function

e.g.




odd function

ii) one to one function




one to one function




onto function

v) Piecewise function:



Polar Equation




Polar co-ordinate Point
 $P(r, \theta)$

$$\frac{y}{r} = \frac{r \sin \theta}{r \cos \theta}$$

$$\tan \theta = \frac{y}{r}$$

$$\therefore \theta = \tan^{-1} \left(\frac{y}{r} \right)$$



$$x^2 + y^2 = r^2 \sin^2 \theta + r^2 \cos^2 \theta$$

$$\text{or, } x^2 + y^2 = r^2 (\sin^2 \theta + \cos^2 \theta)$$

$$\text{or, } r^2 = x^2 + y^2$$


$$\sin \theta = \frac{y}{r} = \frac{y}{h}$$

$$\therefore y = r \sin \theta$$

$$\cos \theta = \frac{x}{r} = \frac{x}{h}$$

$$\therefore x = r \cos \theta$$

Area of the Polar curve:



$$\therefore \text{Area} = \int_{\alpha}^{\beta} \frac{1}{2} r^2 d\theta$$

Polar Equation: \checkmark

Limacons ($r = a \pm b \cos \theta, r = a \pm b \sin \theta$)


→ Inner loops : $\left(\frac{a}{b} < 1 \right)$

→ Cardioid : $\left(\frac{a}{b} = 1 \right)$




→ Dimple : $\left(1 < \frac{a}{b} < 2 \right)$

→ Convex (oval) : $\left(\frac{a}{b} \geq 2 \right)$


1) Inner loop ($\frac{a}{b} < 1$)




$$r = a - b \cos \theta$$




2) cardioid : ($\frac{a}{b} = 1$)




$$r = a - b \cos \theta$$




3. Dimple ($1 < \frac{a}{b} < 2$)




$$r = a - b \cos \theta$$



4. Convex (oval) ($\frac{a}{b} \geq 2$)



$$r = a - b \cos \theta$$



* Find the area of the cardioid: $r = 2(1 - \cos \theta)$

→ Solution,

$$r = 2(1 - \cos \theta)$$

$$r = 2 - 2 \cos \theta$$

$$a = 2, b = 2$$

$$\therefore \frac{a}{b} = 1, \text{ so it is cardioid}$$

Now,

Area of Cardioid is given as,

$$\text{Area (A)} = \int_{0}^{\pi} \frac{1}{2} r^2 d\theta$$

Cardioid $\Rightarrow \theta \rightarrow \pi$

Area is symmetrical divided so,

$$\text{Area (A)} = \frac{1}{2} \times 2 \int_{0}^{\pi} 2^2 (1 - \cos \theta)^2 d\theta$$

$$= 4 \int_{0}^{\pi} (1 - 2 \cos \theta + \cos^2 \theta) d\theta$$

$$= 4 \left[\int_{0}^{\pi} 1 d\theta - 2 \int_{0}^{\pi} \cos \theta d\theta + \int_{0}^{\pi} \frac{1 + \cos 2\theta}{2} d\theta \right]$$

$$= 4 \left[[\theta]_0^{\pi} - 2 [\sin \theta]_0^{\pi} + \frac{1}{2} \left\{ [\theta]_0^{\pi} + \left[\frac{\sin 2\theta}{2} \right]_0^{\pi} \right\} \right]$$

$$= 4 \left\{ (\pi - 0) - 2 (\sin \pi - \sin 0) + \frac{1}{2} (\pi - 0) + \left(\frac{\sin^2 \pi}{2} - \frac{\sin^2 0}{2} \right) \right\}$$

$$= 4 \left\{ (\pi - 0) - 2 (0 - 0) + \frac{1}{2} \pi + \left(\frac{\pi}{2} - \frac{0}{2} \right) \right\}$$


$$= 4 \left\{ \pi + \frac{1}{2} \pi \right\}$$

$$= 6\pi$$

∴ Area of Cardioid is $\frac{6\pi}{2} = 3\pi$ ~~square~~

* Find the area of $r = 4 + 2 \cos \theta$

→ Solution,



$$= \frac{\pi}{3} + \frac{1}{4} - \frac{\sqrt{3}}{2} + 2 \left\{ \frac{\pi}{3} + \frac{0}{2} - \frac{\sqrt{3}}{2} \right\}$$

$$= \frac{\pi}{3} + 2\sqrt{3} + \frac{2\pi}{3} + 2\left(\frac{\sqrt{3}}{2}\right) \times \frac{1}{2}$$

$$= \frac{\pi}{3} + \frac{2\pi}{3} - 2\sqrt{3} + \frac{\sqrt{3}}{2}$$

$$= \frac{3\pi}{3} - \frac{4\sqrt{3} + \sqrt{3}}{2}$$

$$= \frac{8\pi}{3} - \frac{3\sqrt{3}}{2}$$

$$= \frac{\pi}{3} - \frac{3\sqrt{3}}{2}$$

Answer

Hence the area inside the limacon is $\pi - \frac{3\sqrt{3}}{2}$ sq. units.

* find the area outside the loop of limacon.

$$r = 2\cos(\theta + 1)$$

→ Solution

we have, $2\sqrt{3}$

$$\text{Area} = \int \frac{1}{2} \times r^2 d\theta$$

$$2\sqrt{3}$$

$$= \int_{2\sqrt{3}}^0 (1+2\cos\theta)^2 d\theta$$

$$2\sqrt{3}$$


$$= \int_0^{2\sqrt{3}} (1+4\cos\theta+4\cos^2\theta) d\theta$$

$$= \int_0^{2\sqrt{3}} 1 d\theta + 4 \int_0^{2\sqrt{3}} \cos\theta d\theta + 4 \int_0^{2\sqrt{3}} \left(\frac{1+2\cos 2\theta}{2} \right) d\theta$$

$$= [0]_0^{2\sqrt{3}} + 4 [\sin\theta]_0^{2\sqrt{3}} + \frac{4}{2} \left\{ [0]_0^{2\sqrt{3}} + \left[\frac{\sin 2\theta}{2} \right]_0^{2\sqrt{3}} \right\}$$

$$= \frac{2\pi}{3} - 0 + 4 \left[\sin \frac{2\pi}{3} - \sin 0 \right] + 2 \left\{ \frac{2\pi}{3} - 0 + \frac{\sin 2 \cdot \frac{2\pi}{3}}{2} - \frac{\sin 0}{2} \right\}$$

$$= \frac{2\pi}{3} + \frac{4\sqrt{3}}{2} + 2 \left\{ \frac{2\pi}{3} + -\frac{\sqrt{3}}{2} \times \frac{1}{2} - \frac{0}{2} \right\}$$



$$= \frac{2\pi}{3} + \frac{2\sqrt{3}}{2} + \frac{4\pi}{3} - \frac{2\sqrt{3}}{42}$$

$$= \frac{2\pi + 4\pi}{3} + 2\sqrt{3} - \frac{\sqrt{3}}{2}$$

$$= \frac{6\pi}{3} + \frac{4\sqrt{3} - \sqrt{3}}{2}$$

~~2000~~ ~~1000~~

$$= 2\pi + \frac{3\sqrt{3}}{2} \text{ sq. unit.}$$

∴ Hence area outside the loop of limacon is
 $2\pi + \frac{3\sqrt{3}}{2}$ sq. unit

* Find the area of curve $r = 3 + 2 \cos \alpha$.

→ Solution,

Area of curve is given by,


$$= \int_0^{\pi} \frac{1}{2} \times 2 (r)^2 d\alpha$$

$$= \int_0^{\pi} (3 + 2 \cos \alpha)^2 d\alpha$$

$$= \int_0^{\pi} (9 + 12 \cos \alpha + 4 \cos^2 \alpha) d\alpha$$

$$= \int_0^{\pi} 9 d\alpha + \int_0^{\pi} 12 \cos \alpha d\alpha + \int_0^{\pi} 4 \cos^2 \alpha d\alpha$$

$$= 9 \left[\alpha \right]_0^{\pi} + 12 \left[\sin \alpha \right]_0^{\pi} + 4 \int_0^{\pi} \left(\frac{1 + \cos 2\alpha}{2} \right) d\alpha$$



$$= 9(\pi - 0) + 12(\sin \pi - \sin 0) + \frac{9}{2} [0]_0^\pi + \left[\frac{\sin 2\theta}{2} \right]_0^\pi$$

$$= 9\pi + 0 + 2 \left\{ (\pi - 0) + \frac{\sin 2\pi}{2} - \frac{\sin 0}{2} \right\}$$


$$= 9\pi + 2\pi + 0 + 0$$

$$= 11\pi \text{ (sq. unit)}$$


Hence the area of curve is 11π (square unit).

Loop of Lemniscate (Bernoulli)

$$\text{i)} r^2 = a^2 \cos 2\theta$$



$$\text{ii)} r^2 = a^2 \sin 2\theta$$



* Find the area of $r^2 = a^2 \cos 2\theta$.

→ Solution,

$r^2 = a^2 \cos 2\theta$, it is bernouli, so

we have,

$$\theta = \frac{\pi}{4}$$

$$\text{Area } (A) = \int \frac{1}{2} r^2 d\theta$$

$$\theta = 0^\circ$$

$$= \frac{1}{2} \times \frac{a^2}{4} \int_{0^\circ}^{\frac{\pi}{4}} a^2 \cos 2\theta d\theta$$

$$= 2a^2 \int_{0^\circ}^{\frac{\pi}{4}} \frac{\sin^2 \theta}{2} d\theta$$

$$= 2a^2 \left[\frac{\sin^2 \theta}{2} \right]_{0^\circ}^{\frac{\pi}{4}}$$

$$= 2a^2 \left[\frac{\sin^2 \frac{\pi}{4}}{2} - \frac{\sin^2 0}{2} \right]$$

$$= 2a^2 \left(\frac{1}{2} - 0 \right)$$

$$= 2a^2 \times \frac{1}{2}$$

$$= a^2 \text{ Sq. unit}$$

∴ Area of $r^2 = a^2 \cos 2\theta$ is a^2 Sq. unit

* Find the area of $r^2 = a^2 \sin 2\theta$.

→ Solution,

we know, Area of bernouli (A) = $\int \frac{1}{2} r^2 d\theta$

$$\theta = \frac{\pi}{2}$$

$$= \int_{0^\circ}^{\frac{\pi}{2}} \frac{1}{2} \times 2 a^2 \sin^2 \theta d\theta$$

$$= a^2 \left[-\frac{\cos 2\theta}{2} \right]_0^{\frac{\pi}{2}}$$


$$= a^2 \left[-\cos 2 \cdot \frac{\pi}{2} - \frac{\cos 0}{2} \right]$$

$$= a^2 \left(-\left(-\frac{1}{2} \right) - \frac{1}{2} \right)$$

$$= \frac{a^2}{2} + \frac{1}{2} - \frac{1}{2} = a^2 \text{ Sq. unit}$$

* Three leaved Rose:

i) $r = a \cos 3\theta$




max $\theta = 0^\circ$

min when $\theta = \frac{\pi}{6}$

$\therefore \theta_1 = 0^\circ$

$\theta_2 = \frac{\pi}{6}$

ii) $r = a \sin 3\theta$



$\theta = 0^\circ = 0$

min


$\theta = \frac{\pi}{3} = \frac{1}{2}$

$\theta_1 = 0^\circ$

$\theta_2 = \frac{\pi}{3}$

* Four leaved Rose:


i) $r = a \cos 2\theta$



$\therefore \theta_1 = 0^\circ$

$\therefore \theta_2 = \frac{\pi}{4} \quad \times 8$

ii) $r = a \sin 2\theta$




$\theta_1 = 0^\circ$

$\theta_2 = \frac{\pi}{2} \quad \times 9$

* Find the area inside one leaf of the four leaved rose $r = a \cos 2\theta$

→ Solution,

Area of one leaf = $\frac{1}{2} \times 2 \int_0^{\pi/4} (\cos 2\theta)^2 d\theta$



$$\begin{aligned}
 & \text{Q} = \frac{\pi}{4} \quad \text{Q} = \frac{\pi}{4} \\
 & \text{Q} = 0 \quad \text{Q} = 0 \\
 & \int_{0}^{\frac{\pi}{4}} \cos 2\theta d\theta = \int_{0}^{\frac{\pi}{4}} \frac{1 + \cos 4\theta}{2} d\theta = \frac{1}{2} \left[\theta \right]_0^{\frac{\pi}{4}} + \left[\frac{\sin 4\theta}{4} \right]_0^{\frac{\pi}{4}} \\
 & = \frac{1}{2} \left[\frac{\pi}{4} - 0 \right] + \left[\frac{\sin 4 \cdot \frac{\pi}{4}}{4} - \sin 0 \right] \\
 & = \left(\frac{1}{2} \times \frac{\pi}{4} \right) + \frac{1}{2} \left[\frac{\pi}{4} - 0 \right] \\
 & = \frac{\pi}{8} \text{ sq. unit}
 \end{aligned}$$

Hence the area of one leaf on a four leaved rose is $\frac{\pi}{8}$ sq. unit.

* Find the area of three leaved rose $r = 2 \cos 3\theta$
 → Solution,

Area of three leaved rose = $\frac{1}{2} \times 6 \int_{0}^{\frac{\pi}{6}} a^2 \cos^2 3\theta d\theta$

$$= 3a^2 \int_{0}^{\frac{\pi}{6}} \frac{1 + \cos 6\theta}{2} d\theta$$

$$= \frac{3a^2}{2} \left[\theta \right]_0^{\frac{\pi}{6}} + \left[\frac{\sin 6\theta}{6} \right]_0^{\frac{\pi}{6}}$$

$$= \frac{3a^2}{2} \left[\frac{\pi}{6} - 0 \right] + \left[\frac{\sin 6 \cdot \frac{\pi}{6}}{6} - \frac{\sin 6 \cdot 0}{6} \right]$$

$$= \frac{3a^2}{2} \times \frac{\pi}{6}$$

$$= \frac{\pi a^2}{4} \text{ sq. unit}$$

Hence the area of three leaved rose is $\frac{\pi a^2}{4}$ sq. unit

* Find the area of $r = a \sin 3\theta$

→ Solution, $0 = \frac{\pi}{3}$

$$\text{Area (A)} = \int \frac{1}{2} r^2 d\theta$$

$$\theta = 0$$

$$\theta = \frac{\pi}{3}$$

$$= \frac{1}{2} \times 3 \int_{0}^{\frac{\pi}{3}} a^2 \sin^2 3\theta d\theta$$

$$= \frac{3a^2}{2} \int_{0}^{\frac{\pi}{3}} (1 - \cos 6\theta) d\theta$$

$$= \frac{3a^2}{2} \left[\int_{0}^{\frac{\pi}{3}} 1 d\theta - \int_{0}^{\frac{\pi}{3}} \cos 6\theta d\theta \right]$$

$$= \frac{3a^2}{4} \left[[0]_{0}^{\frac{\pi}{3}} - \left[\frac{\sin 6\theta}{6} \right]_{0}^{\frac{\pi}{3}} \right]$$

$$= \frac{3a^2}{4} \times \left(\frac{\pi}{3} - 0 \right) - \left(\frac{\sin^2 \frac{\pi}{3}}{6} - \frac{\sin 0}{6} \right)$$

$$= \frac{3a^2}{4} \left\{ \left(\frac{\pi}{3} \right) - \left(\frac{0}{6} - \frac{0}{6} \right) \right\}$$


$$= \frac{3a^2}{4} \times \frac{\pi}{3}$$

$$= \frac{3\pi a^2}{12}$$

$$= \frac{\pi a^2}{4} \text{ sq. unit}$$

Hence the area of $a \sin 3\theta$
is $\frac{\pi a^2}{4}$ sq. unit.

* Area between Polar Curves:



$$\text{Area between Polar Curves} = \int_{\alpha}^{\beta} \frac{1}{2} r_1^2 d\theta - \int_{\alpha}^{\beta} \frac{1}{2} r_2^2 d\theta$$

$$\text{Area} = \frac{1}{2} \int_{\alpha}^{\beta} (r_1^2 - r_2^2) d\theta$$

* Find the area inside circle $r=1$ but outside the cardioid $r=1-\cos\theta$

→ Solution,

$$r_1 = 1, r_2 = 1 - \cos\theta$$

$$\therefore 1 = 1 - \cos\theta$$

$$\cos\theta = 0$$

$$\therefore \theta = \pm \frac{\pi}{2}$$

$$\text{Area of circle (A_1)} = \int_{0}^{\frac{\pi}{2}} \frac{1}{2} r_1^2 d\theta$$

$$= \frac{1}{2} \times \frac{1}{2} [1]^{\frac{\pi}{2}}_0$$

$$= \left[\frac{\pi}{2} - 0 \right]$$

$$= \frac{\pi}{2} \text{ sq. units}$$

$$\text{Area of Cardioid (A_2)} = \int_{0}^{\frac{\pi}{2}} \frac{1}{2} (1 - \cos\theta)^2 d\theta$$

$$= \int_{0}^{\frac{\pi}{2}} \frac{1}{2} \times 2 (1 - \cos\theta)^2 d\theta$$

$$\begin{aligned}
 & \text{Q. } \frac{\pi}{2} \\
 & = \int (1 - 2\cos\phi + \cos^2\phi) d\phi \\
 & \quad \text{Q. } 0 \\
 & \quad \text{Q. } \frac{\pi}{2} \quad \text{Q. } \frac{\pi}{2} \\
 & = \int_{0}^{\frac{\pi}{2}} 1 d\phi - 2 \int_{0}^{\frac{\pi}{2}} \cos\phi + \int_{0}^{\frac{\pi}{2}} \frac{1 + \cos 2\phi}{2} d\phi \\
 & \quad \text{Q. } 0 \quad \text{Q. } 0 \\
 & = \left[\phi \right]_0^{\frac{\pi}{2}} - 2 \left[\sin\phi \right]_0^{\frac{\pi}{2}} + \frac{1}{2} \left[\phi \right]_0^{\frac{\pi}{2}} + \left[\frac{\sin 2\phi}{2} \right]_0^{\frac{\pi}{2}} \\
 & = \left(\frac{\pi}{2} - 0 \right) - 2 \left[\sin \frac{\pi}{2} - \sin 0 \right] + \frac{1}{2} \left(\frac{\pi}{2} - 0 \right) + \left(\frac{\sin 2 \cdot \frac{\pi}{2}}{2} - \frac{\sin 2 \cdot 0}{2} \right) \\
 & = \frac{\pi}{2} - 2 \times 1 + \frac{1}{2} \times \frac{\pi}{2} + \frac{1}{2} \left(\frac{0}{2} - \frac{0}{2} \right) \\
 & = \frac{\pi}{2} - 2 + \frac{\pi}{4} \\
 & = \frac{3\pi}{4} - 2
 \end{aligned}$$

Now,

Area inside the circle but outside the cardioid

is,

$$\begin{aligned}
 A &= (A_1 - A_2) \\
 &= \frac{\pi}{2} - \left(\frac{3\pi}{4} - 2 \right) \\
 &= \frac{\pi}{2} - \frac{3\pi}{4} + 2 = \frac{2\pi - 3\pi + 8}{4} \\
 &= 2 - \frac{\pi}{4} \text{ sq. unit}
 \end{aligned}$$

* Find the area shared by the circle $r=2$ and cardioid $r=2(1-\cos\phi)$

→ solution,

$$r=2, r=2(1-\cos\phi)$$

$$2=2(1-\cos\phi)$$

$$\text{or, } 1-\cos\phi=1$$

$$\cos\phi=1-1$$

$$\therefore \phi = \pm \frac{\pi}{2}$$

here,

Area of cardioid

$$(A_1) = \int_{0}^{\pi/2} \frac{1}{2} \times 2 (r)^2 d\theta$$

$$= \int_{0}^{\pi/2} 2 (1 - \cos \theta)^2 d\theta$$

$$= 4 \int_{0}^{\pi/2} (1 - 2\cos \theta + \cos^2 \theta) d\theta$$

$$= 4 \left\{ [\theta]_0^{\pi/2} - 2 [\sin \theta]_0^{\pi/2} + \frac{1}{2} \left\{ [\theta]_0^{\pi/2} + \left[\frac{\sin 2\theta}{2} \right]_0^{\pi/2} \right\} \right\}$$

$$= 4 \left\{ \left[\frac{\pi}{2} - 0 \right] - 2 \left[\sin \frac{\pi}{2} - \sin 0 \right] + \frac{1}{2} \left\{ \left[\frac{\pi}{2} - 0 \right] + \sin \frac{2\pi}{2} - \sin 0 \right\} \right\}$$

$$= 4 \frac{\pi}{2} - 4 \times 2 (1 - 0) + \frac{4}{2} \frac{\pi}{2} + 4 \left[\frac{\pi}{2} - 0 \right]$$

$$\Rightarrow 2\pi - 8 + \pi + 0$$

$$\therefore 3\pi - 8$$

Again,

Area of circle (A_2) = $\int_{0}^{\pi/2} \frac{1}{2} \times 2 (2)^2 d\theta$


$$= 4 \int_{0}^{\pi/2} 2 d\theta$$

$$\Rightarrow 4 [\theta]_0^{\pi/2} = 4 \left[\frac{\pi}{2} - 0 \right] = \frac{2\pi}{2} = 2\pi$$


Hence Area Shared by circle & Cardioid is

$$A = A_1 + A_2 = 3\pi - 8 + 2\pi$$

$$= (5\pi - 8) \text{ sq. units}$$



* Length (Perimeter of the Polar Curve) :



$$L = \int_{\theta=\alpha}^{\theta=\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

* Find the length of Cardioid $r = 1 - \cos \theta$

→ Solution,

$$r = 1 - \cos \theta$$

$$r^2 = 1 - 2\cos \theta + \cos^2 \theta$$

$$\left(\frac{dr}{d\theta}\right)^2 = \frac{d(1-\cos \theta)}{d\theta}$$

$$= \frac{d1}{d\theta} - \frac{d\cos \theta}{d\theta}$$

$$= 0 - -\sin \theta$$

$$\therefore \left(\frac{dr}{d\theta}\right)^2 = (\sin \theta)^2$$

$$\theta = \pi$$

$$\therefore \text{length } (L) = \int_{\theta=0^\circ}^{\theta=\pi} \sqrt{1 - 2\cos \theta + \cos^2 \theta + \sin^2 \theta} d\theta$$

$$\theta = 0$$


$$= \int_{\theta=0}^{\theta=\pi} \sqrt{1 - 2\cos \theta + 1} d\theta \quad \therefore \sin^2 \theta = \frac{1 - \cos 2\theta}{2}$$

$$= \int_0^{\pi} \sqrt{2(1 - \cos \theta)} d\theta$$

$$= \int_0^{\pi} \sqrt{2 \cdot 2 \sin^2 \frac{\theta}{2}} d\theta$$

$$\therefore 2 \sin^2 \frac{\theta}{2} = 1 - \cos \theta$$

$$\therefore 2 \sin^2 \frac{\theta}{2} = 1 - \cos \theta$$



$$= \int_0^{\pi/2} 2 \sin \frac{\alpha}{2} d\alpha$$

$$= 2 \left[-\cos \frac{\alpha}{2} \right]_0^{\pi/2} = 2 \cdot 2 \left[-\cos \frac{\pi}{2} + \cos \frac{0}{2} \right]$$

$$= 4 \left[0 - 0 + 1 \right]$$

$$= 4 \times 1 = 4 \times 2 \text{ unit} = 8 \text{ unit}$$

Hence the perimeter of cardioid is 8 unit.

* Find the length of spiral $r = \alpha^2$, $0 \leq \alpha \leq \sqrt{5}$

→ Solution,

$$r = \alpha^2$$

$$r^2 = \alpha^4$$

$$\frac{dr}{d\alpha} = \frac{d\alpha^2}{d\alpha} = 2\alpha$$

$$\left(\frac{dr}{d\alpha} \right)^2 = (2\alpha)^2 = 4\alpha^2$$

$$\begin{aligned} r^2 + \left(\frac{dr}{d\alpha} \right)^2 &= \alpha^4 + 4\alpha^2 \\ &= \alpha^2(\alpha^2 + 4) \end{aligned}$$

Now,

$$\text{length } (L) = \int_0^{\sqrt{5}} \sqrt{r^2 + \left(\frac{dr}{d\alpha} \right)^2} d\alpha$$

$$= \int_0^{\sqrt{5}} \sqrt{\alpha^2(\alpha^2 + 4)} d\alpha$$

$$= \int_0^{\sqrt{5}} \sqrt{\alpha^2 + 4} \alpha d\alpha$$

$$\text{let } y = \alpha^2 + 4$$

diff. $y = \alpha^2 + 4$ w.r.t $d\alpha$

$$\frac{dy}{d\alpha} = 2\alpha \quad \therefore dy = 2\alpha d\alpha \quad \therefore \frac{dy}{2} = \alpha d\alpha$$

$$y = \theta^2 + 4$$

when $\theta = 0$

$$y = 4$$

, when $\theta = 15$

$$y = (\sqrt{5})^2 + 4$$

$$= 9$$

Now, 9

$$\begin{aligned}
 L &= \int \sqrt{y} \cdot \frac{dy}{2} \\
 &= \frac{1}{2} \int y^{1/2} dy \\
 &= \frac{1}{2} \left[\frac{y^{1/2+1}}{\frac{1}{2}+1} \right]_4^9 \\
 &= \frac{1}{2} \left[\frac{y^{3/2}}{3/2} \right]_1^9 \\
 &= \frac{1}{2} \times \frac{2}{3} \left[9^{3/2} - 4^{3/2} \right] \\
 &= \frac{1}{3} \left[3^2 - 2^2 \right]^{3/2} \\
 &= \frac{1}{3} (27 - 8) \\
 &= \frac{19}{3} \\
 &= 6.33 \text{ unit}
 \end{aligned}$$

Hence the length of spiral is 6.33 unit.

* Find the length of cardioid $r = a(1 + \cos\theta)$

→ For Cardioid

$$\theta_1 = 0$$

$$\theta_2 = \pi$$

$$r = a(1 + \cos\theta)$$

$$\rightarrow r^2 = a(1 + \cos\theta)$$

$$r^2 = a^2 (1 + 2\cos\theta + \cos^2\theta)$$

$$\frac{dr}{d\theta} = \frac{a}{r} \frac{d(1 + \cos\theta)}{d\theta} = -a\sin\theta$$

$$\therefore \left(\frac{dr}{d\theta}\right)^2 = a^2 \sin^2\theta$$

Now, $\theta = \pi$

$$\text{Length}(L) = \int_{\theta=0}^{\theta=\pi} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

$$= \int_{\theta=0}^{\theta=\pi} \sqrt{a^2(1 + 2\cos\theta + \cos^2\theta) + a^2 \sin^2\theta} d\theta$$

$$= \int_{\theta=0}^{\theta=\pi} a \sqrt{1 + 2\cos\theta + \cos^2\theta + \sin^2\theta} d\theta$$

$$= a \int_{\theta=0}^{\theta=\pi} \sqrt{2(1 + \cos\theta)} d\theta$$

$$\cos^2\theta = \frac{1 + \cos 2\theta}{2}$$

$$2\cos^2\theta = 1 + \cos 2\theta$$

$$\therefore 2\cos^2\theta = 1 + \cos 2\theta$$

$$= 2 \int_{\theta=0}^{\theta=\pi} \sqrt{2 \cdot 2\cos^2\theta} = 2 \cos\theta$$

$$= 2a \left[\frac{\sin\frac{\theta}{2}}{\frac{1}{2}} \right]_0^{\pi}$$

$$= 2a \times \frac{2}{1} \left[\sin\frac{\pi}{2} - \sin\frac{0}{2} \right]$$

$$= 4a (1 - 0)$$

$$= 4a$$

$$\therefore \text{total length of Cardioid} = 4a \times 2$$

$$= 8a \text{ units}$$

* Length for a function $y = f(u)$, $a \leq u \leq b$.

If a function is continuous and differentiable on closed interval $[a, b]$, then,

$$u=b$$

$$L = \int_{u=a}^{u=b} \sqrt{1 + \left(\frac{dy}{du}\right)^2} du$$

$$y = f(u)$$

For function, $u = f(y)$

$$c \leq y \leq d$$

$$x=d$$

$$L = \int_{x=c}^{x=d} \sqrt{1 + \left(\frac{du}{dy}\right)^2} dy$$

$$x=a$$

$$x=b$$

Note:

If $f(u)$ is not differentiable on closed interval $[a, b]$ then use $f(y)$ and vice-versa.

* Find the length of curve $y = u^{\frac{3}{2}}$, $0 \leq u \leq 1$.

→ Solution,

$$\frac{dy}{du} = \frac{d u^{\frac{3}{2}}}{du} = \frac{3}{2} u^{\frac{3}{2}-1} = \frac{3}{2} u^{\frac{1}{2}}$$

∴ It is differentiable at $u \in [0, 1]$

No@,

$$1 + \left(\frac{dy}{du}\right)^2 = 1 + \left(\frac{3}{2} u^{\frac{1}{2}}\right)^2 = 1 + \frac{9}{4} u = \frac{4+9u}{4}$$

Now,

$$\text{length}(L) = \int_{u=0}^{u=1} \sqrt{1 + \left(\frac{dy}{du}\right)^2} du$$

$$= \int_{0}^{1} \sqrt{\frac{4+9u}{4}} du$$

$$= \frac{1}{2} \int_{0}^{1} \sqrt{4+9u} du$$

$$\begin{aligned}
 &= \frac{1}{2} \int_0^1 (4+9u)^{1/2} du \\
 &= \frac{1}{2} \left[\frac{(4+9u)^{1/2+1}}{\frac{3}{2} \times 9} \right]_0^1 \\
 &= \frac{1}{2} \times \frac{2}{3 \times 9} \left[(4+9u)^{3/2} \right]_0^1 \\
 &= \frac{1}{27} (4+9 \cdot 1)^{3/2} - (4+9 \cdot 0)^{3/2} \\
 &= \frac{1}{27} (13)^{3/2} - (4)^{3/2} \\
 &= \frac{1}{27} \times 46.87 - 2^{3/2} \\
 &= \frac{38.87}{27} \\
 &= 1.439 \text{ unit}
 \end{aligned}$$

\therefore length of curve is 1.439 unit

* Find the length of curve $y = \left(\frac{u}{2}\right)^{2/3}$ from $x=0$ to $x=2$

→ Solution,

$$\frac{dy}{du} = \frac{1}{2} \left(\frac{u}{2}\right)^{\frac{2}{3}-1} \times \frac{1}{3} u^{\frac{2}{3}-1} = \frac{1}{3} \times \frac{u}{2}$$

$$\frac{dy}{dx} = \left(\frac{1}{2}\right)^{\frac{2}{3}} \times \frac{2}{3} (u)^{\frac{2}{3}-1} = 0.62 \times \frac{2}{3 \sqrt[3]{u}} = \frac{0.41}{\sqrt[3]{u}}$$

$$\text{when } u=0 \quad \frac{0.41}{0} = \infty$$

so it is not differentiable at $u \in [0, 2]$

Again,

$$\frac{dy}{dx} = ? \quad \text{so, } y = \left(\frac{u}{2}\right)^{2/3}$$

$$y^{\frac{3}{2}} = \frac{u}{2} \quad ; \quad u = 2y^{\frac{3}{2}}$$

$$u = 2y^{3/2}$$

when $u=0$
 $0 = 2y^{3/2}$
 $\therefore y = 0$

$$\text{when } u=2$$

$$2 = 2y^{3/2} \rightarrow y^{3/2} = 1$$

Now,

$$\frac{du}{dy} = 2 \frac{dy^{3/2}}{dy} = 2 \times \frac{3}{2} y^{3/2-1} = 3y^{1/2}$$

$$\therefore 1 + \left(\frac{dy}{dx} \right)^2 = 1 + (3y^{1/2})^2 = 1 + 9y$$

$$\therefore \text{length}(l) = \int_{y=0}^{y=1} \sqrt{1 + \left(\frac{dy}{dx} \right)^2} dy$$

$$= \int_0^1 \sqrt{1 + 9y} dy$$

$$= \int_0^1 (1 + 9y)^{1/2} dy$$

$$= \int_0^1 \frac{(1 + 9y)^{1/2+1}}{(1/2+1) \cdot 9} dy$$

$$= \frac{2}{3 \cdot 9} (1 + 9 \cdot 1)^{3/2} - (1 + 9 \cdot 0)^{3/2}$$

$$= \frac{2}{27} (10)^{3/2} - 1$$

$$= \frac{61.24}{27}$$

$$= \frac{61.24}{27}$$

$$= 2.26 \text{ unit}$$

Hence the length curve is 2.26 unit.

* Find the length of curve $y = \frac{4\sqrt{2}}{3} u^{\frac{3}{2}} - 1$ $0 \leq u \leq 1$.

→ Solution,

$$\frac{dy}{du} = \frac{d \frac{4\sqrt{2}}{3} u^{\frac{3}{2}} - 1}{du}$$

$$= \frac{4\sqrt{2}}{3} \times \frac{3}{2} u^{\frac{1}{2}} - 0$$

$$= 2\sqrt{2} u^{\frac{1}{2}}$$

$$= 2\sqrt{2}u$$

Now,

$$1 + \left(\frac{dy}{du} \right)^2 = (2\sqrt{2}u)^2 = 1 + 8u$$

Now,

$$\text{length } (l) = \int_0^1 \sqrt{1 + \left(\frac{dy}{du} \right)^2} du$$

$$= \int_0^1 \sqrt{1 + 8u} du$$

$$= \int_0^1 (1 + 8u)^{\frac{1}{2}} du$$

$$= \left[\frac{(1 + 8u)^{\frac{1}{2}}}{\frac{1}{2} + 1 \cdot 8} \right]_0^1$$

$$= (1 + 8u)^{\frac{1}{2}} \times \frac{2}{3} \times 8$$

$$= \frac{2}{3} (1 + 8 \cdot \frac{1}{8})^{\frac{1}{2}} - (1 + 8 \cdot 0)^{\frac{1}{2}}$$

$$= \frac{2}{3} (3)^{\frac{1}{2}} - 1^{\frac{1}{2}}$$

$$= \frac{2 \times 26}{24}$$

$$\Rightarrow 2.16 \text{ unit}$$

∴ Hence the length of Curve is 2.16 unit

→ solution,

$$\frac{dy}{du} = \frac{1}{3} \times \frac{3}{2} (u^2+2)^{\frac{3}{2}-1} = \frac{1}{2} (u^2+2)^{\frac{1}{2}}$$

Now,

$$\begin{aligned} 1 + \left(\frac{dy}{du} \right)^2 &= 1 + \left(\frac{1}{2} (u^2+2)^{\frac{1}{2}} \right)^2 \\ &= 1 + \frac{1}{4} (u^2+2) \\ &= \frac{4+u^2+2}{4} \\ &= \frac{6+u^2}{4} \end{aligned}$$

$$\begin{aligned} \frac{dy}{du} &= \frac{1}{3} \frac{d}{d(u^2+2)} (u^2+2)^{\frac{3}{2}} \times \frac{d}{du} (u^2+2) \\ &= \frac{1}{3} \times \frac{3}{2} (u^2+2)^{\frac{3}{2}-1} \times 2u \\ &= (u^2+2)^{\frac{1}{2}} \cdot 2u \\ 1 + \left(\frac{dy}{du} \right)^2 &= 1 + 4u^2(u^2+2) \end{aligned}$$

→ solution,

$$y = \frac{1}{3} (u^2+2)^{\frac{3}{2}}$$

Now,

$$\begin{aligned} \frac{dy}{du} &= \frac{1}{3} \frac{d}{d(u^2+2)} (u^2+2)^{\frac{3}{2}} \times \frac{d}{du} (u^2+2) \\ &= \frac{1}{3} \times \frac{3}{2} (u^2+2)^{\frac{1}{2}} \times 2u \\ &= u(u^2+2)^{\frac{1}{2}} \end{aligned}$$

$$\begin{aligned} 1 + \left(\frac{dy}{du} \right)^2 &= 1 + \left\{ u(u^2+2)^{\frac{1}{2}} \right\}^2 = 1 + u^2(u^2+2) \\ &= 1 + u^4 + 2u^2 \end{aligned}$$

$$= 1 + 2u^2 + u^4$$

$$= (1+u^2)^2$$

Now,

$$\text{length}(l) = \int_0^3 \sqrt{(1+u^2)^2} du$$

$$= \int_0^3 (1+u^2) du$$

$$= [u]^3_0 + \left[\frac{u^3}{3} \right]^3_0$$

$$= (3-0) + \left(\frac{3^3}{3} - \frac{0^3}{3} \right)$$

$$= 3 + \frac{27}{3}$$

$$= 12 \text{ unit.}$$

Hence the length of curve is 12 unit.

* Find the length of curve $u = \left(\frac{y^4}{4}\right) + \frac{1}{8}y^2 \quad 1 \leq y \leq 2$

→ Solution,

$$\frac{du}{dy} = \frac{d}{dy} \left(\frac{y^4}{4} \right) + \frac{1}{8}y^2$$

$$= \frac{1}{4} \times 4y^3 + \frac{1}{8} \times (-2)y^{-3}$$

$$= y^3 - \frac{1}{4}y^{-3}$$

Now,

$$1 + \left(\frac{du}{dy} \right)^2 = 1 + \left(y^3 - \frac{1}{4}y^{-3} \right)^2$$

$$= \left\{ 1 + y^3 - 2 \cdot y^3 \cdot \frac{1}{4}y^{-3} + \left(\frac{1}{4}y^{-3} \right)^2 \right\}$$

$$= 1 - \frac{1}{2} + y^3 + \left(\frac{1}{4}y^{-3} \right)^2$$

$$= (y^3)^2 + 2 \cdot y^3 \cdot \frac{1}{4}y^{-3} + \left(\frac{1}{4}y^{-3} \right)^2$$

$$= \left(y^3 + \frac{1}{4y^3} \right)$$

Now,

$$\text{Length} = \int_1^2 \sqrt{1 + \left(\frac{dy}{dx} \right)^2} dy$$

$$= \int_1^2 \sqrt{\left(y^3 + \frac{1}{4y^3} \right)^2} dy$$

$$= \int_1^2 \left(y^3 + \frac{1}{4y^3} \right) dy$$

$$= \left[\frac{y^4}{4} \right]_1^2 + \frac{1}{4} \left[\frac{y^{-3+1}}{-3+1} \right]_1^2$$

$$= \left(\frac{2^4 - 1}{4} \right) + \frac{1}{4} \left[\frac{2^{-2} - 1^{-2}}{-2} \right]$$

$$= \left(\frac{16-1}{4} \right) + \frac{1}{-8} \left[\frac{1}{2^2} - \frac{1}{1^2} \right]$$

$$= \frac{15}{4} - \frac{1}{8} \left(\frac{1}{4} - 1 \right)$$

$$= 3.75 - 0.125 (-0.75)$$

$$= 3.75 + 0.093$$

$$= 3.84375 \text{ unit}$$

Hence the length of curve is 3.84 unit.

* Find the length of curve $y = \frac{y^3}{3} + \frac{1}{4y}$, $1 \leq y \leq 3$

→ Solution,

$$\frac{dy}{dx} = \frac{d\left(\frac{y^3}{3}\right) + \frac{1}{4y}}{dy}$$

$$= \frac{1}{3} \times 3y^2 + \frac{1}{-4} y^{-2}$$

$$= y^2 - \frac{1}{4y^2}$$

Now,

$$1 + \left(\frac{dy}{dx} \right)^2 = 1 + \left(y^2 - \frac{1}{4y^2} \right)^2$$

$$= 1 + y^4 - 2 \cdot y^2 \cdot \frac{1}{4y^2} + \left(\frac{1}{4y^2} \right)^2$$

$$= 1 - \frac{1}{2} + (y^2)^2 + \left(\frac{1}{4y^2} \right)^2$$

$$= \frac{1}{2} + (y^2)^2 + \left(\frac{1}{4y^2} \right)^2$$

$$= (y^2)^2 + 2 \cdot y^2 \cdot \frac{1}{4y^2} + \left(\frac{1}{4y^2} \right)^2$$

$$= \left(y^2 + \frac{1}{4y^2} \right)^2$$

Now,

$$\text{length (L)} = \int_1^3 \sqrt{1 + \frac{dy}{dx}} dy$$

$$= \int_1^3 \sqrt{\left(y^2 + \frac{1}{4y^2} \right)^2} dy$$

$$= \int_1^3 \left(y^2 + \frac{1}{4y^2} \right) dy + \frac{1}{4y^2} = \frac{1}{4} y^2 = \frac{-2}{y^3}$$

$$= \left[\frac{y^3}{3} \right]_1^3 + \frac{-2}{4} \left[\frac{y^{-2+1}}{-2+1} \right]_1^3$$

$$= \left(\frac{3 \times 3 \times 3}{3} - \frac{1}{3} \right) + \frac{-2}{4} \left[\frac{y^{-1}}{-1} \right]_1^3$$

$$= \frac{26}{3} + \frac{-2}{4} \left[\frac{1}{3} - \frac{1}{1} \right]$$

$$= 8.66 - 0.25 \times (-0.666)$$

$$= 8.66 + 0.1666$$

$$= 8.766 \text{ unit}$$

∴ The length of curve is 8.766 unit.

* Find the length of the circle $x^2 + y^2 = a^2$

→ solution

~~parametric~~

$$u^2 + y^2 = a^2$$

differentiating both sides w.r.t u

we get

$$\frac{du^2}{du} + \frac{dy^2}{dy} \times \frac{dy}{du} = \frac{da^2}{du}$$

$$2u + 2y \frac{dy}{du} = 0$$

$$\therefore 2y \frac{dy}{du} = -2u$$

$$\therefore \frac{dy}{du} = -\left(\frac{u}{y}\right)$$

$$\left(\frac{dy}{du}\right)^2 = \frac{u^2}{y^2}$$

Now,

$$\text{length } (L) = \int_{x=0}^{x=a} \sqrt{1 + \left(\frac{dy}{du}\right)^2} du$$

$$= \int_{x=0}^{x=a} \sqrt{1 + \frac{u^2}{y^2}} du$$


$$= \int_{x=0}^{x=a} \sqrt{\frac{u^2 + y^2}{y^2}} du$$

$$= \int_{x=0}^{x=a} \sqrt{\frac{a^2}{a^2 - u^2}} du$$

$$= \int_{x=0}^{x=a} \frac{a}{\sqrt{a^2 - u^2}} du$$

For $\sqrt{a^2 - u^2}$,

Put $u = a \sin \theta$



$$\begin{aligned} \therefore x^2 + y^2 &= a^2 \\ \therefore y^2 &= a^2 - u^2 \end{aligned}$$

diff both sides

$$\frac{du}{d\theta} = \frac{a \sin \theta}{a}$$

$$\therefore du = a \cos \theta d\theta$$

when $\theta = 0$

$$0 = a \sin 0$$

$$\therefore 0 = \sin^{-1}(0) = 0$$

when $\theta = \pi/2$

$$a = a \sin \theta$$

$$\therefore \sin \theta = 1$$

$$\therefore 0 = \sin^{-1}(1) = \frac{\pi}{2}$$

Now

$$\text{Length}(l) = \int_0^{\pi/2} \sqrt{a^2 - u^2} du$$

$$= \int_0^{\pi/2} \frac{a \cdot a \cos \theta d\theta}{\sqrt{a^2 - a^2 \sin^2 \theta}}$$

$$= \int_0^{\pi/2} \frac{a^2 \cos \theta d\theta}{\sqrt{a^2(1 - \sin^2 \theta)}}$$

$$= a^2 \int_0^{\pi/2} \frac{\cos \theta d\theta}{\sqrt{\cos^2 \theta}}$$

$$= a \int_0^{\pi/2} \frac{\cos \theta d\theta}{\cos \theta}$$

$$= a \left[\theta \right]_0^{\pi/2}$$

$$= a [\pi/2 - 0]$$

$$= \frac{\pi a}{2}$$

$$\therefore \text{Total Area of circle: } \frac{\pi a}{2} \times 4 = 2\pi a \text{ unit}^2$$

Hypocycloid

$$\left(\frac{u}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} = 1$$

(0, b)

(-a, 0)

(a, 0)

(0, -b)

Astroid ($a=b$)

$$\left(\frac{u}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{a}\right)^{\frac{2}{3}} = 1$$

$$-(x)^{\frac{2}{3}} + (y)^{\frac{2}{3}} = a^{\frac{2}{3}}$$

(0, a)

(-a, 0)

(a, 0)

(0, -a)

* Find the perimeter (length) of astroid $u^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$

→ Solution:

differentiating both sides, we get,

$$\frac{2}{3}u^{\frac{2}{3}-1} + \frac{2}{3}y^{\frac{2}{3}-1} \frac{dy}{du} = 0$$

$$\frac{2}{3\sqrt[3]{u}} + \frac{2}{3\sqrt[3]{y}} \frac{dy}{du} = 0$$

$$\frac{dy}{du} = -\frac{1}{\sqrt[3]{u}} \times \frac{\sqrt[3]{y}}{\sqrt[3]{u}}$$

$$\therefore \left(\frac{dy}{du}\right)^2 = \left(\frac{\sqrt[3]{y}}{\sqrt[3]{u}}\right)^2 = \frac{y^{\frac{2}{3}}}{u^{\frac{2}{3}}}$$

Now,

$$\text{length}(L) = \int_0^a \sqrt{1 + \frac{y^{\frac{2}{3}}}{u^{\frac{2}{3}}}} du$$

$$[u^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}]$$

$$= \int_0^a \sqrt{\frac{u^{\frac{2}{3}} + y^{\frac{2}{3}}}{u^{\frac{2}{3}}}} du$$

$$= \int_0^a \sqrt{\frac{a^{\frac{2}{3}}}{u^{\frac{2}{3}}}} du$$

$$= \int_0^a \sqrt{\left(\frac{a}{u}\right)^{\frac{2}{3}}} du$$

$$= \int_0^a \left(\frac{a}{u}\right)^{\frac{2}{3} \times \frac{1}{2}} du$$

$$= a^{\frac{1}{3}} \int_0^a \left(\frac{1}{u}\right)^{\frac{1}{3}} du$$

~~$$= a^{\frac{1}{3}} \int_0^a u^{-\frac{1}{3}} du$$~~

$$= a^{\frac{1}{3}} \left[\frac{u^{-\frac{1}{3}+1}}{-\frac{1}{3}+1} \right]_0^a$$

$$= a^{\frac{1}{3}} \times \frac{3}{2} \left[u^{\frac{2}{3}} \right]_0^a$$

$$= \frac{3a^{\frac{1}{3}}}{2} \left[a^{\frac{2}{3}} - 0^{\frac{2}{3}} \right]$$

$$= \frac{3a^{\frac{1}{3}}}{2} \cdot a^{\frac{2}{3}}$$

$$= \frac{3}{2} \times a^{\frac{1}{3} + \frac{2}{3}}$$

$$= \frac{3a}{2}$$

Total area of the asteroid is $= 4 \times \frac{3a}{2} = 6a$ unit.

~~total area of an asteroid~~

Length of parametric curve

If $x = f(t)$ and $y = g(t)$ and $a \leq t \leq b$

then,

$$L = \int_{t=a}^{t=b} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

* Find the length of asteroid $x = \cos^3 t$ and $y = \sin^3 t$

$$0 \leq t \leq 2\pi/3$$

→ Solution,

$$\frac{dx}{dt} = \frac{d \cos^3 t}{d \cos t} \times \frac{d \cos t}{dt}$$

$$= 3 \cos^2 t \times (-\sin t)$$

$$\therefore \left(\frac{dx}{dt}\right)^2 = -9 \sin^2 t \cdot \cos^2 t = 9 \sin^2 t \cdot \cos^4 t$$

$$\frac{dy}{dt} = \frac{d \sin^3 t}{d \sin t} \times \frac{d \sin t}{dt}$$

$$= 3 \sin^2 t \times \cos t$$

$$\therefore \left(\frac{dy}{dt}\right)^2 = 9 \sin^4 t \cdot \cos^2 t = 9 \sin^4 t \cdot \cos^2 t$$

$$\text{Now, } t \in [0, \pi/2]$$

$$L = \int_{t=0}^{t=\pi/2} \sqrt{9 \sin^2 t \cdot \cos^4 t + 9 \sin^4 t \cdot \cos^2 t} dt$$

$$= \int_{t=0}^{t=\pi/2} \sqrt{9 \sin^2 t \cos^2 t (\cos^2 t + \sin^2 t)} dt$$

$$= 3 \int_{t=0}^{t=\pi/2} \sin t \cos t dt = \frac{3}{2} \left[-\frac{\cos 2t}{2} \right]_0^{\pi/2}$$

$$= \frac{3}{2} \int_{t=0}^{t=2\pi} 2 \sin t \cos t dt$$

$$= \frac{3}{2} \int_{t=0}^{t=2\pi} \sin 2t dt$$

$$= \frac{3}{2} \left[-\frac{\cos 2t}{2} \right]_0^{2\pi}$$

$$= \frac{3}{4} (0) = 0$$

$$= \frac{3}{4}$$

\therefore Total length of asteroid = $4x$

Now, $t = \frac{\pi}{2}$

$$L = \int_{t=0}^{\pi/2} \sqrt{9 \sin^2 t \cdot \cos^2 t \cdot (\cos^2 t + \sin^2 t)} dt$$

$$= 3 \int_0^{\pi/2} \sin t \cdot \cos t \cdot dt$$

$$= \frac{3}{2} \int_0^{\pi/2} 2 \cdot \sin t \cdot \cos t \cdot dt$$

$$= \frac{3}{2} \int_0^{\pi/2} \sin 2t \cdot dt$$

$$= \frac{3}{2} \left[-\frac{\cos 2t}{2} \right]_0^{\pi/2}$$

$$= \frac{3}{2} \left[-\frac{\cos 2 \cdot \pi/2}{2} + \frac{\cos 2 \cdot 0}{2} \right]$$

$$= \frac{3}{4} [+1 + 1]$$

$$= \frac{3}{4} \times 2$$

$$= \frac{3}{2}$$

\therefore Total length of asteroid = $24x \frac{3}{2}$

$$= 6 \text{ units}$$


* Find the length of circle.

$$x = a \cos t, y = a \sin t, 0 \leq t \leq 2\pi$$

→ Solution,

here,

we have,



$$v = a \cos t$$

$$\therefore \frac{dv}{dt} = -a \sin t$$

$$y = a \sin t$$

$$\frac{dy}{dt} = a \cos t$$

Now,

$$\text{Length} \rightarrow \int_0^{2\pi} \sqrt{\left(\frac{dv}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$= \int_0^{2\pi} \sqrt{a^2 \sin^2 t + a^2 \cos^2 t} dt$$

$$= \int_0^{2\pi} \sqrt{a^2 (\sin^2 t + \cos^2 t)} dt$$

$$= \int_0^{2\pi} a dt$$

$$= 2 \left[t \right]_0^{2\pi}$$

$$= 2 \times [2\pi - 0]$$

$$= 2\pi a \text{ unit.}$$

Hence length of circle is $2\pi a$ unit.

Mean Value Theorem

1. Rolle's Theorem

2. Langrange's Mean Value Theorem

1.1 Rolle's Theorem :-

If a function $f(x)$ is,


a) continuous in closed interval $[a, b]$

b) differentiable (derivable) in open interval (a, b)

c) $f(a) = f(b)$,

then there exists at least a point $c \in (a, b)$

such that,



* Verify Rolle's Theorem,

$$a) f(u) = u(u-3)^2, u \in [0, 3]$$

→ Solution,

$$f(u) = u(u^2 - 6u + 9)$$

$$= u(u^2 - 6u + 9)$$

$$\therefore f(u) = u^3 - 6u^2 + 9u$$

function $f(u)$ is a polynomial function which exists for all $u \in [0, 3]$

so it is continuous.

$$f'(u) = 3u^2 - 12u + 9$$

function $f(u)$ is differentiable for all $u \in (0, 3)$

then,

$$f(a) = f(b)$$

$$f(a) = f(0) = 0^3 - 6 \times 0^2 + 9 = 0$$

$$f(b) = f(3) = 3^3 - 6 \times 3^2 + 9 \cdot 3 = 27 - 54 + 27 = 0$$

$$\therefore f(a) = f(b)$$

then

there exists at least a point $c \in (0, 3)$, such that

$$f'(c) = 0$$

$$\text{or, } 3c^2 - 12c + 9 = 0 \quad \text{or, } c(c-3) - 1(c-3) = 0$$

$$\text{or, } c^2 - 4c + 3 = 0 \quad \text{or, } (c-3)(c-1) = 0$$

$$\text{or, } c^2 - 3c - c + 3 = 0$$

Either,

$$c = 3, 1$$

$$\therefore c = 3 \notin (0, 3)$$

$$\therefore c = 1 \in (0, 3)$$

Hence Rolle's theorem is Verified.

b) $f(u) = \sqrt{16-u^2}, u \in [-4, 4]$

→ Solution,

$$f(u) = \sqrt{16-u^2}$$

function $f(u)$ is polynomial function, so it is continuous, which exists for all $[-4, 4]$

$$f'(u) = \frac{d(16-u^2)^{1/2}}{d(16-u^2)} \times \frac{d(16-u^2)}{du}$$

$$= \frac{1}{2} (16-u^2)^{1/2-1} (x-2u)$$

$$= -\frac{u}{\sqrt{16-u^2}}$$

function $f(u)$ is differentiable at open interval $u \in (-4, 4)$

Now,

$$f(a) = f(-4) = \sqrt{16 - (-4)^2} = 0$$

$$f(b) = f(4) = \sqrt{16 - 4^2} = 0$$

$$\therefore f(a) = f(b)$$

For more notes visit <https://collegenote.pythonanywhere.com>
then there exists at least a point $c \in (-4, 4)$,
such that,

$$f'(c) = 0$$

$$\text{or } \frac{-c}{\sqrt{16 - c^2}} = 0$$

$$\text{or, } c = 0 \in (-4, 4)$$

Hence Rolle's theorem is verified.

c) Verify Rolle's theorem,

$$f(u) = \sin u, u \in [0, \pi]$$

→ function $f(u)$ is trigonometry function which exists for all $u \in [0, \pi]$

so it is continuous.

$$f'(u) = \cos u$$

function $f(u)$ is differentiable for all $u \in (0, \pi)$

then,

$$f(a) = f(0) = \sin 0 = 0$$

$$f(b) = f(\pi) = \sin \pi = 0$$

then there exists at least a point $c \in (0, \pi)$,

such that,

$$f'(c) = 0$$

$$\cos c = 0$$

$$\therefore c = \cos^{-1}(0)$$

$$\therefore c = \frac{\pi}{2} \in (0, \pi)$$

Hence Rolle's theorem is Verified.

d) $f(u) = \cos 2u, u \in [\pi, -\pi]$

→ solution

function $f(u)$ is trigonometry function which exists for all $u \in [\pi, -\pi]$

so it is continuous.

$f'(u) = -2\sin 2u$
 \Leftrightarrow function $f(u)$ is differentiable for $u \in (-\pi, \pi)$

Now,

$$f(a) = f(-\pi) = \cos 2(-\pi) = 1$$

$$f(b) = f(\pi) = \cos 2\pi = 1$$

$$\therefore f(a) = f(b)$$

then there exists at least a point constant $c \in (-\pi, \pi)$ such that,

$$f'(c) = 0$$

$$-2\sin 2c = 0$$

$$\sin 2c = 0$$

$$2c = \sin'(0)$$

$$\therefore c = 0 = 0 \in (-\pi, \pi)$$

Hence Rolle's theorem Verified.

$$\Leftrightarrow f(u) = e^u (\sin u - \cos u) \quad u \in \left[\frac{\pi}{4}, \frac{5\pi}{4} \right]$$

→ Solution,

function $f(u)$ is trigonometry and exponential function which exists for all $u \in \left[\frac{\pi}{4}, \frac{5\pi}{4} \right]$

so, it is continuous.

$$\frac{d(u \cdot v)}{du} = u \frac{dv}{du} + v \frac{du}{du}$$

$$f'(u) = e^u \frac{d(\sin u - \cos u)}{du} + (\sin u - \cos u) \frac{de^u}{du}$$

$$= e^u (\cos u + \sin u) + (\sin u - \cos u) e^u$$

$$= e^u (\cos u + \sin u + \sin u - \cos u)$$

$$= e^u 2\sin u$$

function $f(u)$ is differentiable for all $u \in \left(\frac{\pi}{4}, \frac{5\pi}{4} \right)$

then $f(a) = f(b)$

then,

$$f(a) = f\left(\frac{\pi}{4}\right) = e^{\frac{\pi}{4}} \left(\sin \frac{\pi}{4} - \cos \frac{\pi}{4} \right) = e^{\frac{\pi}{4}} \times (0.7 - 0.7) = 0$$

$$f(b) = f\left(\frac{5\pi}{4}\right) = e^{\frac{5\pi}{4}} \left(\sin \frac{5\pi}{4} - \cos \frac{5\pi}{4}\right)$$

$$= e^{\frac{5\pi}{4}} (-0.70 + 0.70)$$

$$= 0$$

$$\therefore f(a) = f(b)$$

then,

there exists at least a constant point

$$c \in \left(\frac{\pi}{4}, \frac{5\pi}{4}\right)$$

such that,

$$f'(c) = 0$$

$$\text{or, } e^c 2 \sin c = 0$$

$$\text{or, } \sin c = \frac{0}{2e^c}$$

$$\text{or, } c = \sin^{-1} 0$$

~~0, \pi/2, \pi, 3\pi/2~~

$$\text{or, } c = 0, \pi$$

$$c = 0 \notin \left(\frac{\pi}{4}, \frac{5\pi}{4}\right)$$

$$c = \pi \in \left(\frac{\pi}{4}, \frac{5\pi}{4}\right)$$

Hence Rolle's theorem is verified.

Lagrange Mean Value Theorem (MVT)

If a function $f(u)$ is

a) continuous in closed interval $[a, b]$


b) differentiable in open interval (a, b)

then,

there exists at least a point $c \in (a, b)$

such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$



* Verify Mean Value Theorem for the following functions.

a) $f(u) = u(u-1)(u-2)$ $u \in [0, \frac{1}{2}]$

→ Solution,

$$\begin{aligned}
 f(u) &= u^3 - u^2 - 2u \\
 &= u^2(u-1) - u(u-2) \\
 &= u^3 - 2u^2 + u^2 + 2u \\
 &= u^3 - 3u^2 + 2u
 \end{aligned}$$

Since the function $f(u)$ is polynomial function which exists for all $u \in [0, \frac{1}{2}]$

$$f'(u) = 3u^2 - 6u + 2$$

function $f(u)$ is differentiable for all $u \in (0, \frac{1}{2})$

then

there exists at least a point $c \in (0, \frac{1}{2})$

such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f(a) = f(0) = 0^3 - 3 \cdot 0^2 + 2 \cdot 0 = 0$$

$$f(b) = f\left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)^3 - 3\left(\frac{1}{2}\right)^2 + \frac{1}{2} \cdot 2$$

$$= \frac{1}{8} - \frac{3}{4} + 1$$

$$= \frac{1}{8} - \frac{3}{4} + 1 \Rightarrow \frac{1-6+8}{8} = \frac{3}{8}$$

Now,

$$\frac{f'(c)}{b-a} = f(b) - f(a)$$

$$3c^2 - 6c + 2 = \frac{\frac{3}{8} - 0}{\frac{1}{2} - 0}$$

$$3c^2 - 6c + 2 = \frac{3}{8} \times \frac{2}{1}$$

$$12c^2 - 24c + 8 = 3$$

$$12c^2 - 24c + 5 = 0 \dots \dots \text{(i)}$$

Comparing eqn (i) with $ax^2 + bx + c$

where $a = 12, b = -24, c = 5$

Now,

$$c = -\frac{b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-24 \pm \sqrt{(-24)^2 - 4 \times 12 \times 5}}{2 \times 12}$$

$$= \frac{-24 \pm 18.33}{24}$$

$$= \frac{42.33}{24}, \frac{5.67}{24}$$

$$= 1.76, 0.236$$

$$\therefore c = 1.76 \notin (0, \frac{1}{2})$$

$$\therefore c = 0.236 \in (0, \frac{1}{2})$$

Hence Mean value Theorem is verified.

b) $f(u) = \sqrt{u^2 - 4}$, $u \in [2, 4]$

→ Solution,

$$f(u) = (u^2 - 4)^{\frac{1}{2}}$$

function $f(u)$ is polynomial function which exists for all $u \in [2, 4]$

so it is continuous

$$f'(u) = \frac{d(u^2 - 4)^{\frac{1}{2}}}{d(u^2 - 4)} \times \frac{d(u^2 - 4)}{du}$$

$$= \frac{1}{2} (u^2 - 4)^{\frac{1}{2}-1} \times 2u$$

$$= \frac{1}{2\sqrt{u^2 - 4}} \times 2u$$

$$= \frac{u}{\sqrt{u^2 - 4}}$$

function $f(u)$ is differentiable at open interval

for all $u \in (2, 4)$

then,

there exists at least a point $c \in (2, 4)$

such that,

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f(a) = f(2) = \sqrt{2^2 - 4} = 0$$

$$f(b) = f(4) = \sqrt{16 - 4} = \sqrt{12} = 2\sqrt{3}$$

Now,

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$\text{or, } \frac{c}{\sqrt{c^2 - 4}} = \frac{2\sqrt{3} - 0}{4 - 2}$$

squaring on both sides

$$\frac{c^2}{c^2 - 4} = \frac{2\sqrt{3}}{2}$$

$$c^2 = 3c^2 - 12$$

$$\text{or, } 2c^2 - 12 = 0$$

$$\text{or, } c^2 - 6 = 0$$

$$\therefore c^2 = 6$$

$$\therefore c = \sqrt{6}$$

$$\therefore c = \pm 2.44$$

$$c = 2.44 \in (2, 4)$$

Hence, Mean Value Theorem verified.

c) $f(u) = Au^2 + Bu + C, u \in [a, b]$

→ Solution.

function $f(u)$ is polynomial function which exists for all $u \in [a, b]$

so it is continuous.

$$f'(u) = 2Au + B$$

function $f(u)$ is differentiable for all $u \in (a, b)$ then,

there exists at least a point $c \in (a, b)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f(a) = f(a) = Aa^2 + Ba + C$$

$$f(b) = Ab^2 + Bb + C$$

$$2Ac + B = \frac{Ab^2 + Bb - (Aa^2 + Ba)}{b - a} - C$$

$$2Abc + Bb - 2Aac - Ba = Ab^2 + Bb + C - Aa^2 - Ba - C$$

$$2Ac(b - a) = A(b^2 - a^2)$$

$$2c = \frac{(b - a)(b + a)}{(b - a)}$$

$$c = \frac{(a + b)}{2} \in (a, b)$$

Hence Mean Value Theorem is verified.

d) $f(u) = e^u, u \in [0, 1]$

→ Solution.

$f(u) = e^u$ is exponential function

which exists for all $u \in [0, 1]$

so it is continuous.

$$f'(u) = e^u$$

function $f(u)$ is differentiable for all $u \in (a, b)$

then,

there exists at ^{least one} ~~constant~~ point $c \in (0, 1)$

such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f(a) = f(0) = e^0 = 1$$

$$f(b) = f(1) = e^1 = 2.71$$

then,

$$e^c = \frac{2.71 - 1}{1 - 0}$$

$$e^c = 1.71$$

$$c = \ln(1.71)$$

$$c = 0.53 \in (0, 1)$$

Hence Mean Value Theorem is verified.

e) $f(u) = \log u$, $u \in [1, e]$

→ Solution,

function $f(u)$ is logarithm function which exists for all $u \in [1, e]$. So it is continuous.

$$f'(u) = \frac{d \log u}{du} = \frac{1}{u}$$

which is differentiable for all $u \in (1, e)$

then,

there exists at least a point $c \in (1, e)$

such that,

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$\therefore f(a) = f(1) = \log(1) = 0$$

$$\therefore f(b) = f(e) = \log(e) = \log(2.71) = 0.996$$

$$\therefore f'(c) = \frac{f(b) - f(a)}{b-a}$$

$$\text{or, } \frac{1}{c} = \frac{0.996 - 0}{e-1}$$

$$\text{or, } 0.996c = e-1$$

$$\text{or, } 0.996c > 2.71 - 1$$

$$\text{or, } c = \frac{1.71}{0.996} = 1.71 \in (1, e)$$

$$\therefore c = 1.71 \in (1, e)$$

Hence Mean Value Theorem is verified.

$$f: f(u) = u^3 + u^2 - 6u, u \in [-1, 4]$$

→ solution,

function $f(u)$ is polynomial function which exists for all $u \in [-1, 4]$,

so it is continuous.

$$f'(u) = 3u^2 + 2u - 6$$

function $f(u)$ is differentiable for all $u \in (-1, 4)$ then,

there exists at least a point $c \in (-1, 4)$, such that,

$$f'(c) = \frac{f(b) - f(a)}{b-a}$$

$$f(a) = f(-1) = (-1)^3 + (-1)^2 - 6 \times (-1) = -1 + 1 + 6 = 5$$

$$f(b) = f(4) = 4^3 + 4^2 - 6 \times 4 = 64 + 16 - 24 = 56$$

Now,

$$f'(c) = \frac{f(b) - f(a)}{b-a}$$

$$3c^2 + 2c - 6 = \frac{56 - 5}{4 + 1}$$

$$\text{or, } 3c^2 + 2c - 6 = \frac{50}{5}$$

$$\text{or, } 3c^2 + 2c - 16 = 0$$

$$\text{or, } 3c^2 + 8c - 6c - 16 = 0$$

$$\text{or, } c(3c+8) - 2(3c+8) = 0$$

$$\text{or, } (3c+8)(c-2) = 0$$

either

$$c = +2 \in (-1, 4)$$

$$c = -\frac{8}{3}$$

Hence MVT verified.

$$\text{g) } f(u) = (u-1)(u-2)(u-3), u \in [1, 4]$$

→ Solution,

$$\text{function } f(u) = (u-1)(u-2)(u-3)$$

$$= u(u-2) - 1(u-2)(u-3)$$

$$= (u^2 - 2u - u + 2)(u-3)$$

$$= (u^2 - 3u + 2)(u-3)$$

$$= u(u^2 - 3u + 2) - 3(u^2 - 3u + 2)$$

$$= u^3 - 3u^2 + 2u - 3u^2 + 9u - 6$$

$$= u^3 - 6u^2 + 11u - 6$$

function $f(u)$ is polynomial function which exists for all $u \in [1, 4]$, so

It is continuous.

$$f'(u) = 3u^2 - 12u + 11$$

so, it is differentiable at open interval for all $u \in (1, 4)$ then,

there exists at least a point $c \in (1, 4)$

such that

$$\frac{f'(c)}{b-a} = \frac{f(b) - f(a)}{b-a}$$

$$\text{or, } f(b) = f(4) = 4^3 - 6 \times 4^2 + 11 \times 4 - 6 = 6$$

$$f(a) = f(1) = 1^3 - 6 \times 1^2 + 11 \times 1 - 6 = 0$$

Now,

$$3c^2 - 12c + 11 = \frac{6-0}{4-1}$$

$$3c^2 - 12c + 11 = 2$$

$$\text{or } 3c^2 - 12c + 9 = 0$$

$$\text{or } c^2 - 4c + 3 = 0$$

$$\text{or } c^2 - 3c - c + 3 = 0$$

$$\text{or, } c(c-3) - 1(c-3) = 0$$

$$\text{or, } (c-1)(c-3) = 0$$

$$\therefore c = 1, 3$$

$$\text{or } c = 3 \in (1, 4)$$

Hence Mean Value Theorem is verified.

$$\text{h. } f(u) = u + \frac{1}{u}, u \in [\frac{1}{2}, 2]$$

→ Solution,

function $f(u)$ is polynomial function which exists for all $u \in [\frac{1}{2}, 2]$, so it is continuous

$$\begin{aligned} f'(u) &= 1 + (-1)u^{-2} \\ &= 1 - \frac{1}{u^2} \end{aligned}$$

it is differentiable at open interval $u \in (\frac{1}{2}, 2)$ then,

there exists at least a point $c \in (\frac{1}{2}, 2)$ such that,

$$\frac{f(c) - f(b) - f(a)}{b-a}$$

$$\therefore f(a) = f(\frac{1}{2}) = \frac{1}{2} + \frac{1}{\frac{1}{2}} = 0.5 + 2 = 2.5$$

$$\therefore f(b) = f(2) = 2 + \frac{1}{2} = 2 + 0.5 = 2.5$$

$$f(c) = \frac{f(b) - f(a)}{b-a}$$

$$\text{or, } 1 - \frac{1}{c^2} = 2.5 - 2.5$$

$$\text{or, } c^2 - 1 = 0$$

$$\text{or, } c^2 = 1$$

$$\therefore c = \sqrt{1} = \pm 1$$

$$c = +1 \in \left(\frac{1}{2}, 2\right)$$

Hence Mean Value Theorem is Verified.

$$\therefore f(u) = u^{\frac{2}{3}}, u \in [0, 1]$$

→ Solution,

function $f(u)$ is polynomial function which exists for all $u \in [0, 1]$,

so it is continuous.

$$f'(u) = \frac{2}{3} u^{\frac{2}{3}-1}$$

$$= \frac{2}{3} u^{\frac{1}{3}}$$

$$= \frac{2}{3 \sqrt[3]{u}}$$

It is differentiable for all $u \in (0, 1)$.

then,

there exists at least a point $c \in (0, 1)$, it

(Such that)

$$f'(c) = \frac{f(b) - f(a)}{b-a}$$

$$\therefore f(a) = f(0) = 0$$

$$f(b) = f(1) = 1$$

$$\text{or, } \frac{2}{3 \sqrt[3]{c}} = \frac{1-0}{1-0}$$

Cubing on both sides

$$8 = 27c$$

$$\therefore c = \frac{8}{27} = 0.29 \in (0, 1)$$

Hence MVT is Verified.

* Proof for Rolle's Theorem:


→ Function $f(x)$ is continuous in $x \in [a, b]$.

It has greater value

say $f(c) = M$ and

least value say

$f(d) = m$



a. If $M = m$,

then function is constant and $f'(u) = 0$

b. If $M \neq m$, then M is different from either $f(a)$ or $f(b)$. Since it is differentiable.

$$f'(u) = \lim_{h \rightarrow 0} \frac{f(u+h) - f(u)}{h}$$

$$\text{a. } f'(c) = \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h}$$

If $f(c)$ is greater than $f(c+h)$ for +ve and -ve value of h .

If h is positive:

$$\frac{f(c+h) - f(c)}{h} \leq 0$$

If h is negative:

$$\frac{f(c+h) - f(c)}{h} \geq 0$$

taking limit $h \rightarrow 0$ for Right hand Derivatives and LHD.

$$\lim_{h \rightarrow 0^+} \frac{f(c+h) - f(c)}{h} \leq 0, \text{ for RHD}$$

$$\lim_{h \rightarrow 0^-} \frac{f(c+h) - f(c)}{h} \geq 0, \text{ for LHD}$$


Derivatives exists when $\text{RHD} = \text{LHD} = 0$
Hence $f'(c) = 0$

Proved

Geometrical Interpretation: (Rolle's Theorem)

→ In a continuous curve of function $f(u)$, tangent can be drawn from each point of the curve.


There exists at least a point where tangent can be drawn which is parallel to x -axis.



Geometrical Interpretation: (Langrange MVT)

→ In the continuous curve of function $f(u)$, tangent can be drawn from each point of the curve.

There exists at least a point where tangent can be drawn which is parallel to the secant joining the end points of the function.



Proof (Lagrange Mean Value Theorem)

Defining a new function $\Phi(u)$ as

$$\Phi(u) = f(u) + Au \quad \text{--- (i)}$$


where A is a constant to be determined such that,

$$\Phi(a) = \Phi(b)$$

$$\text{or, } f(a) + Aa = f(b) + Ab$$

$$\text{or, } A(a-b) = f(b) - f(a)$$

$$\text{or, } A = -\frac{f(b) - f(a)}{b-a}$$

Putting the value of A in eqn (i),

$$\Phi(u) = f(u) - \frac{f(b) - f(a)}{b-a} u$$

and

$$\Phi'(u) = f'(u) - \frac{f(b) - f(a)}{b-a}$$

Since the function $\Phi(u)$ is continuous in $u \in [a, b]$, differentiable in $u \in (a, b)$ and $\Phi(a) = \Phi(b)$. Then by Rolle's Theorem there exists at least a point $c \in (a, b)$,

such that,

$$\Phi'(c) = 0$$

$$\text{or, } f'(c) - \frac{f(b) - f(a)}{b-a} = 0$$

$$\text{or, } f'(c) = \frac{f(b) - f(a)}{b-a}$$

Hence proved

* Verify Rolle's Theorem:

$$f(u) = u(u+3)e^{-u/2}, u \in [-3, 0]$$

→ Solution,

function $f(u)$ is polynomial function which exists for all $u \in [-3, 0]$.

so it is continuous.

$$f(u) = (u^2 + 3u)e^{-u/2}$$

$$f'(u) = (u^2 + 3u) \frac{d}{du} e^{-u/2} + e^{-u/2} \frac{d}{du} (u^2 + 3u)$$

$$= (u^2 + 3u) \left(-\frac{1}{2}\right) e^{-u/2} + e^{-u/2} (2u + 3)$$

$$= e^{-u/2} \left(\frac{(u^2 + 3u - 1) \cdot (2u + 3)}{2} \right)$$

$$= e^{-u/2} \left(\frac{u^2 + 3u - 2u - 6}{2} \right)$$

$$= \frac{e^{-u/2}}{2} (u^2 - u - 6)$$

which is differentiable for all $u \in (-3, 0)$

then,

$$f(a) = f(-3) = (-3)^2 + 3(-3) e^{-3/2}$$

$$= 9 - 9 e^{-3/2}$$

$$= 0$$

$$f(b) = f(0) = 0$$

$$f(a) = f(b) = 0$$

then there exists at least a point $c \in (-3, 0)$

such that

$$f'(c) = 0$$

$$\frac{e^{-c/2}}{2} (c^2 - c - 6) = 0$$

$$c^2 - c - 6 = 0$$

$$\text{Q. } c^2 - 3c + 2c - 6 = 0$$

$$\text{or } c(c-3) + 2(c-3)$$

$$\text{or } (c-3)(c+2) = 0$$

either,

$$c = +3 \notin (-3, 0)$$

$$c = -2 \in (-3, 0)$$

Hence Rolle's theorem is verified.

Conic Section

1. Parabola ($e=1$)


2. Ellipse ($e < 1$)

3. Hyperbola ($e > 1$)


"Parabola"

$$1. y^2 = 4au$$


eqn of dir $\Rightarrow u = -a$




$$2. y^2 = -4au$$



$$3. u^2 = 4ay$$



47 $x^2 = -4ay$



The important results of parabola are tabulated as,

Equation of Parabola	Vertex	Focus	Equation of directrix	length of latus rect.
1. $y^2 = 4ax$	(0, 0)	(a, 0)	$x = -a$	4a
$y^2 = -4ax$		(-a, 0)	$x = a$	
2. $x^2 = 4ay$	(0, 0)	(0, a)	$y = -a$	4a
$x^2 = -4ay$		(0, -a)	$y = a$	
3. $(y-k)^2 = 4a$ (u-h)	(h, k)	(h+a, k)	$x = h-a$	4a
4. $(u-h)^2 = 4a$ (y-k)	(h, k)	(h, k+a)	$y = k-a$	4a

* Find the focus and directrix of parabola $y^2 = 10x$.

→ Solution,

$$y^2 = 10x$$

Comparing $y^2 = 10x$ with $y^2 = 4ax$

$$\therefore 4a = 10$$

$$\therefore a = \frac{5}{2}$$

$$\therefore \text{Focus} = (a, 0) = \left(\frac{5}{2}, 0\right)$$

$$\text{eqn of directrix} = x + a = 0$$

$$\text{or, } x + \frac{5}{2} = 0$$

$$\therefore 2x + 5 = 0$$

parabola.

2. Find the focus and directrix of parabola.

a) $y^2 = 12u$

→ Solution

Comparing $y^2 = 12u$ with $y^2 = 4au$

$$4a = 12$$

$$\therefore a = 3$$

$$\therefore \text{Focus (F)} = (3, 0)$$

equation of directrix is $u = -a$

$$u + 3 = 0 \quad u$$

b) $u^2 = 6y$

Comparing $u^2 = 6y$ with eqn ~~$u^2 = 4ay$~~

$$\therefore 4a = 6$$

$$\therefore a = \frac{6}{4} = \frac{3}{2}$$

$$\therefore \text{Focus (F)} = (0, \frac{3}{2}) \quad \text{directrix} = y = -\frac{3}{2}$$

$$y + \frac{3}{2} = 0 \quad u$$

c) $u^2 = -8y$

→ Solution,

Comparing $u^2 = -8y$ with $u^2 = -4ay$

$$\therefore 4a = 8$$

$$\therefore a = 2$$

$$\therefore \text{Focus (F)} = (0, -2)$$

$$\text{directrix} = y = 2$$

$$\therefore y - 2 = 0$$

d) $y = -8u^2$

→ Solution,

$$y = -8u^2$$

$$u^2 = -\frac{1}{8}y \quad \dots \text{(i)}$$

Comparing (i) with $u^2 = -4ay$

$$\therefore 4a = \frac{1}{8}$$

$$\therefore a = \frac{1}{32}$$

$$\text{Focus (F)} = (0, -\frac{1}{32})$$

$$\text{directrix (y)} = \frac{1}{32}$$

$$\Rightarrow u = -3y$$

$$\text{a. } y^2 = -\frac{1}{3}u \quad \dots \quad (1)$$

Comparing eqⁿ (1) with $y^2 = -4au$

$$4a = \frac{1}{3} \quad \therefore a = \frac{1}{12} = \frac{1}{12}$$

$$\text{Focus (F)} = \left(-\frac{1}{12}, 0\right)$$

$$\text{directrix (u)} = \frac{1}{3 \times 4} = \frac{1}{12} \#$$

$$\text{b. } u = 2y^2$$

$$\Rightarrow y^2 = \frac{1}{2}u \quad \dots \quad (1)$$

Comparing eqⁿ (1) with $y^2 = 4au$

$$\therefore 4a = \frac{1}{2}$$

$$\therefore a = \frac{1}{8} \quad \text{so Focus} = \left(\frac{1}{8}, 0\right)$$

$$\text{directrix (u)} = -\frac{1}{8}$$

$$\text{g. } (y+3)^2 = 2(u+2)$$

→ Solution,

Comparing $(y+3)^2 = 2(u+2)$ with $(y-k)^2 = 4a(u-h)$

where,

$$h = -2, k = -3$$

$$4a = 2 \quad \therefore a = \frac{1}{2}$$

we have eqⁿ of directrix (u) = h - a

$$\text{or, } u = -2 - \frac{1}{2}$$

$$\text{vertex } (h, k) = (-2, -3)$$

$$= \frac{-9-1}{2}$$

$$\text{Focus (F)} = (h+a, k)$$

$$= -\frac{5}{2} \#$$

$$= \left(-2 + \frac{1}{2}, -3\right)$$

$$= \left(-\frac{3}{2}, -3\right) \#$$

$$h) u^2 + 4u + 4y + 16 = 0$$

→ Solution,

$$u^2 + 2 \cdot u \cdot 2 + (2)^2 - (2)^2 + 4(y+4) = 0$$

$$\therefore (u+2)^2 - 4 + 4(y+4) = 0$$

$$\therefore (u+2)^2 - 4 + 4y + 16 = 0$$

$$\therefore (u+2)^2 + 4y + 12 = 0$$

$$\therefore (u+2)^2 + 4(y+3) = 0 \quad \text{--- (i)}$$

Comparing eqn (i) with $(u-h)^2 = 4a(y-k)$

$$h = -2$$

$$F = (h, k+2)$$

$$k = -3$$

$$4a = -4$$

$$y = k+2$$

$$\therefore a = -1$$

$$\text{Focus } (F) = (h, k+2)$$

$$= (-2, -3-1)$$

$$= (-2, -4)$$

$$\text{eqn of directrix } (y) = k-a$$

$$y = -3+1 = -2$$

* Find the equation of Parabola:

$$a) \text{ vertex } (0,0), \text{ focus } (-4,0)$$

→ Solution,

Since y-coordinates of vertex and focus are equal, so it is parallel to x-axis

here,

$$V(0,0), F(-4,0) \text{ is of } y^2 = 4au$$

$$\therefore a = -4$$

Now

$$\text{eqn of Parabola} = y^2 = 4au$$

$$y^2 = 4 \times (-4)u$$

$$\boxed{y^2 = -16u}$$

b) vertex $(0,0)$, eqⁿ of directrix $(y) = 2$

→ solution,

eqⁿ of directrix $(y) = 2$

so,

$$y = +a \quad \therefore a = 2$$

$$u^2 = 4ay$$

$$u^2 = -4 \times 2y$$

$u^2 + 8y = 0$ is req eqⁿ of parabola

c) vertex $(-2,0)$, eqⁿ of directrix $(u) = 2$

→ solution,

vertex $(h,k) = (-2,0)$ eqⁿ of directrix $(u) = 2$ so eqⁿ of P is $(y-k)^2$

we have,

$$(y-k)^2 = 4a(u-h)$$

$$\therefore (y-0)^2 = 4 \times a(u+2)$$

For a

we have,

eqⁿ of directrix $(u) = \underline{\underline{h-a}}$

$$2 = -2 - a$$

$$a = 4$$

∴ the required eqⁿ of parabola is $y^2 = -4 \times 4(u+2)$

$$y^2 = -16u - 32$$

$$\therefore y^2 + 16u + 32 = 0$$

d) vertex $(3,2)$, and end of latusrectum $(5,6)$ and $(5,-2)$.


→ solution:


Mid point of latus rectum
is Focus. so

$$\text{Mid-point} = \frac{u_1+u_2}{2}, \frac{y_2+y_1}{2}$$

$$= \frac{5+5}{2}, \frac{6-2}{2}$$

$$= 5, 2$$





Focus $(5, 2)$ vertex $(3, 2)$

Since y -coordinate of the vertex and focus are equal. So the axis is parallel to x -axis. $\therefore h = 3$ and $(u-h)^2 = 4a(y-k)$

here $(h, k) = (3, 2)$ $y = 5$ and $(y-2)^2 = 4a(y-2)$
and $(h+a, k) = (5, 2)$

$$\therefore h+a = 5$$

$$a = 5 - 3 = 2$$

Now,

Eqⁿ of parabola is $(y-2)^2 = 4a(u-3)$

$$(y-2)^2 = 4a(u-3)$$

$$y^2 - 4y + 4 = 4 \times 2(u-3)$$

$$y^2 - 4y + 4 = 8u - 24$$

$$y^2 - 4y - 8u + 28 = 0$$

which is req. eqⁿ of parabola.

Assignment: 3

Attempt all questions from 2065-2072 'read only'

Year 2065

$$\text{Group 'A'} = 2 \times 10 = 20$$

1. Verify Rolle's theorem for the function $f(u) = \frac{u^3}{3} - 3u$ on interval $[-3, 3]$.

→ Solution:

Given function $f(u)$ is polynomial function which exist for all $u \in [-3, 3]$

so it is continuous.

$$f'(u) = \frac{1}{3} \times 3u^2 - 3$$

$$\therefore f'(u) = u^2 - 3$$

so it is differentiable at open interval $u \in (-3, 3)$ then,

$$f(a) = f(-3) = -\frac{3^3}{3} + 3 \times 3 = -9 + 9 = 0$$

$$f(b) = f(3) = \frac{3^3}{3} - 9 = 9 - 9 = 0$$

$$\therefore f(a) = f(b) = 0$$

then,

there exists at least a point $c \in (-3, 3)$, such that

$$f'(c) = 0$$

$$c^2 - 3 = 0$$

$$\therefore c^2 = 3$$

$$\therefore c = \sqrt{3}$$

$$\therefore c = 1.73 \in (-3, 3)$$

Hence Rolle's theorem is verified.

2. Find the eccentricity of the hyperbola $9u^2 - 16y^2 = 144$

→ Solution,

$$9u^2 - 16y^2 = 144$$

$$\text{or, } \frac{9u^2}{144} - \frac{16y^2}{144} = 1$$

$$\frac{u^2}{16} + \frac{y^2}{9} = 1$$

Comparing $\frac{u^2}{(4)^2} + \frac{y^2}{(3)^2} = 1$ with $\frac{u^2}{a^2} + \frac{y^2}{b^2} = 1$

$$\therefore a = 4, b = 3$$

Now,

we have,

$$\text{eccentricity of hyperbola} = \sqrt{1 + \frac{b^2}{a^2}}$$

$$\therefore e = \sqrt{1 + \frac{9}{16}} = \sqrt{\frac{16+9}{16}}$$

$$e = \sqrt{\frac{25}{16}} = \frac{5}{4}$$

∴ eccentricity of hyperbola is $\frac{5}{4}$.

3. Find the area enclosed by the curve $r^2 = 4 \cos 2\theta$

→ Solution

$$r^2 = 4 \cos 2\theta$$

$$\theta = \frac{\pi}{4}$$

we have,

$$\text{Area of curve} = \int \frac{1}{2} r^2 d\theta$$

$$\theta_1 = \frac{\pi}{4}$$

$$= \int_{0}^{\frac{\pi}{4}} \frac{1}{2} 4 \cos 2\theta d\theta$$

$$\theta_2 = \frac{\pi}{4}$$


$$= 2 \int_{0}^{\frac{\pi}{4}} \cos 2\theta d\theta$$

$$= 2 \left[\frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{4}}$$

$$= 2 \times \left[\frac{\sin 2 \cdot \frac{\pi}{4}}{2} - \frac{\sin 2 \cdot 0}{2} \right]$$

$$= 2 \times \left[\frac{1}{2} - \frac{0}{2} \right]$$

$$= 1 \text{ sq. unit}$$



∴ The total area enclosed by curve $= 1 \times 4 = 4$ sq. unit.

Year 2066

Group A: $2 \times 10 = 20$

1. Find the length of curve $y = u^{3/2}$ from $u=0$ to $u=4$.

→ Solution,

$$y = u^{3/2}$$

$$\frac{dy}{du} = \frac{3}{2} u^{3/2 - 1} \quad \therefore 1 + \left(\frac{dy}{du} \right)^2 = 1 + \left(\frac{3\sqrt{u}}{2} \right)^2$$

$$= \frac{3}{2} u^{1/2}$$

$$= \frac{3\sqrt{u}}{2}$$

$$u=4$$

$$\text{Length (L)} = \int_{u=0}^{u=4} \sqrt{1 + \left(\frac{dy}{du} \right)^2} du$$

$$= \int_{u=0}^{u=4} \sqrt{1 + \frac{9u}{4}} du$$

$$= \int_{u=0}^{u=4} \left(1 + \frac{9u}{4} \right)^{1/2} du$$

$$= \left[\frac{\left(1 + \frac{9u}{4} \right)^{1/2 + 1}}{\left(\frac{1}{2} + 1 \right) \cdot \frac{9}{4}} \right]_{u=0}^{u=4}$$

$$= \frac{\left(1 + \frac{9u}{4} \right)^{3/2}}{\frac{3}{2} \times}$$

5.10- Equation of an ellipse

	Centre	Vertex	Focus	Eccentricities	Length of latus rectum
1. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ $a > b > 0$	(0,0)	($\pm a, 0$)	($\pm ae, 0$)	$e = \sqrt{1 - \frac{b^2}{a^2}}$	$\frac{2b^2}{a}$
2. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ $b > a > 0$	(0,0)	(0, $\pm b$)	(0, $\pm be$)	$e = \sqrt{1 - \frac{a^2}{b^2}}$	$\frac{2a^2}{b}$
3. $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$ $a > b > 1$	(h,k)	($h \pm a, k$)	($h \pm ae, k$)	$e = \sqrt{1 - \frac{b^2}{a^2}}$	$\frac{2b^2}{a}$
4. $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$ $b > a > 1$	(h,k)	($h, k \pm b$)	($h, k \pm be$)	$e = \sqrt{1 - \frac{a^2}{b^2}}$	$\frac{2a^2}{b}$

Hyperbola

5.10- Equation of a Hyperbola

	Centre	Vertex	Focus	Trans. axis	Conj. axis	Eccentricity
1. $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$	($\frac{a^2}{b^2}$, 0)	($\pm a, 0$)	($\pm ae, 0$)	2a	2b	$e = \sqrt{1 + \frac{b^2}{a^2}}$
2. $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ $\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$	(0,0)	(0, $\pm b$)	(0, $\pm be$)	2b	2a	$e = \sqrt{1 + \frac{a^2}{b^2}}$
3. $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$	(h,k)	($h \pm a, k$)	($h \pm ae, k$)	2a	2b	$e = \sqrt{1 + \frac{b^2}{a^2}}$


Polar Equation

$$\cos \theta = \frac{x}{r}$$

$$\therefore x = r \cos \theta$$

$$\sin \theta = \frac{y}{r}$$

$$\therefore y = r \sin \theta$$



$$P(x, y)$$

Polar coordinates

$$x^2 + y^2 = r^2 \cos^2 \theta + r^2 \sin^2 \theta$$

$$x^2 + y^2 = r^2 (\cos^2 \theta + \sin^2 \theta)$$

$$\therefore r^2 = x^2 + y^2$$

$$\frac{y}{x} = \frac{r \sin \theta}{r \cos \theta}$$

$$\therefore \tan \theta = \frac{y}{x}$$

$$\therefore \theta = \tan^{-1} \left(\frac{y}{x} \right)$$

Cartesian equation: (x, y)

Find Cartesian eqⁿ from polar eqⁿ

$$r \sin(\theta + \frac{\pi}{6}) = 2$$

$$\text{on } r \{ \sin \theta \cdot \cos \frac{\pi}{6} + \sin \frac{\pi}{6} \cdot \cos \theta \} = 2$$

$$\text{a. } r \left(\sin \theta \cdot \frac{\sqrt{3}}{2} + \frac{1}{2} \cos \theta \right) = 2$$

$$\sin(C + D) = \sin C \cdot \cos D + \cos C \cdot \sin D$$

$$\text{a. } r \sin \theta \sqrt{3} + r \cos \theta = 4$$

$$\therefore x + y\sqrt{3} = 4$$




Fig: Polar eqⁿ

1) $r = \frac{ke}{1 + e \cos \theta}$

and $x = k$ is the eqⁿ of directrix.

2) $r = \frac{ke}{1 - e \cos \theta}$

and $x = -k$ is the eqⁿ of directrix.

3) $r = \frac{ke}{1 + e \sin \theta}$

and $y = k$ is the equation of directrix.

4) $r = \frac{ke}{1 - e \sin \theta}$

and $y = -k$ is the eqⁿ of directrix.


* find Polar eqⁿ of $e=2$ $n=2$

$e=1$, so parabola

$n=k=2$

\therefore Polar eqⁿ = $r = \frac{ke}{1 + e \cos \theta} = \frac{2 \times 1}{1 + 1 \cos \theta} = \frac{2}{1 + \cos \theta}$

$e=1 \rightarrow$ Parabola
 $e > 1 \rightarrow$ Hyperbola
 $e < 1 \rightarrow$ Ellipse

"Cylindrical coordinates"

→ Cylindrical co-ordinates represent a point 'P' in Space where

a) r and α are polar co-ordinates for the vertical projection of P on u, y plane.

b) z is the rectangular vertical coordinate.

here,

$$1. u = r \cos \alpha \quad \tan \alpha = \frac{y}{u}$$

$$2. y = r \sin \alpha \quad 4. \alpha = \tan^{-1} \left(\frac{y}{u} \right)$$

$$3. z = z \quad 5. u^2 + y^2 = r^2$$

* Convert into cylindrical system eqⁿ.

$$9) u^2 + y^2 + (z - \frac{1}{2})^2 = \frac{1}{4}$$

$$\Rightarrow u^2 + y^2 + z^2 - 2z \frac{1}{2} + \frac{1}{4} = \frac{1}{4}$$

$$\text{or, } u^2 + y^2 + z^2 - 2z = 0$$

$$\text{or, } u^2 + y^2 + z^2 = 2z \neq 0$$

$$\cos \alpha = ar$$

$$r \cos \alpha = ar^2$$

$$u = ar^2$$

$$u = a(u^2 + y^2)$$

$$\therefore u = au^2 + ay^2$$

→ convert rect. coordinates $(1, 0, 0)$ in cylindrical coordinates

→ rect. coordinates $= (u, y, z) = (1, 0, 0)$ $\sqrt{1^2 + 0^2 + 0^2}$.

$$r^2 = u^2 + y^2 =$$

$$r = \sqrt{u^2 + y^2} = 1$$

$$\tan \alpha = \frac{y}{u}$$

$$\therefore \alpha = \tan^{-1} \left(\frac{0}{1} \right) = 0,$$

spherical co-ordinates




fig: spherical coordinates

→ Spherical coordinates represent a point P on the space by ordered triples (s, ϕ, θ) where,

a) s is a distance from P to origin (0).

b) ϕ is the angle that \vec{OP} makes with +ve z axis.

c) θ is the angle from cylindrical coordinates.

$$\cos \phi = \frac{z}{s}$$

$$\sin \phi = \frac{r}{s}$$

$$\therefore z = s \cos \phi$$

$$\therefore r = s \sin \phi$$

$$1. u = r \cos \theta = s \sin \phi \cdot s \cos \phi \quad * \text{Find Sp. co eqn}$$

$$2. y = r \sin \theta = s \sin \phi \cdot s \sin \phi \quad \text{a) } u^2 + y^2 + (z-1)^2 = 1$$

$$3. z = s \cos \phi$$

$$\text{or } u^2 + y^2 + z^2 \rightarrow 2z + 1 = 1$$

$$4. s^2 = u^2 + y^2 + z^2$$

$$\text{or } s^2 - 2s \cos \phi = 0$$

$$5. \tan \theta = \left(\frac{y}{u} \right)$$

$$\text{or } s(s - 2 \cos \phi) = 0$$

$$\therefore s = 0$$

$$s = 2 \cos \phi$$

Vector And Vector Valued Function

Dot Product: $\vec{a} = (a_1, a_2, a_3), \vec{b} = (b_1, b_2, b_3)$

$$\vec{a} \cdot \vec{b} = (a_1 \vec{i} + a_2 \vec{j} + a_3 \vec{k}) \cdot (b_1 \vec{i} + b_2 \vec{j} + b_3 \vec{k})$$


$$= (a_1 b_1 \vec{i} + a_2 b_2 \vec{j} + a_3 b_3 \vec{k})$$

Cross Product: $\vec{a} (a_1, a_2, a_3) \vec{b} (b_1, b_2, b_3)$

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \vec{i}(0-0) + \vec{j}(0-0) + \vec{k}(a_1 b_2 - a_2 b_1)$$


$$= (a_1 b_2 - a_2 b_1) \vec{k}$$

* Angle betⁿ two Vectors :-



$$\cos \alpha = \frac{\vec{u} \cdot \vec{v}}{|\vec{u}| |\vec{v}|}$$

Orthogonal Vectors :



$$\vec{u} \cdot \vec{v} = 0$$

Unit Vector = $\frac{\vec{v}}{|\vec{v}|}$

Q. Find angle betⁿ $\vec{u} = 4\vec{i} - 2\vec{j} - \vec{k}$ and $\vec{v} = 4\vec{i} - 2\vec{j} + 4\vec{k}$

$$\rightarrow |\vec{u}| = \sqrt{16+4+1} = \sqrt{21}$$

$$|\vec{v}| = \sqrt{16+4+16} = \sqrt{36} = 6$$

Now,

$$\cos \alpha = \frac{\vec{u} \cdot \vec{v}}{|\vec{u}| |\vec{v}|} = \frac{(4\vec{i} - 2\vec{j} - \vec{k}) \cdot (4\vec{i} - 2\vec{j} + 4\vec{k})}{6 \times \sqrt{21}} = \frac{16+4-4}{6\sqrt{21}}$$

$$\therefore \alpha = \cos^{-1} \left(\frac{8}{3\sqrt{21}} \right) = 54.41^\circ$$

* Find the measure of triangle whose vertices are $A(-1, 0)$, $B(2, 1)$ and $C(1, -2)$.

$$\angle A = ?$$

$$\rightarrow \cos \angle A = \frac{\vec{AB} \cdot \vec{AC}}{|\vec{AB}| \cdot |\vec{AC}|}$$

$$\vec{AB} = \vec{OB} - \vec{OA} = (2, 1) - (-1, 0) = (3, 1)$$

$$|\vec{AB}| = \sqrt{9+1} = \sqrt{10}$$

$$\vec{AC} = \vec{OC} - \vec{OA} = (1, -2) - (-1, 0) = (2, -2)$$

$$|\vec{AC}| = \sqrt{4+4} = \sqrt{8}$$

$$\therefore \cos \angle A = \frac{\vec{AB} \cdot \vec{AC}}{|\vec{AC}| \cdot |\vec{AB}|} = \frac{(3, 1) \cdot (2, -2)}{\sqrt{10} \times \sqrt{8}} = \frac{6-2}{\sqrt{80}}$$

$$\therefore \angle A = \cos^{-1} \left(\frac{4}{\sqrt{80}} \right) = 63.43^\circ$$


Similarly,

$$\text{find } \cos \angle B = \frac{\vec{BA} \cdot \vec{BC}}{|\vec{BA}| \cdot |\vec{BC}|} = 53.13^\circ$$

$$\cos \angle C = \frac{\vec{CA} \cdot \vec{CB}}{|\vec{CA}| \cdot |\vec{CB}|} = 63.43^\circ$$


* Cross Vector:

$$|\vec{u} \times \vec{v}| = |\vec{u}| \cdot |\vec{v}| \sin \theta$$



$$\therefore \sin \theta = \frac{|\vec{u} \times \vec{v}|}{|\vec{u}| |\vec{v}|}$$

Area of parallelogram:



$$\text{Area of parallelogram } (A) = |\vec{a} \times \vec{b}|$$

$$\text{Area of triangle} = \frac{1}{2} |\vec{a} \times \vec{b}|$$

$$\frac{\vec{u} \times \vec{v}}{|\vec{u} \times \vec{v}|}$$


* Find vector perpendicular to plane of $P(1, -1, 0)$,
 $Q(2, 1, -1)$ and $R(-1, 1, 2)$.

→ Solution,

let O be the position vector,

$$\vec{PO} = \vec{OQ} - \vec{OP} = (2, 1, -1) - (1, -1, 0) \\ = (1, 2, -1)$$

$$\vec{PR} = \vec{OR} - \vec{OP} = (-1, 1, 2) - (1, -1, 0) \\ = (-2, 2, 2)$$



Now,

$$\vec{PO} \times \vec{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & -1 \\ -2 & 2 & 2 \end{vmatrix} \quad |\vec{PO} \times \vec{PR}| = \sqrt{36+36} \\ = \sqrt{72}$$

$$= \mathbf{i}(4+2) - \mathbf{j}(2-2) + \mathbf{k}(2+2) \\ = 6\mathbf{i} + 0\mathbf{j} + 6\mathbf{k} \\ = 6\mathbf{i} + 6\mathbf{k}$$

Unit vector of $\vec{PO} \times \vec{PR} = \frac{\vec{PO} \times \vec{PR}}{|\vec{PO} \times \vec{PR}|}$

$$= \frac{6\mathbf{i} + 6\mathbf{k}}{\sqrt{72}}$$

$$= \frac{6\mathbf{i}}{6\sqrt{2}} + \frac{6\mathbf{k}}{6\sqrt{2}}$$

$$= \frac{\mathbf{i}}{\sqrt{2}} + \frac{\mathbf{k}}{\sqrt{2}}$$

* Find area of triangle with vertices $P(4, 2, 0)$, $Q(1, 3, 0)$ and $R(1, 1, 3)$

→ Solution,

let O be the position vector.

$$\vec{PO} = \vec{OQ} - \vec{OP} = (1, 3, 0) - (4, 2, 0) = (-3, 1, 0)$$

$$\vec{PR} = \vec{OR} - \vec{OP} = (1, 1, 3) - (4, 2, 0) = (-3, -1, 3)$$

Now,

$$|\vec{AB}| = |\vec{PO} \times \vec{PR}| = ?$$

$$\vec{PO} \times \vec{PR} = \begin{vmatrix} i & j & k \\ -3 & 1 & 0 \\ -3 & -1 & 3 \end{vmatrix}$$

$$= i(3+0) - j(0+9) + k(-3+3)$$

$$= 3\vec{i} - 9\vec{j} + 0\vec{k}$$

$$|\vec{PO} \times \vec{PR}| = \sqrt{9+81+36}$$

$$= \sqrt{126}$$

$$\therefore \text{Area of triangle} = \frac{\sqrt{126}}{2} \text{ Sq. unit.}$$

* Find area of Parallelogram:

$$a) A(1,0), B(0,1), C(-1,0), D(0,-1)$$

→ Solution,

$$\vec{AB} = (\vec{OB} - \vec{OA}) = (0,1) - (1,0) = (-1,1)$$

$$\vec{AC} = (\vec{OC} - \vec{OA}) = (-1,0) - (1,0) = (-2,0)$$

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} i & j & k \\ -1 & 1 & 0 \\ -2 & 0 & 0 \end{vmatrix}$$

$$= i(0-0) + j(0-0) + k(0+2)$$

$$= 2\vec{k}$$

Now,

$$|\vec{AB} \times \vec{AC}| = \sqrt{2^2} = 2$$

$$\therefore \text{Area of Parallelogram} = |\vec{AB} \times \vec{AC}|$$

$$= 2 \text{ Sq. unit.}$$

OR

$$\vec{AC} \times \vec{AD} \text{ w.}$$

Triple Scale Product (Box Product):

 $|(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}|$ = Volume of box.

$$\vec{U} = U_1 \vec{i} + U_2 \vec{j} + U_3 \vec{k}$$

$$\vec{V} = V_1 \vec{i} + V_2 \vec{j} + V_3 \vec{k}$$

$$\vec{W} = W_1 \vec{i} + W_2 \vec{j} + W_3 \vec{k}$$

$$(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} = \begin{vmatrix} U_1 & U_2 & U_3 \\ V_1 & V_2 & V_3 \\ W_1 & W_2 & W_3 \end{vmatrix}$$

* find the Vol. of box determined by

a) $\vec{U} = \vec{i} + 2\vec{j} - \vec{k}$

$\vec{V} = -2\vec{i} + 3\vec{j}$


$\vec{W} = 7\vec{j} - 4\vec{k}$

$$\rightarrow \text{Vol. of box} = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} = \begin{vmatrix} 1 & 2 & -1 \\ -2 & 0 & 3 \\ 0 & 7 & -4 \end{vmatrix}$$

$$= 1 \left| (0 - 21) - 2(8 - 0) - 1(-4 - 0) \right|$$

$= 1 - 23$

= 23 unit cubed.

Equation of Plane passing through point $P(x_1, y_1, z_1)$ and normal to $\vec{n} = A\vec{i} + B\vec{j} + C\vec{k}$.

$$\therefore \text{Eqn of plane} = A(x - x_1) + B(y - y_1) + C(z - z_1) = 0$$

* find the eqn of plane passing through $A(0, 0, 1)$ $B(2, 0, 0)$ $C(0, 3, 0)$

→ Solution,

$$\vec{AB} = \vec{OB} - \vec{OA} = (2, 0, 0) - (0, 0, 1) = (2, 0, -1)$$

$$\vec{AC} = \vec{OC} - \vec{OA} = (0, 3, 0) - (0, 0, 1) = (0, 3, -1)$$

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 0 & -1 \\ 0 & 3 & -1 \end{vmatrix}$$

$$= \mathbf{i}(0+3) - \mathbf{j}(-2-0) + \mathbf{k}(0-0) \\ = 3\mathbf{i} + 2\mathbf{j} + 0\mathbf{k} \quad \therefore \mathbf{i} = 3$$

$$\text{Point } P = (0, 0, 1) = (u, y, z) \quad \begin{matrix} B=2 \\ C=6 \end{matrix}$$

$$\therefore \text{eqn of Plane} = A(u-0) + B(y-0) + C(z-1) \\ = 3(u-0) + 2(y-0) + 6(z-1) \\ = 3u + 2y + 6z = 6 \quad \#$$

Parametric eqn of a line.

→ Parametric eqn of a line passing through Point $P(u_1, y_1, z_1)$ and Parallel to vector $v = (A\mathbf{i} + B\mathbf{j} + C\mathbf{k})$

∴ Parametric eqn are,

$$u = u_1 + At$$

$$y = y_1 + Bt$$

$$z = z_1 + Ct$$

$$v = (A\mathbf{i} + B\mathbf{j} + C\mathbf{k})$$

* Find the parametric eqn passing through $A(-3, 2, -3)$
 $B(2, -1, 4)$.

→ Solution:

$$\vec{AB} = \vec{OB} - \vec{OA} = (1, -1, 4) - (-3, 2, -3) \\ = (4, -3, 7)$$

$$\therefore A=4, \quad B=-3 \quad C=7$$

$$\text{let line } (u, y, z) = (-3, 2, -3)$$

∴ Parametric eqn is

$$u = u_1 + At = -3 + 4t$$

$$y = y_1 + Bt = 2 - 3t$$

$$z = z_1 + Ct = -3 + 7t$$

when $(u, y, z) = (2, -1, 4)$

$$u = 1 + 4t$$

$$y = -1 - 3t$$

$$z = 4 + 7t$$

* Find the intersecting point where the line $x = \frac{3}{8}t + 2$,
 $y = -2t$ and $z = 1+t$ intersect the plane $3u + 2y + 6z = 6$.

→ Solution,

Substituting the value of u, y, z in eqn of plane,

$$3u + 2y + 6z = 6$$

$$\text{a, } 3\left(\frac{8}{3}t + 2\right) + 2(-2t) + 6(1+t) = 6$$

$$\text{or } 8 + 6t - 4t + 6 + 6t = 6$$

$$\text{or } 8t + 8 = 0$$

$$\text{or } t = -1$$

Now,

$$u = \frac{8}{3}t + 2 = \frac{8}{3} + 2 \times (-1) = \frac{2}{3}$$


$$y = -2t = -2 \times -1 = 2$$

$$z = 1+t = 1-1 = 0$$

∴ Required point is $(\frac{2}{3}, 2, 0)$.

From 2nd term

* Distance from a point S to a line through P parallel to vector \vec{v} .



* find the distance from point $S(1,1,5)$ to the line $u=1+t, y=3-t, z=2t$.
 → Solution,

Comparing $u=1+t, y=3-t, z=2t$ with parametric eqⁿ $u=u_1+At, y=y_1+Bt, z=z_1+Ct$

$$P \therefore (u_1, y_1, z_1) = (1, 3, 0)$$

$$\text{vector Parallel } (\vec{v}) = (A, B, C) = (1, -1, 2) \\ = \vec{i} - \vec{j} + 2\vec{k}$$

Given point $S = (1, 1, 5)$

Now,

$$\text{distance } (d) = \frac{|\vec{PS} \times \vec{v}|}{|\vec{v}|}$$

$$\vec{PS} = \vec{OS} - \vec{OP} = (1, 1, 5) - (1, 3, 0) = (0, -2, 5)$$

$$\therefore d = \frac{|\vec{PS} \times \vec{v}|}{|\vec{v}|}$$

$$\vec{PS} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 0 & -2 & 5 \\ 1 & -1 & 2 \end{vmatrix}$$

$$= \vec{i}(-4+5) - \vec{j}(0-5) + \vec{k}(0+2)$$

$$= \vec{i} + 5\vec{j} + 2\vec{k}$$

$$|\vec{PS} \times \vec{v}| = \sqrt{1+25+4} = \sqrt{30}$$

$$|\vec{v}| = \sqrt{1+1+4} = \sqrt{6}$$

$$\therefore \text{distance } (d) = \frac{|\vec{PS} \times \vec{v}|}{|\vec{v}|} = \frac{\sqrt{30}}{\sqrt{6}} = \sqrt{\frac{30}{6}} = \sqrt{5} \text{ unit}$$

Distance from a point S to a plane.

$$A\vec{i} + B\vec{j} + C\vec{k} \cdot S$$

$$P(u, v, z)$$

$$d = \left| \vec{PS} \cdot \frac{\vec{n}}{|\vec{n}|} \right|$$

where, $\vec{n} = A\vec{i} + B\vec{j} + C\vec{k}$

P = Point lies on the plane (unknown)
to find P

Put $u=0, z=0$ and $y=u$.

* Find the distance from S(1, 1, 3) to plane $3x+2y+6z=6$

→ Solution,

Given S(1, 1, 3)

normal vector $(\vec{n}) = 3\vec{i} + 2\vec{j} + 6\vec{k}$

for point P, $|\vec{n}| = \sqrt{9+4+36} = \sqrt{49} = 7$

when $u=0, z=0$

$$3x0 + 2y + 6x0 = 6$$

$$\therefore 2y = 6 \therefore y = 3$$

∴ Point P(0, 3, 0)

Now,

$$\vec{PS} = \vec{OS} - \vec{OP} = (1, 1, 3) - (0, 3, 0) = \vec{i} - 2\vec{j} + 3\vec{k}$$

$$\text{Then, } \frac{\vec{n}}{|\vec{n}|} = \frac{3\vec{i}}{7} + \frac{2\vec{j}}{7} + \frac{6\vec{k}}{7}$$

Now,

$$\text{distance (an)} = \left| \vec{PS} \cdot \frac{\vec{n}}{|\vec{n}|} \right|$$


$$= \left| \left(\frac{3}{7}\vec{i} - \frac{2}{7}\vec{j} + \frac{6}{7}\vec{k} \right) \cdot (3\vec{i} + 2\vec{j} + 6\vec{k}) \right|$$

$$= \frac{3}{7} \cdot \frac{4}{7} + \frac{9}{7} = \frac{3}{7} + \frac{4}{7} + \frac{18}{7}$$

$$= \frac{25}{7} \text{ unit}$$

(25/7) ~~25/7~~

* Angle between two planes :



$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|}$$

where,

θ = angle betⁿ two planes

\mathbf{n}_1 and \mathbf{n}_2 are normal for two plane

* Find the angle between two planes

$$a. 3x - 6y - 2z = 15 \text{ and } 2x + y - 2z = 5.$$

→ Solution,

normal for two vectors are,

$$\mathbf{n}_1 = 3\hat{i} - 6\hat{j} - 2\hat{k}$$

$$|\mathbf{n}_1| = \sqrt{9+36+4} = \sqrt{49} = 7$$

$$\mathbf{n}_2 = 2\hat{i} + \hat{j} - 2\hat{k}$$

$$|\mathbf{n}_2| = \sqrt{4+1+4} = \sqrt{9} = 3$$

Now,

$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|} = \frac{(3\hat{i} - 6\hat{j} - 2\hat{k}) \cdot (2\hat{i} + \hat{j} - 2\hat{k})}{7 \times 3}$$

$$= \frac{6\hat{i}}{21} - \frac{6\hat{j}}{21} + \frac{4\hat{k}}{21}$$

$$\therefore \cos \theta = \frac{4}{21}$$


$$\therefore \theta = \cos^{-1} \left(\frac{4}{21} \right) = 79.01^\circ$$

∴ Angle betⁿ two plane is 79.01° *

Vector Parallel to the line of intersection of two plane.

$$\text{Formula} = \mathbf{n}_1 \times \mathbf{n}_2$$

$$\boxed{\text{Formula} = \mathbf{n}_1 \times \mathbf{n}_2}$$



* Find the vector parallel to line of intersection of plane $3u - 6y - 2z = 15$ and $2u + y - 2z = 5$.

→ Solution,

$$\text{Normal eqn are, } \mathbf{n}_1 = 3\mathbf{i} - 6\mathbf{j} - 2\mathbf{k}$$

$$\mathbf{n}_2 = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$$

Now,

$$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -6 & -2 \\ 2 & 1 & -2 \end{vmatrix}$$

$$\text{Req. Veet.} = 14\mathbf{i} + 2\mathbf{j} + 15\mathbf{k}$$

Vector Valued Function:

A vector is defined by $\mathbf{r}(t) = f(t)\mathbf{i} + g(t)\mathbf{j} + h(t)\mathbf{k}$ is known as Vector Valued function in t .

Where,

t = Parameters

$f(t), g(t), h(t)$ are components of $\mathbf{r}(t)$.

→ if $\mathbf{r}(t)$ is a position vector, particle moving along a smooth curve in space, then,

(a) Velocity (\mathbf{v}) = $\frac{d\mathbf{r}}{dt}$

(b) Speed (s) = $|\mathbf{v}|$

(c) Acceleration (\mathbf{a}) = $\frac{d\mathbf{v}}{dt}$

* Find particle velocity and acceleration vectors at the given value of t .

① $\mathbf{r}(t) = (t+1)\mathbf{i} + (t^2-1)\mathbf{j} + 2t\mathbf{k}$, $t=1$.

→ Solution,

$$\text{velocity } (\mathbf{v}) = \frac{d\mathbf{r}}{dt}$$

$$= \frac{d}{dt} \{ (t+1)\mathbf{i} + (t^2-1)\mathbf{j} + 2t\mathbf{k} \}$$

$$= 1\mathbf{i} + 2t\mathbf{j} + 2\mathbf{k}$$

when $t=1$

$$\text{velocity } (\mathbf{v}) = \mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$$

$$\text{speed } (s) = |\mathbf{v}| = \sqrt{1+4+4} = \sqrt{9} = 3$$

$$\text{acceleration } (\mathbf{a}) = \frac{d\mathbf{v}}{dt}$$

$$= \frac{d}{dt} (1\mathbf{i} + 2t\mathbf{j} + 2\mathbf{k})$$

$$= 2\mathbf{j}$$

② $\mathbf{r}(t) = (2 \cos t)\mathbf{i} + (3 \sin t)\mathbf{j} + 4t\mathbf{k}$, $t=\pi/2$.

→ Solution,

$$\text{Velocity } (\mathbf{v}) = \frac{d\mathbf{r}}{dt} = -2 \sin t \mathbf{i} + 3 \cos t \mathbf{j} + 4 \mathbf{k}$$

when $t=\pi/2$

$$= -2 \sin(\pi/2) \mathbf{i} + 3 \cos(\pi/2) \mathbf{j} + 4 \mathbf{k}$$

$$= -2 \times 1 \mathbf{i} + 3 \times 0 \mathbf{j} + 4 \mathbf{k}$$

$$\therefore \text{velocity } (\mathbf{v}) = -2\mathbf{i} + 4\mathbf{k}$$

$$\text{acceleration } (\mathbf{a}) = \frac{d\mathbf{v}}{dt} = \frac{d}{dt} (-2 \sin t) \mathbf{i} + 3 \cos t \mathbf{j} + 4 \mathbf{k}$$

$$= -2 \cos t \mathbf{i} + 3 \sin t \mathbf{j}$$

where, $t = \pi/2$

$$\mathbf{a} = -2 \cos \pi/2 \hat{i} - 3 \sin \pi/2 \hat{j} \\ = -3 \hat{j}$$

③ $\mathbf{r}(t) = e^t \hat{i} + \frac{2}{9} e^{2t} \hat{j}$, $t = \ln 3$
 $\rightarrow \text{so in}$

$$\text{velocity} (\mathbf{v}) = \frac{d\mathbf{r}}{dt} = e^t \hat{i} + \frac{4}{9} e^{2t} \hat{j}$$

when $t = \ln 3$

$$\mathbf{v} = e^{\ln 3} \hat{i} + \frac{4}{9} e^{2\ln 3} \hat{j}$$

$$= 3 \hat{i} + \frac{9 \times 4}{9} \hat{j}$$

$$= 3 \hat{i} + 4 \hat{j} \cancel{\#}$$

$$\text{acceleration} (\mathbf{a}) = \frac{d\mathbf{v}}{dt} = e^t \hat{i} + \frac{8}{9} e^{2t} \hat{j}$$

= when $t = \ln 3$

$$\mathbf{a} = e^{\ln 3} \hat{i} + \frac{8}{9} e^{2\ln 3} \hat{j}$$

$$= 3 \hat{i} + \frac{8 \times 9}{9} \hat{j}$$

$$\therefore \mathbf{a} = 3 \hat{i} + 8 \hat{j} \cancel{\#}$$

* Evaluate the integral of vector value function.

i) $\int_0^1 \{ 3t^2 \hat{i} + 2 \hat{j} + (-3) \hat{k} \} dt$

$$= \left[3 \frac{t^3}{3} \right]_0^1 + 2 \left[t \right]_0^1 + \left[\frac{t^2}{2} - 3t \right]_0^1$$

In Calculus

do $\ln 3$

first

then,

$\ln 3$ w

or

$\ln 3$

then $2 \times \ln 3$

after

e^{Ans} w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

e^{Ans}

w

or

$$= (1^3 - 0)\vec{i} + 2(1-0)\vec{j} + \left(\frac{1}{2} - 3 - 0 + 0\right)\vec{k}$$

$$= \vec{i} + 2\vec{j} - \frac{5}{2}\vec{k} \quad \text{W.}$$

$$\text{ii)} \int_{1}^2 \left\{ (6-6t)\vec{i} + 3\sqrt{t}\vec{j} + \frac{4}{t^2}\vec{k} \right\}$$

$$= \left[6t - 6\frac{t^2}{2} \right]_1^2 \vec{i} + 3 \left[\frac{t^{3/2}}{3/2} \right]_1^2 + 4 \int_1^2 t^{-2} \vec{k}$$

$$= \left(12 - 6 \times \frac{4}{2} - \left(6 + \frac{6}{2} \right) \right) \vec{i} + \frac{3}{\frac{1}{2}} \left(2^{3/2} - 1 \right) \vec{j} + 4 \left[\frac{t^{-2+1}}{-2+1} \right]_1^2 \vec{k}$$

$$= -3\vec{i} + 3.65\vec{j} - 4 \left[\frac{1}{2} - 1 \right] \vec{k}$$

$$= -3\vec{i} + 3.65\vec{j} + 2\vec{k} \quad \text{W.}$$

$$4.) \int_{1}^4 \left[\frac{1}{t}\vec{i} + \frac{1}{5-t}\vec{j} + \frac{1}{2t}\vec{k} \right] dt$$

$$\rightarrow \left[\ln t \vec{i} \right]_1^4 + \left[\frac{1}{-(5-t)} \vec{j} \right]_1^4 + \left[\frac{1}{2} \vec{k} \right] \ln t$$

$$= (\ln 4 - \ln 1)\vec{i} - \left[\ln (4-5) \right]_1^4 \vec{j} + \frac{1}{2} \left[\ln t \right]_1^4 \vec{k}$$

$$= \ln \left(\frac{4}{1} \right) \vec{i} - (\ln 1 - \ln (-4)) \vec{j} + \frac{1}{2} (\ln 4 - \ln 1) \vec{k}$$

$$= \ln 4 \vec{i} + (\ln (-4) - \ln (-1)) \vec{j} + \frac{1}{2} (\ln \frac{4}{1}) \vec{k}$$

$$= \ln 4 \vec{i} + \ln \left(-\frac{4}{1} \right) \vec{j} + \frac{1}{2} \ln 4 \vec{k}$$

$$= \ln 4 \vec{i} + \ln 4 \vec{j} + \frac{1}{2} \ln 4 \vec{k} \quad \text{W.}$$

Length of a smooth curve (Arc length formula)


Length of smooth curve is

$$r(t) = f(t) \vec{i} + g(t) \vec{j} + h(t) \vec{k}, \quad a \leq t \leq b$$

$$\therefore L = \int_{t=a}^{t=b} \sqrt{\left(\frac{df}{dt}\right)^2 + \left(\frac{dg}{dt}\right)^2 + \left(\frac{dh}{dt}\right)^2} dt$$

$$L = \int_{t=a}^{t=b} |v| dt$$

$$\text{where } v = \frac{dr}{dt}$$



* find the length of 1 turn of helix.

$$r(t) = (\cos t) \vec{i} + (\sin t) \vec{j} + t \vec{k}$$

→ Solution,

$$v = \frac{dr}{dt} = \frac{d}{dt} \{ (\cos t) \vec{i} + (\sin t) \vec{j} + t \vec{k} \}$$

$$= -\sin t \vec{i} + \cos t \vec{j} + \vec{k}$$

$$|v| = \sqrt{(-\sin t)^2 + (\cos t)^2 + 1^2}$$

$$= \sqrt{\sin^2 t + \cos^2 t + 1}$$

$$= \sqrt{1+1} = \sqrt{2}$$

$$\therefore \text{length of helix} = \int_{t=0}^{t=2\pi} \sqrt{2} dt$$

$$= \sqrt{2} \left[t \right]_0^{2\pi}$$

$$= \sqrt{2} [2\pi - 0]$$

$$= 2\pi \sqrt{2} \text{ unit}$$

* Find the length of indicated portion of curve:

① $r(t) = (2 \cos t) \vec{i} + (2 \sin t) \vec{j} + \sqrt{5} \vec{k}, 0 \leq t \leq \pi$

→ Solution,

we have, $\text{Length } (L) = \int_0^\pi |V| dt$

$$V = \frac{dr}{dt} = (2 \sin t) \vec{i} + (2 \cos t) \vec{j} + \sqrt{5} \vec{k}$$

$$\begin{aligned} |V| &= \sqrt{(-2 \sin t)^2 + (2 \cos t)^2 + (\sqrt{5})^2} \\ &= \sqrt{4(\sin^2 t + \cos^2 t) + 5} \\ &= \sqrt{9} = 3 \end{aligned}$$

$$\therefore L = \int_0^\pi 3 dt = 3 [t]_0^\pi = 3\pi \neq$$

∴ Required length is 3π unit.

② $r(t) = (6 \sin 2t) \vec{i} + (6 \cos 2t) \vec{j} + 5 \vec{k}, 0 \leq t \leq \pi$

→ Solution,

we have, $\text{Length } (L) = \int_0^\pi |V| dt$

$$\begin{aligned} V = \frac{dr}{dt} &= (6 \times 2 \cos 2t) \vec{i} + (-6 \times 2 \sin 2t) \vec{j} + 0 \vec{k} \\ &= (12 \cos 2t) \vec{i} + (-12 \sin 2t) \vec{j} + 0 \vec{k} \end{aligned}$$

$$\begin{aligned} |V| &= \sqrt{144 \cos^2 2t + 144 \sin^2 2t + 0} \\ &= \sqrt{144 + 25} = \sqrt{169} = 13 \end{aligned}$$

$$\therefore L = \int_0^\pi 13 dt$$

$$= 13 [t]_0^\pi$$

$$= 13\pi \text{ unit}$$

∴ Required length of curve is 13π unit.

$$\textcircled{c} \quad r(t) = t \vec{i} + \frac{2}{3} t^{\frac{3}{2}} \vec{k}, \quad 0 \leq t \leq 8.$$

→ Solution,

$$\text{length } (l) = \int_0^8 |v| dt$$

$$v = \frac{dr}{dt} = \vec{i} + \frac{2}{3} \times \frac{3}{2} t^{\frac{3}{2}-1} \vec{k}$$

$$= \vec{i} + t^{\frac{1}{2}} \vec{k}$$

$$|v| = \sqrt{(1)^2 + (t^{\frac{1}{2}})^2} = (1+t) \frac{1}{2}$$

$$\vec{i} + \frac{2}{3} \times \frac{3}{2} t^{\frac{3}{2}-1}$$

$$\vec{i} + \sqrt{t}$$

$$|v| = \sqrt{(1)^2 + (t^{\frac{1}{2}})^2} = \sqrt{1+t}$$

$$\therefore \text{length} = \int_0^8 (1+t)^{\frac{1}{2}} dt = \frac{1}{\frac{1}{2}+1} (1+t)^{\frac{1}{2}}$$

$$= \left[\frac{[1+t]^{\frac{1}{2}+1}}{(\frac{1}{2}+1) \times 1} \right]_0^8$$

$$= \left[\frac{(1+t)^{\frac{3}{2}}}{\frac{3}{2} \times 1} \right]_0^8$$

$$= \frac{2}{3} \left[\left[\frac{(1+t)^{\frac{3}{2}}}{\frac{3}{2}} \right] \right]_0^8$$

$$= \frac{2}{3} \left[\left(\frac{9}{2} - 1 \right) \right]$$

$$= \frac{2}{3} \left\{ \left(\frac{9}{2} - 1 \right) \right\}$$

$$= \frac{2}{3} \left[\frac{7}{2} \right]$$

$$= \frac{2}{3} \left\{ 3^{\frac{3}{2}} - 1 \right\}$$

$$= \frac{2}{3} (26)$$

$$= \frac{2}{3} (27 - 1)$$

$$= \frac{2 \times 26}{3}$$

$$= 17.33 \text{ unit}$$

$$v = \frac{dr}{dt} = \frac{d \cos 3t}{dt} \vec{i} + \frac{d \sin 3t}{dt} \vec{k}$$

$$= 3 \cos 3t \vec{i} + (-3 \sin 3t) \vec{k}$$

∴ Required length is 17.33 unit.

$$4) \quad r(t) = (\cos^3 t) \vec{j} + (\sin^3 t) \vec{k}, \quad 0 \leq t \leq \frac{\pi}{2}$$

→ Solution,

$$\text{length} = \int_0^{\frac{\pi}{2}} |v| dt$$

$$v = \frac{dr}{dt} = \frac{d \cos^3 t}{dt} \vec{j} + \frac{d \sin^3 t}{dt} \vec{k}$$

$$= 3 \cos^2 t (-\sin t) \vec{j} + 3 \sin^2 t (\cos t) \vec{k}$$

$$\begin{aligned}
 \mathbf{v} = \frac{d\mathbf{r}}{dt} &= \frac{d(\cos^3 t)}{dt} \mathbf{j} + \frac{d(\sin^3 t)}{dt} \mathbf{j} \\
 &= \left(\frac{d \cos^3 t}{dt} \times \frac{d \cos t}{dt} \right) \mathbf{j} + \left(\frac{d \sin^3 t}{dt} \times \frac{d \sin t}{dt} \right) \mathbf{j} \\
 &= 3 \cos^2 t \cdot (-\sin t) \mathbf{j} + 3 \sin^2 t \cdot \cos t \mathbf{j}
 \end{aligned}$$

$$|\mathbf{v}| = \sqrt{9 \cos^4 t \cdot \sin^2 t + 9 \sin^4 t \cdot \cos^2 t}$$

$$\begin{aligned}
 &= \sqrt{9 \cos^2 t \cdot \sin^2 t (\cos^2 t + \sin^2 t)} \\
 &= 3 \sin t \cdot \cos t \\
 &= \frac{3}{2} 2 \sin t \cdot \cos t
 \end{aligned}$$

$$= \frac{3}{2} \sin 2t$$

$$L = \int_0^{\pi/2} \frac{3}{2} \sin 2t \, dt$$

$$= \frac{3}{2} \left[-\frac{\cos 2t}{2} \right]_0^{\pi/2}$$

$$= \frac{3}{4} [-\cos 2 \cdot \pi/2 + \cos 2 \cdot 0]$$

$$= \frac{3}{4} (-1 + 1)$$

$$= \frac{3}{4} \times 2$$

$$= \frac{3}{2} \text{ unit}$$

∴ Required length is $\frac{3}{2}$ unit.

$$5. \mathbf{r}(t) = t \mathbf{i} + \frac{\sqrt{6}}{2} t^2 \mathbf{j} + t^3 \mathbf{k}, -1 \leq t \leq 1$$

→ Solution,

$$\text{Length (L)} = \int |\mathbf{v}| dt$$

$$\mathbf{v} = \frac{d\mathbf{r}}{dt} = \mathbf{i} + \frac{\sqrt{6}}{2} \times 2t \mathbf{j} + 3t^2 \mathbf{k}$$

$$|V| = \sqrt{1+6t^2+9t^4} = \sqrt{(1+3t^2)^2} = (1+3t^2)$$

$$L = \int_{-1}^1 (1+3t^2) dt$$

$$= [t]_{-1}^1 + 3 \left[\frac{t^3}{3} \right]_{-1}^1 = [1 - (-1)] + \frac{3}{3} (1^3 - (-1)^3) \\ = (1+1) + (1+1) = 2+2 = 4 \text{ unit.}$$

∴ Required length of curve is 4 unit.

$$\boxed{\text{Unit Tangent Vector (T)} = \frac{V}{|V|}}$$

→ It determines the direction of the motion of curve.

$$T = \frac{V}{|V|}, \text{ where } V = \frac{dr}{dt}$$

$$\boxed{\text{Curvature: } k = \frac{1}{|V|} \left| \frac{dT}{dt} \right|}$$

The rate at which the unit tangent vector (T) turns per unit length along the curve is known as curvature. i.e.

$$k = \frac{1}{|V|} \left| \frac{dT}{dt} \right|$$

$$\text{where } T = \text{unit tangent} = \frac{V}{|V|}$$

* Show that circle of radius a is $\frac{1}{a}$.

→ Solution,

always for circle

we have,

$$r(t) = a(\cos t) \hat{i} + a(\sin t) \hat{j}$$

$$r(t) = a(\cos t) \hat{i} + a(\sin t) \hat{j}$$

$$V = \frac{dr}{dt} = (-a \sin t) \hat{i} + a \cos t \hat{j}$$

$$|V| = \sqrt{a^2 \sin^2 t + a^2 \cos^2 t} = \sqrt{a^2} = a$$

Again,

$$\text{Unit Vector (T)} = \frac{V}{|V|} = \frac{(-a \sin t) \hat{i} + a \cos t \hat{j}}{a} \\ = -\sin t \hat{i} + \cos t \hat{j}$$

Again,

$$\text{Curvature } (k) = \frac{1}{|V|} \left| \frac{dT}{dt} \right|$$

$$\frac{dT}{dt} = -\cos t \vec{i} - \sin t \vec{j}$$

$$\left| \frac{dT}{dt} \right| = \sqrt{\cos^2 t + \sin^2 t} = 1$$

$$\therefore k = \frac{1}{|V|} \left| \frac{dT}{dt} \right| = \frac{1}{a} \times 1 = \frac{1}{a}$$

\therefore The radius of curvature is a .

* Find the Curvature of helix:

$$r(t) = (a \cos t) \vec{i} + (a \sin t) \vec{j} + bt \vec{k}, a, b \geq 0, a^2 + b^2 \neq 0.$$

→ Solution,

$$V = \frac{dr}{dt} = (-a \sin t) \vec{i} + (a \cos t) \vec{j} + b \vec{k}$$

$$|V| = \sqrt{a^2 \sin^2 t + a^2 \cos^2 t + b^2} \\ = \sqrt{a^2 (\sin^2 t + \cos^2 t) + b^2} = \sqrt{a^2 + b^2}$$

$$\text{Unit tangent Vector } (T) = \frac{V}{|V|} = \frac{(-a \sin t) \vec{i} + (a \cos t) \vec{j} + b \vec{k}}{\sqrt{a^2 + b^2}}$$

$$\frac{dT}{dt} = \frac{(-a \cos t) \vec{i} - (a \sin t) \vec{j} + 0}{\sqrt{a^2 + b^2}}$$


$$\therefore \text{Curvature } (k) = \frac{1}{|V|} \left| \frac{dT}{dt} \right|$$

$$\left| \frac{dT}{dt} \right| = \sqrt{\frac{a^2 \cos^2 t + a^2 \sin^2 t}{(\sqrt{a^2 + b^2})^2}} = \sqrt{\frac{a^2}{a^2 + b^2}} = \frac{a}{\sqrt{a^2 + b^2}}$$

$$\therefore k = \frac{1}{|V|} \left| \frac{dT}{dt} \right|$$

$$= \frac{1}{\sqrt{a^2 + b^2}} \times \frac{a}{\sqrt{a^2 + b^2}} = \frac{a}{a^2 + b^2}$$

\therefore Required Curvature of Helix is $\frac{a}{a^2 + b^2}$.



* Length of a smooth curve (Arc length formula):

Length of smooth curve is,


$$r(t) = f(t)\hat{i} + g(t)\hat{j} + h(t)\hat{k}, \quad a \leq t \leq b$$

$$\therefore L = \int_{t=a}^{t=b} \sqrt{\left(\frac{df}{dt}\right)^2 + \left(\frac{dg}{dt}\right)^2 + \left(\frac{dh}{dt}\right)^2} dt$$

OR

$$\text{Length} = \int_{t=a}^{t=b} |v| dt$$

$$\text{where, } v = \frac{dr}{dt}$$



* Unit tangent Vector: (T)

It determines the direction of the motion of the curve. It is denoted by T and given by,

$$T = \frac{v}{|v|}, \text{ where, } v = \frac{dr}{dt}$$

* Curvature: (k)

The rate at which the unit tangent vector (T) turns per unit length along the curve is known as curvature.

$$k = \frac{1}{|v|} \left| \frac{dt}{dt} \right|$$

$$k = \frac{|v \times a|}{|v|^3}$$

where,

$$T = \text{unit vector} = \frac{v}{|v|}$$

$$\text{for circle } r(t) = a(\cos t)\hat{i} + a(\sin t)\hat{j}$$

* Curvature of a straight line:-

The curvature of straight line is zero because the unit tangent vector (T) always points in same direction, so its components are equal.

$$k = \left| \frac{dT}{ds} \right| = \frac{1}{|v|} \left| \frac{dT}{dt} \right|$$

when,

$$\frac{dT}{dt} = 0 \quad \therefore k = 0$$

* Principal Unit Normal (N):-

$$N = \frac{\frac{dT}{dt}}{\left| \frac{dT}{dt} \right|}$$

$$T = \frac{v}{|v|} \quad N = \frac{dT}{dt}$$

Defn:-

Let $r(t)$ be a differentiable vector valued function and $T(t)$ be unit T.V. The P.U.N. = $\frac{\frac{dT}{dt}}{\left| \frac{dT}{dt} \right|}$

* Find principle Unit Normal Vector and Curvature.

a) $r(t) = (\cos 2t) \vec{i} + (\sin 2t) \vec{j}$

$$v = \frac{dr}{dt} = (-2 \sin 2t) \vec{i} + (2 \cos 2t) \vec{j}$$

$$\begin{aligned} |v| &= \sqrt{4 \sin^2 2t + 4 \cos^2 2t} \\ &= \sqrt{4} \\ &= 2 \end{aligned}$$

Now,

$$T = \frac{v}{|v|} = \frac{(-2\sin 2t)\vec{i}}{2} + \frac{(2\cos 2t)\vec{j}}{2}$$

$$T = (-\sin 2t)\vec{i} + \frac{\cos 2t}{2}\vec{j}$$

$$\frac{dT}{dt} = (-2\cos 2t)\vec{i} + (-2\sin 2t)\vec{j}$$

$$\left| \frac{dT}{dt} \right| = \sqrt{4\cos^2 2t + 4\sin^2 2t} = \sqrt{4} = 2$$

Now,

$$\text{Principal Unit Normal vector } (N) = \frac{\frac{dT}{dt}}{\left| \frac{dT}{dt} \right|} =$$

$$= \frac{(-2\cos 2t)\vec{i}}{2} + \frac{(-2\sin 2t)\vec{j}}{2}$$

$$\therefore N = (-\cos 2t)\vec{i} - (\sin 2t)\vec{j}$$

$$\text{Curvature } (k) = \frac{1}{|v|} \left| \frac{dT}{dt} \right| = \frac{1}{2} \times 2 = 1 \text{ unit} \#$$

$$\text{b) } r(t) = (3\sin t)\vec{i} + (3\cos t)\vec{j} + 4t\vec{k}$$

$$\rightarrow N = (-\sin t)\vec{i} - (\cos t)\vec{j}$$

$$k = \frac{3}{25} \#$$

$$\text{c) } r(t) = (2t + 3)\vec{i} + (5 - t^2)\vec{j}$$

solution,

$$v = \frac{dr}{dt} = 2\vec{i} - 2t\vec{j}, |v| = \sqrt{4 + 4t^2} = 2\sqrt{1+t^2}$$

$$\text{Now, } T = \frac{v}{|v|} = \frac{2\vec{i} - 2t\vec{j}}{2\sqrt{1+t^2}}$$

$$= \frac{2(\vec{i} - t\vec{j})}{2\sqrt{1+t^2}^{1/2}}$$

$$\frac{dT}{dt} = \left(\frac{d(1+t^2)^{-1/2}}{d(1+t^2)} \times \frac{d(1+t^2)}{dt} \right) \vec{i} - \frac{d(1+t^2)^{-1/2}}{dt} \vec{j} +$$

Rough:-

1st part

$$\frac{d(1+t^2)^{-1/2}}{d(1+t^2)} \times \frac{d(1+t^2)}{dt} = -\frac{1}{2} (1+t^2)^{-3/2} \cdot 2t = -\frac{t}{(1+t^2)^{3/2}}$$

2nd Part,

$$\frac{d\left(\frac{1}{(1+t^2)^{-1/2}}\right)}{dt} \quad \boxed{\frac{u}{v} = \sqrt{\frac{du}{dv}} - \frac{u}{v} \frac{dv}{du}}$$

$$= (1+t^2)^{1/2} \frac{dt}{dt} - + \frac{d(1+t^2)^{1/2}}{dt}$$

$$(1+t^2)^{1/2}$$

$$= (1+t^2)^{1/2} - + \frac{1}{2} (1+t^2)^{-1/2} \times 2t / \sqrt{2}$$

$$= (1+t^2)^{1/2} - +^2 (1+t^2)^{-1/2} / \sqrt{2}$$

$$= (1+t^2)^{1/2} - \frac{1^2}{(1+t^2)^{1/2}} / (1+t^2)^2$$

$$= \frac{(1+t^2) - 1^2}{(1+t^2)^{1/2}}$$

$$= \frac{1+t^2 - 1^2}{(1+t^2)^{1/2} \cdot (1+t^2)} = \frac{1}{(1+t^2)^{3/2}}$$

$$\therefore \frac{dT}{dt} = -\frac{t}{(1+t^2)^{3/2}} \vec{i} - \frac{1}{(1+t^2)^{3/2}} \vec{j}$$

$$|\frac{dT}{dt}| = \sqrt{\left\{ -\frac{t}{(1+t^2)^{3/2}} \right\}^2 + \left\{ -\frac{1}{(1+t^2)^{3/2}} \right\}^2}$$

$$\begin{aligned}
 &= \sqrt{t^2(1+t^2)^{-3} + (1+t^2)^{-3}} \\
 &= \sqrt{\frac{t^2}{(1+t^2)^3} + \frac{1}{(1+t^2)^3}} \\
 &= \sqrt{\frac{t^2+1}{(1+t^2)^3}} \\
 &= \sqrt{(t^2+1)^{1-3}} \\
 &= \sqrt{(t^2+1)^{-2}} \\
 &= \frac{1}{t^2+1} \quad \because \left| \frac{dT}{dt} \right| = \frac{1}{1+t^2}
 \end{aligned}$$

Now,

$$\begin{aligned}
 N &= \frac{dT}{dt} \\
 \frac{dT}{dt} &= -\frac{t}{\sqrt{1+t^2}} \vec{i} - \frac{1}{\sqrt{1+t^2}} \vec{j}
 \end{aligned}$$

$$\therefore \text{Curvature } (k) = \frac{1}{2} (1+t^2)^{3/2}$$

Unit Binomial Vector And Torsion
 Unit Binomial Vector:-




fig: TNB frame

$$B = T \times N$$

Unit Binormal Vector: \mathbf{B}

* \mathbf{TNB} are mutually orthogonal to each other and it defines a moving right handed vector frame that play a significant role in calculation paths of particles moving through space. It is called as frenet frame \mathbf{TNB} frame. $\mathbf{B} = \mathbf{T} \times \mathbf{N}$

Torsion (τ):

Torsion measures how the curve twist. It is negative scalar product of change of binormal with respect to Arc length and principle Unit normal Vector.

$$\tau = -\frac{d\mathbf{B}}{ds} \cdot \mathbf{N} \quad \text{OR}$$

$$\therefore \tau = -\frac{\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \dddot{x} & \dddot{y} & \dddot{z} \end{vmatrix}}{|\mathbf{v} \times \mathbf{a}|^2}$$

$$\dot{x} = 1^{\text{st}} \text{ derivative} = \frac{du}{dt}$$

$$\ddot{x} = 2^{\text{nd}} \text{ derivative} = \frac{d^2u}{dt^2}$$

$$\dddot{x} = 3^{\text{rd}} \text{ derivative} = \frac{d^3u}{dt^3}$$

$$\mathbf{r}(t) = u(t) \mathbf{i} + y(t) \mathbf{j} + z(t) \mathbf{k}$$

$$\dot{x} = \frac{du}{dt}, \dot{y} = \frac{dy}{dt}, \dot{z} = \frac{dz}{dt}$$

$$\text{Similarly } \ddot{x} = \frac{d^2u}{dt^2}, \ddot{y} = \frac{d^2y}{dt^2}$$

Alternative,

$$\text{Curvature } (k) = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3}$$

Q. Find the Curvature and Torsion for:

$$a) r(t) = (a \cos t) \vec{i} + (a \sin t) \vec{j} + bt \vec{k}, a, b \geq 0, a^2 + b^2 \neq 0.$$

→ Solution,

$$v = \frac{dr}{dt} = (-a \sin t) \vec{i} + (a \cos t) \vec{j} + b \vec{k}$$

$$a = \frac{dv}{dt} = (-a \cos t) \vec{i} + (-a \sin t) \vec{j}$$

$$|v| = \sqrt{(-a \sin t)^2 + (a \cos t)^2 + b^2} = \sqrt{a^2(\sin^2 t + \cos^2 t) + b^2} \\ = \sqrt{a^2 + b^2}$$

Now,

$$v \times a = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -a \sin t & a \cos t & b \\ -a \cos t & -a \sin t & 0 \end{vmatrix}$$

$$= \vec{i} \begin{pmatrix} a \cos t & b \\ -a \sin t & 0 \end{pmatrix} - \vec{j} \begin{pmatrix} -a \sin t & b \\ -a \cos t & 0 \end{pmatrix} + \vec{k} \begin{pmatrix} -a \sin t & a \cos t \\ -a \cos t & -a \sin t \end{pmatrix} \\ = (b \sin t) \vec{i} - (a b \cos t) \vec{j} + (a^2 \sin^2 t + a^2 \cos^2 t) \vec{k} \\ = (a b \sin t) \vec{i} - (a b \cos t) \vec{j} + a^2 \vec{k}$$

Again,

$$|v \times a| = \sqrt{(a b \sin t)^2 + (-a b \cos t)^2 + (a^2)^2} \\ = \sqrt{a^2 b^2 (\sin^2 t + \cos^2 t) + a^4} \\ = \sqrt{a^2 (a^2 + b^2)} \\ = a \sqrt{a^2 + b^2}$$

$$\therefore \text{Curvature} = \frac{|v \times a|}{|v|^3} = \frac{a \sqrt{a^2 + b^2}}{(\sqrt{a^2 + b^2})^3} = \frac{a}{(\sqrt{a^2 + b^2})^{3-1}} = \frac{a}{a^2 + b^2}.$$

here,

$$x = a \cos t$$

$$y = a \sin t$$

$$z = bt$$

we have,

$$\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \dddot{x} & \ddot{y} & \ddot{z} \end{vmatrix} = \begin{vmatrix} -a \sin t & a \cos t & b \\ -a \cos t & -a \sin t & 0 \\ a \sin t & -a \cos t & 0 \end{vmatrix}$$

$$= b \begin{vmatrix} -a \cos t & -a \sin t & -0+0 \\ a \sin t & -a \cos t & 0 \end{vmatrix}$$

$$= b (a^2 \cos^2 t + a^2 \sin^2 t)$$

$$= a^2 b$$

$$\therefore \text{Torsion } (\tau) = \frac{\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \dddot{x} & \ddot{y} & \ddot{z} \end{vmatrix}}{|v \times a|^2}$$

$$= \frac{a^2 b}{(a\sqrt{a^2+b^2})^2}$$

$$= \frac{ab}{2(\sqrt{a^2+b^2})^2} \#$$

$$= \frac{b}{a^2+b^2} \#$$

$$\text{b)} \quad r(t) = (a \cos t) \vec{i} + (a \sin t) \vec{j} + at \vec{k}$$

→ Solution,

$$v = \frac{dr}{dt} = (-a \sin t) \vec{i} + (a \cos t) \vec{j} + a \vec{k}$$

$$a = \frac{dv}{dt} = (-a \cos t) \vec{i} + (-a \sin t) \vec{j}$$

$$|v| = \sqrt{a^2 \sin^2 t + a^2 \cos^2 t + a^2}$$

$$= \sqrt{2a^2}$$

$$v \times a = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -a \sin t & a \cos t & a \\ -a \cos t & -a \sin t & 0 \end{vmatrix}$$

$$= \vec{i} (a^2 \sin t) - \vec{j} (a^2 \cos t) + \vec{k} (a^2 \sin^2 t + a^2 \cos^2 t)$$

$$= (a^2 \sin t) \vec{i} - (a^2 \cos t) \vec{j} + a^2 \vec{k}$$

$$|v \times a| = \sqrt{a^4 \sin^2 t + a^4 \cos^2 t + a^4}$$

$$= \sqrt{2a^4}$$

$$= a^2 \sqrt{2}$$

$$\text{Curvature } (k) = \frac{|v \times a|}{|v|^3} = \frac{a^2 \sqrt{2}}{2a^3 \sqrt{2}} = \frac{a^2 \sqrt{2}}{2^3 \cdot 2 \cdot \sqrt{2}}$$

$$= \frac{a^2 \sqrt{2}}{(2\sqrt{2})^3}$$

$$= \frac{a^2 \sqrt{2}}{2^3 \cdot 2\sqrt{2}}$$

$$= \frac{1}{2a} \#$$

Now,

$$x = a \cos t, \quad y = a \sin t, \quad z = at$$

$$\begin{vmatrix} x & y & z \\ x' & y' & z' \\ x'' & y'' & z'' \end{vmatrix} = \begin{vmatrix} -a \sin t & a \cos t & a \\ -a \cos t & -a \sin t & 0 \\ a \sin t & -a \cos t & 0 \end{vmatrix}$$

$$= a \begin{vmatrix} -a \cos t & -a \sin t \\ a \sin t & -a \cos t \end{vmatrix} - 0 + 0$$

$$= a (a^2 \cos^2 t + a^2 \sin^2 t)$$

$$= a^3$$

Now,

$$T = \frac{a^3}{(a^2 \sqrt{2})^2} = \frac{a^3}{a^4 \cdot 2} = \frac{1}{2a} \#$$

$$\Rightarrow \mathbf{r}(t) = (3\sin t)\mathbf{i} + (3\cos t)\mathbf{j} + 4t\mathbf{k}$$

→ Solution,

$$\mathbf{v} = \frac{d\mathbf{r}}{dt} = (3\cos t)\mathbf{i} + (-3\sin t)\mathbf{j} + 4\mathbf{k}$$

$$\mathbf{a} = \frac{d\mathbf{v}}{dt} = (-3\sin t)\mathbf{i} + (-3\cos t)\mathbf{j} + \mathbf{0}$$

$$\begin{aligned} |\mathbf{v}| &= \sqrt{(3\cos t)^2 + (3\sin t)^2 + (4)^2} \\ &= \sqrt{9(\cos^2 t + \sin^2 t) + 16} \\ &= \sqrt{25} \end{aligned}$$

$$|\mathbf{v} \times \mathbf{a}| = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3\cos t & -3\sin t & 4 \\ -3\sin t & -3\cos t & 0 \end{vmatrix}$$

$$\begin{aligned} &= \mathbf{i} (12\cos t) - \mathbf{j} (12\sin t) + \mathbf{k} (-9\cos^2 t - 9\sin^2 t) \\ &= (12\cos t)\mathbf{i} - (12\sin t)\mathbf{j} - 9(\cos^2 t + \sin^2 t)\mathbf{k} \\ &= (12\cos t)\mathbf{i} - (12\sin t)\mathbf{j} - 9\mathbf{k} \end{aligned}$$

$$\begin{aligned} |\mathbf{v} \times \mathbf{a}| &= \sqrt{144\cos^2 t + 144\sin^2 t + 81} \\ &= \sqrt{144(\cos^2 t + \sin^2 t) + 81} = \sqrt{225} = 15 \end{aligned}$$

Now,

$$\text{curvature } (k) = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3} = \frac{15}{(5)^3} = \frac{15}{125} = 0.12$$

$$u = 3\sin t \quad z = 4t$$

$$y = 3\cos t$$

$$\therefore \begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \ddot{\dot{x}} & \ddot{\dot{y}} & \ddot{\dot{z}} \end{vmatrix} = \begin{vmatrix} 3\cos t & -3\sin t & 4 \\ -3\sin t & -3\cos t & 0 \\ -3\cos t & 3\sin t & 0 \end{vmatrix}$$

$$= 4 \begin{vmatrix} -3\sin t & -3\cos t \\ -3\cos t & 3\sin t \end{vmatrix} - 0 + 0$$

$$\begin{aligned} &= 4(-9\sin^2 t - 9\cos^2 t) \\ &= 4 \times (-9)(\sin^2 t + \cos^2 t) = -36 \end{aligned}$$

$$\therefore \text{Torsion } (\tau) = \frac{\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \dddot{x} & \dddot{y} & \dddot{z} \end{vmatrix}}{|\mathbf{v} \times \mathbf{a}|^2} = \frac{-36}{(15)^2} = \frac{-36}{225} = -0.16 \text{ #}$$

Also find unit binormal vector,

$$\mathbf{B} = \mathbf{T} \times \mathbf{N}$$

$$\mathbf{r}(t) = (3\sin t)\mathbf{i} + (3\cos t)\mathbf{j} + 4t\mathbf{k}$$

→ Solution,

$$\mathbf{v} = \frac{d\mathbf{r}}{dt} = (3\cos t)\mathbf{i} + (-3\sin t)\mathbf{j} + 4\mathbf{k}$$

$$|\mathbf{v}| = \sqrt{9\cos^2 t + 9\sin^2 t + 16} = \sqrt{9+16} = \sqrt{25} = 5$$

Now,

$$\text{Unit tangent Vector } (\mathbf{T}) = \frac{\mathbf{v}}{|\mathbf{v}|} = \frac{(3\cos t)\mathbf{i} + (-3\sin t)\mathbf{j} + 4\mathbf{k}}{5}$$

Again,

$$\frac{d\mathbf{T}}{dt} = \frac{(-3\sin t)\mathbf{i} + (-3\cos t)\mathbf{j}}{5}$$

$$\left| \frac{d\mathbf{T}}{dt} \right| = \sqrt{\frac{9\sin^2 t + 9\cos^2 t}{25}} = \sqrt{\frac{9}{25}} = \frac{3}{5}$$

∴ Unit binormal vector = $\mathbf{T} \times \mathbf{N}$

$$\mathbf{N} = \frac{\frac{d\mathbf{T}}{dt}}{\left| \frac{d\mathbf{T}}{dt} \right|} = \frac{(-3\sin t)\mathbf{i} + (-3\cos t)\mathbf{j} \times \frac{5}{3}}{\frac{3}{5}} = (-\sin t)\mathbf{i} + (-\cos t)\mathbf{j}$$

$$\therefore \mathbf{B} = \mathbf{T} \times \mathbf{N} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{3\cos t}{5} & -\frac{3\sin t}{5} & \frac{4}{5} \\ -\sin t & -\cos t & 0 \end{vmatrix}$$

$$= (4\cos t)\mathbf{i} - (4\sin t)\mathbf{j} + \mathbf{k}(-3\cos^2 t - 3\sin^2 t)$$

$$= (4\cos t)\mathbf{i} - (4\sin t)\mathbf{j} - \frac{3}{5}\mathbf{k}(\cos^2 t + \sin^2 t)$$

$$\therefore \text{Unit binormal vector} = (4\cos t)\mathbf{i} - (4\sin t)\mathbf{j} - \frac{3}{5}\mathbf{k} \text{ #}$$

$$d\vec{r}(t) = (e^t \cos t) \vec{i} + (e^t \sin t) \vec{j} + 2 \vec{k}$$

→ Solution,

$$\begin{aligned} \frac{d(e^t \cos t)}{dt} &= e^t \cdot \frac{d \cos t}{dt} + \cos t \cdot \frac{de^t}{dt} \\ &= e^t (-\sin t) + \cos t \cdot e^t \\ &= e^t (\cos t - \sin t) \end{aligned}$$

$$\begin{aligned} \frac{d(e^t \sin t)}{dt} &= e^t \cdot \frac{d \sin t}{dt} + \sin t \cdot \frac{de^t}{dt} \\ &= e^t \cdot \cos t + \sin t \cdot e^t \\ &= e^t (\sin t + \cos t) \end{aligned}$$

$$\begin{aligned} v = \frac{d\vec{r}}{dt} &= e^t (\cos t - \sin t) \vec{i} + e^t (\sin t + \cos t) \vec{j} + 0 \\ &= e^t (\cos t - \sin t) \vec{i} + e^t (\sin t + \cos t) \vec{j} \end{aligned}$$

$$\begin{aligned} \frac{d(e^t (\cos t - \sin t))}{dt} &= e^t (-\sin t - \cos t) + (\cos t - \sin t) e^t \\ &= e^t (-\sin t - \cos t + \cos t - \sin t) \\ &= e^t (-2 \sin t) \end{aligned}$$

$$\begin{aligned} \frac{d(e^t (\sin t + \cos t))}{dt} &= e^t (\cos t - \sin t) + (\sin t + \cos t) \cdot e^t \\ &= e^t (\cos t - \sin t + \sin t + \cos t) \\ &= e^t (2 \cos t) \end{aligned}$$

$$a = \frac{dv}{dt} = e^t (-2 \sin t) \vec{i} + e^t (2 \cos t) \vec{j}$$

$$\begin{aligned} |v| &= \sqrt{(e^t)^2 (\cos t - \sin t)^2 + (e^t)^2 (\sin t + \cos t)^2} \\ &= e^t \sqrt{\cos^2 t - 2 \cos t \cdot \sin t + \sin^2 t + \sin^2 t + 2 \sin t \cdot \cos t + \cos^2 t} \\ &= e^t \sqrt{2(\cos^2 t + \sin^2 t)} \\ &= e^t \sqrt{2} \end{aligned}$$

$$v \times a = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ e^t (\cos t - \sin t) & e^t (\sin t + \cos t) & 0 \\ e^t (-2 \sin t) & e^t (2 \cos t) & 0 \end{vmatrix}$$

$$\begin{aligned}
 &= (et)^2 (2\cos^2 t - 2\sin t \cdot \cos t) - (et)^2 (-2\sin^2 t - 2\sin t \cdot \cos t) \\
 &= (et)^2 \{ 2\cos^2 t - 2\sin t \cos t + 2\sin^2 t + 2\sin t \cos t \} \\
 &= (et)^2 2 (\sin^2 t + \cos^2 t) \\
 &= 2(et)^2.
 \end{aligned}$$

Now,

$$\text{curvature } (k) = \frac{|V \times a|}{|V|^3}$$

$$= \frac{2e^2 t^2}{e^{3t^3} (\sqrt{2})^3}$$

$$= \frac{2}{et^2 \sqrt{2}}$$

$$= \frac{1}{et \sqrt{2}} \#$$

$$x = e^t \cos t, y = e^t \sin t, z = 2$$

$$\begin{vmatrix} x & y & z \end{vmatrix} = \begin{vmatrix} e^t (\cos t - \sin t) & e^t (\sin t + \cos t) & 0 \\ e^t (-2\sin t) & e^t (2\cos t) & 0 \\ e^t (-2\cos t) - (2\sin t) & e^t (2\cos t) - (2\sin t) & 0 \end{vmatrix}$$

$$= 0$$

$$\therefore \text{Now, Torsion } (T) = \frac{0}{(2(et)^2)^2}$$

$$= 0 \#$$

Formulae from Vector Valued function.

1.> Velocity (v) = $\frac{dr}{dt}$

2.> Speed (s) = $|v|$

3.> Acceleration (a) = $\frac{dv}{dt}$

4.> Length (L) = $\int_{t=a}^{t=b} |v| dt$

5.> Unit Tangent Vector (T) = $\frac{v}{|v|}$

6.> Curvature (k) = $\frac{1}{|v|} \left| \frac{dT}{dt} \right|$

or, Curvature (k) = $\frac{|v \times a|}{|v|^3}$

7.> Principle Unit Normal (N) = $\frac{\frac{dT}{dt}}{\left| \frac{dT}{dt} \right|}$

8.> Unit Binormal Vector (B) = $T \times N$

9.> Torsion (τ) =
$$\frac{\begin{vmatrix} \ddot{x} & \ddot{y} & \ddot{z} \\ \ddot{\ddot{x}} & \ddot{\ddot{y}} & \ddot{\ddot{z}} \\ \ddot{\ddot{\ddot{x}}} & \ddot{\ddot{\ddot{y}}} & \ddot{\ddot{\ddot{z}}} \end{vmatrix}}{|v \times a|^2}$$

\ddot{x} = first der.
 $\ddot{\ddot{x}}$ = 2nd der.
 $\ddot{\ddot{\ddot{x}}}$ = 3rd der.

10. Radius of curvature (R) = $\frac{1}{k}$

Vector Formula:

i) $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$

ii) $\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|}$

iii) Unit Vect = $\frac{\vec{v}}{|\vec{v}|}$

iv) Unit Vect = $\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$

v) Area of parallelogram = $|\vec{a} \times \vec{b}|$

vi) Equation of plane = $A(u-u_1) + B(y-y_1) + C(z-z_1) = 0$

vii) Parametric eqn ; $u = u_1 + At$ $y = y_1 + Bt$ $z = z_1 + Ct$

viii) Distance from s to line (d) = $|\vec{ps} \times \vec{v}|$

ix) Distance from s to a plane (d) = $|\vec{ps} \cdot \frac{n}{|n|}|$

x) Angle between two plane,

$\cos \theta = \frac{n_1 \cdot n_2}{|n_1| |n_2|}$

xi) Vector parallel to line = $n_1 \times n_2$

xii) Polar equation

$u = r \cos \theta$ $y = r \sin \theta$ $r^2 = u^2 + y^2$

$\theta = \tan^{-1} (y/u)$

xiii) $r = \frac{ke}{1+e \cos \theta}$, $x = k$ xiv) $r = \frac{ke}{1-e \sin \theta}$, $y = -k$

xv) Cylindrical co-ordinates = (r, θ, z)

xvi) Spherical coordinates = (ρ, ϕ, θ) , $\rho^2 = u^2 + y^2 + z^2$, $z = \rho \cos \phi$

$\tan \phi = \frac{y}{u}$

$u = \rho \sin \phi \cdot \cos \theta$
 $y = \rho \sin \phi \cdot \sin \theta$

Infinite Series:

Expression in the form of

$a_1 + a_2 + a_3 + \dots + a_n + \dots$ is called infinite series, where, a_n is n^{th} term of infinite series.

Sequence form;

$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

⋮
⋮

$$S_n = a_1 + a_2 + a_3 + \dots + a_n$$

$$\therefore S_n = \sum_{n=1}^n a_n$$

is a sequence of partial sum.

where,

S_n is a n^{th} partial sum.

Convergent Series:

If the sequence of the partial sum of series converges to the certain value or limit, then this series is known as convergent series.

$$a_1 + a_2 + a_3 + \dots + a_n + \dots = \sum_{n=1}^{\infty} a_n = L$$

$$\text{or, } \sum_{n=1}^{\infty} a_n = 0, \text{ it is called divergent series.}$$

Test:

- i) Geometric Series Test
- ii) Ratio test
- iii) Root Test
- iv) Limit Comparison test
- v) Integral test.

* Geometric Series test :-

$$a + ar + ar^2 + \dots + ar^{n-1} + \dots$$

$$\text{Common ratio (r)} = \frac{t_2}{t_1} = \frac{ar}{a} = r$$

when,

$$|r| < 1$$

then the Series converges to

$$S_{\infty} = \frac{a}{1-r}$$

$$|r| > 1 \text{ or infinite,}$$

then the series is diverges.

Q. Test whether the given series are convergent or divergent.

$$1> \sum_{n=0}^{\infty} \frac{(-1)^n}{4^n}$$

→ Solution,

$$\text{series} = 1 - \frac{1}{4} + \frac{1}{4^2} - \dots + \infty$$

$$\text{common ratio (r)} = \frac{t_2}{t_1} = -\frac{1}{4} = -\frac{1}{4}$$

$$|r| = \frac{1}{4} < 1, \text{ so series is converges.}$$

$$\text{Sum of series (S}_{\infty}\text{)} = \frac{1}{1 + \frac{1}{4}} = \frac{4}{5} = 0.8$$

$$2> \sum_{n=1}^{\infty} \frac{7}{4^n}$$

$$\rightarrow \text{series} = \frac{7}{4} + \frac{7}{4^2} + \frac{7}{4^3} + \dots + \infty$$

$$\therefore \text{common diff (r)} = \frac{t_2}{t_1} = \frac{7}{4^2} \times \frac{4}{7} = \frac{1}{4} \quad \left| \text{sum (S}_{\infty}\text{)} = \frac{1}{1 - r} \right.$$

$$|r| = \frac{1}{4} < 1, \text{ so it is convergent}$$

$$= \frac{7}{4} \times \frac{4}{3} = \frac{7}{3} \neq .$$

$$3) \sum_{n=0}^{\infty} (-1)^n \frac{5}{4^n}$$

$$\rightarrow \text{Series} = +5 - \frac{5}{4} + \frac{5}{4^2} + \dots + \infty$$

$$r = \frac{t_2}{t_1} = -\frac{5}{4} \times \frac{1}{5} = -\frac{1}{4}$$

$|r| = \frac{1}{4} < 1$, so it is convergent

$$\text{Sum of series (S}_{\infty}) = \frac{a}{1-r} = \frac{5}{1+\frac{1}{4}} = \frac{5}{\frac{5}{4}} = 4 \neq .$$

$$4) \sum_{n=1}^{\infty} 9^{-n+2} \cdot 4^{n+1}$$

$$\rightarrow \text{Series} = 9 \cdot 4^2 + 9 \cdot 4^3 + 9^{-1} \cdot 4^4 + \dots + \infty$$

$$r = \frac{t_2}{t_1} = \frac{64}{144} = 0.44$$

$$-1|r| = 0.44 < 1.$$

$$\text{So, it is convergent, sum} = \frac{a}{1-r} = \frac{144}{1-0.44} = \frac{144}{0.56} = 259.2$$

$$5) \sum_{n=6}^{\infty} (-1)^n \frac{(2^{n+3})}{3^n}$$

$$\rightarrow \text{Series} = \frac{2^9}{3^6} - \frac{2^{10}}{3^7} + \frac{2^{11}}{3^8} - \dots + \infty$$

$$|r| = \left| \frac{t_2}{t_1} \right| = \left| -\frac{2^{10}}{3^7} \times \frac{3^6}{2^9} \right| = \frac{2^{10-9}}{3^{7-6}} = \frac{2}{3} = 0.66$$

$$= 7464.96$$

$|r| < 1$, so Converged 1119744

$$S_{\infty} = \frac{a}{1-r} = \frac{0.702}{1+0.66} = 0.42 \neq .$$

$$6. \sum_{n=1}^{\infty} \frac{2^{n+1} + 9^{n+2}}{5^n}$$

$$\rightarrow \text{series} = \frac{2^2 + 9^3}{5} + \frac{5^3 + 9^4}{5^2} + \frac{5^4 + 9^5}{5^3} + \dots$$

$$t_1 = \frac{2^2 + 9^3}{5} = 146.6$$

$$r = \frac{t_2}{t_1} = \frac{267.44}{146.6} = 1.82$$

$|r| > 1$, so it is divergent.

$$7. \sum_{n=0}^{\infty} \left(\frac{1}{2^n} + \frac{(-1)^n}{5^n} \right)$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2^n} \right) + \sum_{n=0}^{\infty} \frac{(-1)^n}{5^n}$$

$$= \left(1 + \frac{1}{2} + \frac{1}{2^2} + \dots \right) + \left(1 - \frac{1}{5} + \frac{1}{5^2} - \dots \right)$$

$$\text{common difference } (r_1) = \frac{1}{2} \times 1 = \frac{1}{2}$$

$$\text{common diff. } (r_2) = -\frac{1}{5} \times 1 = -\frac{1}{5}$$

$|r_1| < 1$, so it is convergent.

$|r_2| < 1$, so it is also convergent

$$\text{sum}(S_{\infty 1}) = \frac{a_1}{1-r_1} = \frac{1}{1-\frac{1}{2}} = \frac{1}{1} \times \frac{2}{1} = 2$$

$$\text{sum}(S_{\infty 2}) = \frac{a_1}{1-r_2} = \frac{1}{1+\frac{1}{5}} = \frac{1}{1.2} = 0.83$$

$$\therefore \text{sum}(S_{\infty}) = 2 + 0.83 = 2.83$$

$$8. \sum_{n=0}^{\infty} \frac{(-4)^{3^n}}{5^{n-1}}$$

$$\text{Series} = 1 - \frac{64}{1} + \frac{4096}{5} + \dots \infty$$

$r = -\frac{64}{1} = -64$ $|r| = 64 > 1$, so it is divergent.

$$9. \sum_{n=1}^{\infty} \frac{3^{n-1}}{6^{n-1}} \rightarrow 1$$

→ Solution,

$$\begin{aligned} & \sum_{n=1}^{\infty} \left(\frac{3^{n-1}}{6^{n-1}} - \frac{1}{6^{n-1}} \right) \\ &= \sum_{n=1}^{\infty} \left(\frac{1}{2^{n-1}} \right) - \sum_{n=1}^{\infty} \frac{1}{6^{n-1}} \\ &= \left(1 + \frac{1}{2} + \frac{1}{2^2} + \dots \right) - \left(1 + \frac{1}{6} + \frac{1}{6^2} + \dots \right) \end{aligned}$$

Common diff (r_1) = $\frac{1}{2}$ $\because |r| < 1$ so,

Common diff (r_2) = $\frac{1}{6}$ $|r| < 1$, they are convergent

$$\text{Sum}(S_{\infty}) = \frac{a_1}{1-r_1} = \frac{1}{1-\frac{1}{2}} = 2$$

$$\text{Sum}(S_{\infty}) = \frac{a_1}{1-r_2} = \frac{1}{1-\frac{1}{6}} = 2 \times \frac{1}{1} \times \frac{6}{5} = 1.2$$

$$\therefore \text{Sum}(S_{\infty}) = 2 - 1.2 = 0.8$$

* Ratio Test :

Let $\sum a_n$ be series with positive terms such that,

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = L$$

then,

a) If $L < 1$, then convergent series

b) If $L > 1$, then divergent series.

c) If $L = 1$, then test is inconclusive

Note: When the question is in factorial form, then Use ratio test. $n!$...

* Investigate the convergence of following series.

a) $\sum_{n=0}^{\infty} \frac{2^n + 5}{3^n}$

→ Solution,

Given that, $a_n = \frac{2^n + 5}{3^n}$

$$a_{n+1} = \frac{2^{n+1} + 5}{3^{n+1}}$$

Now,

$$L = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \lim_{n \rightarrow \infty} \frac{2^{n+1} + 5}{3^{n+1}} \Big/ \frac{2^n + 5}{3^n}$$

∴, $L = \lim_{n \rightarrow \infty} \frac{2^n \cdot 2 + 5}{3^n \cdot 3} \times \frac{3^n}{2^n + 5}$

∴, $L = \lim_{n \rightarrow \infty} \frac{2^n (2 + \frac{5}{2^n})}{3 \cdot 2^n (1 + \frac{5}{2^n})}$

take n common now
as you can take n common
always

∴ $L = \frac{(2 + \frac{5}{\infty})}{3(1 + \frac{5}{\infty})}$

$\left[\infty + 1 = \infty \right]$ $2^\infty = 2^\infty = \infty$
 $\left[\infty = 0 \right]$ $\frac{1}{\infty} = 0$

∴, $L = \frac{2}{3}$


∴ $L < 1$, so it is convergent.

b) $\sum_{n=1}^{\infty} \frac{2^n}{n!}$

→ Given, $a_n = \frac{2^n}{n!}$ $a_{n+1} = \frac{2^{n+1}}{(n+1)!}$

$$n! = n(n-1)(n-2)!$$

$$[\infty + 1 = \infty] \quad \frac{1}{\infty} = 0$$



Now,

$$L = \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \frac{2^{n+1}}{(n+1)!} \times \frac{n!}{2^n} = \frac{2^n \cdot 2}{(n+1) \cdot n!} \times \frac{n!}{2^n}$$

$$\therefore L = \lim_{n \rightarrow \infty} \frac{2}{n+1}$$

$$\therefore L = \frac{2}{\infty} = 0 < 1,$$

so it is convergent.

$$3. \sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}$$

$$\rightarrow a_n = \frac{4^n n! n!}{(2n)!}, a_{n+1} = \frac{4^{n+1} (n+1)! (n+1)!}{(2n+2)!}$$

Now,

$$L = \lim_{n \rightarrow \infty} \frac{4^n \cdot 4 (n+1)! (n+1)! \times (2n)!}{(2n+2)! 4^n n! n!}$$

$$= \lim_{n \rightarrow \infty} \frac{4^n \cdot 4 (n+1) n! (n+1) (n)! \times (2n)!}{(2n+2) (2n+1) (2n)! 4^n n! n!}$$

$$= \lim_{n \rightarrow \infty} \frac{4 \cdot \cancel{4} (1 + \frac{1}{n}) \cancel{n} (1 + \frac{1}{n})}{\cancel{4} (2 + \frac{1}{n}) \cancel{n} (2 + \frac{1}{n})}$$

$$= \frac{4 (1+0) (1+0)}{(2+0) (2+0)}$$

$$= \frac{4}{4}$$

$$= 1$$

$\therefore L = 1$, so it is inconclusive.

$$4. \sum_{n=1}^{\infty} \frac{n^2}{3^n}$$

$$\rightarrow \text{solution } a_n = \frac{n^2}{3^n}, a_{n+1} = \frac{(n+1)^2}{3^{n+1}}$$

$$L = \lim_{n \rightarrow \infty} \frac{(n+1)^2}{3(n+1)} \times \frac{3^n}{n^2}$$

$$= \lim_{n \rightarrow \infty} \frac{n^2 + 2n + 1}{3^n \cdot 3} \times \frac{3^n}{n^2}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{3} \left(\frac{n^2 + 2n + 1}{n^2} \right)$$

$$= \lim_{n \rightarrow \infty} \frac{1}{3} \left(1 + \frac{2}{n} + \frac{1}{n^2} \right)$$

$$= \frac{1}{3} (1 + 0 + 0) = L = \frac{1}{3} < 1, \text{ so it is convergent.}$$

$$5. \sum_{n=1}^{\infty} \frac{(2n)!}{n!n!}$$

$$\rightarrow a_n = \frac{(2n)!}{n!n!}, \quad a_{n+1} = \frac{(2n+1)!}{(n+1)!(n+1)!}$$

Now,

$$L = \lim_{n \rightarrow \infty} \frac{(2n+1)!}{(n+1)!(n+1)!} \times \frac{n!n!}{(2n)!}$$

$$= \lim_{n \rightarrow \infty} \frac{(2n+2)(2n+1)!}{(n+1)!!(n+1)!!} \times \frac{n!n!}{(2n)!}$$

$$= \lim_{n \rightarrow \infty} \frac{\pi(2 + \frac{2}{n})\pi(2 + \frac{1}{n})}{\pi(1 + \frac{1}{n})\pi(1 + \frac{1}{n})}$$

$$= (2+0)(2+0)$$

$$= 4$$

$$\therefore L = 4 > 1, \text{ so it is divergent.}$$

$$6. \sum_{n=1}^{\infty} \frac{n!}{n^n} \quad \left(\text{Formula: } \lim_{n \rightarrow \infty} \left(1 + \frac{u}{n}\right)^n = e^u \right)$$

$$\rightarrow a_n = \frac{(n)!}{n^n}, \quad a_{n+1} = \frac{(n+1)!}{(n+1)^{n+1}}$$

Now,

$$\begin{aligned} L &= \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \frac{(n+1)!}{(n+1)^{n+1}} \times \frac{n^n}{n!} \times \left(1 + \frac{1}{n}\right)^n, e^n \\ &= \lim_{n \rightarrow \infty} \frac{(n+1) \cdot n!}{(n+1)^n \cdot (n+1)^n} \times \frac{n^n}{n!} \times \left(1 + \frac{1}{n}\right)^n, e^n \\ &= \lim_{n \rightarrow \infty} \left(\frac{n}{n+1} \right)^n \times \left(1 + \frac{1}{n}\right)^n, e^n \\ &= \lim_{n \rightarrow \infty} \left[\frac{n}{\pi(1 + \frac{1}{n})} \right]^n \\ &= \lim_{n \rightarrow \infty} \left(\frac{1}{(1 + \frac{1}{n})^n} \right), \quad \left[\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e \right] \\ &= \frac{1}{e} \end{aligned}$$

$$\Rightarrow \sum_{n=1}^{\infty} \frac{n 2^n (n+1)!}{3^n n!}$$

$$\rightarrow a_n = \frac{n 2^n (n+1)!}{3^n n!}, \quad a_{n+1} = \frac{(n+1) 2^{n+1} (n+2)!}{3^{n+1} (n+1)!}$$

Now

$$\begin{aligned} L &= \lim_{n \rightarrow \infty} \frac{(n+1) 2^n \cdot 2 (n+2)!}{3^n \cdot 3 (n+1)!} \times \frac{3^n n!}{n 2^n (n+1)!} \\ &= \lim_{n \rightarrow \infty} \frac{(n+1) 2 (n+2) (n+1)!}{3 (n+1) (n)!} \times \frac{n!}{n (n+1)!} \\ &= \lim_{n \rightarrow \infty} \frac{2}{3} \frac{2(1 + \frac{1}{n}) \cdot \pi(1 + \frac{2}{n})}{2(1 + \frac{1}{n}) \cdot \pi} \\ &= \frac{2}{3} \frac{(1+0)(1+0)}{1+0} \end{aligned}$$

$L = \frac{2}{3} < 1$, so it is convergent.

$$3! = 3 \times 2 \times 1 = 6$$

$$8) \sum_{n=1}^{\infty} \frac{(n+3)!}{3! \cdot n! \cdot 3^n}$$

→ Solution

$$a_n = \frac{(n+3)!}{3! \cdot n! \cdot 3^n}$$

$$a_{n+1} = \frac{(n+1+3)!}{3! \cdot (n+1)! \cdot 3^{n+1}}$$

$$\begin{aligned} L &= \frac{a_{n+1}}{a_n} = \lim_{n \rightarrow \infty} \frac{(n+4)!}{3! \cdot (n+1)! \cdot 3^n \cdot 3} \times \frac{3! \cdot n! \cdot 3^n}{(n+3)!} \\ &= \lim_{n \rightarrow \infty} \frac{(n+4) \cdot (n+3)!}{(n+1) \cdot (n)! \cdot 3} \times \frac{n!}{(n+3)!} \\ &= \lim_{n \rightarrow \infty} \frac{n \cdot (1 + \frac{4}{n})}{n \cdot (1 + \frac{4}{n}) \cdot 3} \\ &= \frac{1+0}{(1+0) \cdot 3} \\ \therefore L &= \frac{1}{3} < 1, \text{ so it is convergent.} \end{aligned}$$

Root Test:-

Let $\sum a_n$ be a series, such that,

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = L$$

$$\sqrt[n]{a_n} = (a_n)^{\frac{1}{n}}$$

$$\begin{cases} L &= 1 \\ n &= 1 \end{cases}$$

then,

- a) if $L < 1$, the series is convergent.
- b) if $L > 1$, or infinite, then series is divergent.
- c) if $L = 1$, then inconclusive.

Note:-

$$\text{if } \lim_{n \rightarrow \infty} \sqrt[n]{n} = n^{\frac{1}{n}} = 1$$

$$\lim_{n \rightarrow \infty} \left(1 + \frac{u}{n}\right)^n = e^u$$

$$\lim_{n \rightarrow \infty} \frac{d(a^n)}{du} = a^n \ln a$$

Test the convergence of the following series.

1.
$$\sum_{n=1}^{\infty} \frac{n^2}{2^n}$$

→ Solution, $a_n = \frac{n^2}{2^n}$

$$L = \lim_{n \rightarrow \infty} (a_n)^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \left(\frac{n^2}{2^n}\right)^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \left(\frac{n^2}{2^n}\right)^{\frac{1}{n}}$$

$$= \lim_{n \rightarrow \infty} \frac{(n^2)^{\frac{1}{n}}}{2^n \times \frac{1}{n}}$$

∴ $L = \frac{1}{2} < 1$, so it is convergent.

2.
$$\sum_{n=1}^{\infty} \frac{2^n}{n^2}$$

→ Solution, $a_n = \frac{2^n}{n^2}$

$$\hookrightarrow L = \lim_{n \rightarrow \infty} (a_n)^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \left(\frac{2^n}{n^2}\right)^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{2^{\frac{1}{n}}}{(n^{\frac{1}{n}})^2} = \frac{2}{1}$$

∴ $L = 2 > 1$, so it is divergent.

3.
$$\sum_{n=1}^{\infty} \left(\frac{1}{1+n}\right)^n$$

→ Solution,

$$a_n = \left(\frac{1}{1+n}\right)^n$$

$$L = \lim_{n \rightarrow \infty} \left(\frac{1}{1+n}\right)^{n \times \frac{1}{n}} = \frac{1}{1+\infty} = \frac{1}{\infty} = 0$$

∴ $L = 0 < 1$, so it is convergent.

4.
$$\sum_{n=1}^{\infty} \frac{n^{10}}{10^n}$$

→ Solution;

$$\begin{aligned}
 L &= \lim_{n \rightarrow \infty} \left(\frac{n^{10}}{10^n} \right)^{\frac{1}{n}} \\
 &= \lim_{n \rightarrow \infty} \frac{(n^{\frac{1}{n}})^{10}}{10^{n \times \frac{1}{n}}} \\
 &= \frac{1^{10}}{10}
 \end{aligned}$$

$\therefore L = \frac{1}{10} = 0.1 < 1$, so it is convergent.

$$5. \sum_{n=1}^{\infty} \left(\frac{n}{3n+1} \right)^n$$

$$\rightarrow a_n = \left(\frac{n}{3n+1} \right)^n$$

$$L = \lim_{n \rightarrow \infty} \left(\frac{n}{3n+1} \right)^{n \times \frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n}{n(3+\frac{1}{n})} = \frac{1}{3+0} = \frac{1}{3}$$

$\therefore L = \frac{1}{3} < 1$, so it is convergent.

$$6. \sum_{n=1}^{\infty} \left(1 + \frac{1}{n} \right)^{-n^2}$$

$$\rightarrow \text{Solution, } a_n = \left(1 + \frac{1}{n} \right)^{-n^2}$$

$$L = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^{-n^2 \times \frac{1}{n}}$$

$$= \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^{-n}$$

$$\left[\because \left(1 + \frac{u}{n} \right)^n = e^u \right]$$

$\therefore L = e^{-1} = \frac{1}{e} = 0.36 < 1$, so it is convergent.

$$7. \sum_{n=1}^{\infty} \left(\frac{n}{n+1} \right)^{n^2}$$

$$\text{or, } a_n = \left(\frac{n}{n+1} \right)^{n^2}$$

$$\text{or, } L = \lim_{n \rightarrow \infty} \left(\frac{n}{n+1} \right)^{n^2 \times \frac{1}{n}}$$

$$\text{or, } L = \lim_{n \rightarrow \infty} \left(\frac{n}{n+1} + \frac{1}{n} \right)^n$$

$$= \lim_{n \rightarrow \infty} (1 +$$

$$\textcircled{a}, L = \lim_{n \rightarrow \infty} \left(\frac{\pi}{\pi(1 + \frac{1}{n})} \right)^n$$

$$\textcircled{a} L = \lim_{n \rightarrow \infty} \frac{1}{(1 + \frac{1}{n})^n}$$

$\textcircled{a} L = \frac{1}{e} = 0.36 < 1$, so it is convergent.

$$8. \sum_{n=1}^{\infty} \frac{n^n}{2^{n^2}} \quad \left[\frac{d(a^n)}{dn} = a^n \ln a \right]$$

$$\rightarrow \text{Solution, } a_n = \frac{n^n}{2^{n^2}}$$

$$\therefore L = \lim_{n \rightarrow \infty} \left(\frac{n}{2^n} \right)^{n \times \frac{1}{n}} = \lim_{n \rightarrow \infty} \left(\frac{n}{2^n} \right)$$

$$= \lim_{n \rightarrow \infty} \frac{1}{2^n \ln 2} \quad [\because \text{L'Hospital Rule}]$$

$$= \frac{1}{\infty}$$

$\therefore L = 0 < 1$, so, it is convergent.

* Integral Test (P-Series Test) :-

$$\int_{u=a}^{u=b} f(u) = du = \text{finite value} \Rightarrow \text{convergent series.}$$

$$u=a$$

$$x=b$$

$$\int_{x=a}^{x=b} f(u) = du = \text{infinite value} \Rightarrow \text{Divergent series.}$$

$$x=a$$

✓ P-Series Test (P-Test) :-

$$\left(\sum_{n=1}^{\infty} \frac{1}{n^p} = \frac{1}{n} + \frac{1}{n^2} + \frac{1}{n^3} + \dots \right)$$

$$\int_{u=1}^{u=\infty} \frac{1}{u^p} du = \left[\frac{u^{-p+1}}{-p+1} \right]_1^{\infty}$$

$$= \lim_{a \rightarrow \infty} \left[\frac{u^{-p+1}}{-p+1} \right]_1^a$$

$$= \frac{1}{1-p} \lim_{a \rightarrow \infty} \left[u^{-p+1} \right]_1^a$$

when,

$$P \cancel{>} 1 \text{ i.e. } P < 1 \quad u^{-2+\frac{1}{p}} = u^{-\frac{1}{p}} = \frac{1}{u^{\frac{1}{p}}}$$

$$\lim_{a \rightarrow \infty} = \frac{1}{1-p} \left[\frac{1}{u^{p-1}} \right]_1^a \quad P > 1$$

$$= \frac{1}{1-p} \lim_{a \rightarrow \infty} \left[\frac{1}{a^{p-1}} - \frac{1}{1^{p-1}} \right]$$

$$= \frac{1}{1-p} (0 - 1)$$

$$= \frac{1}{p-1} = \text{(finite)}$$

Hence the Series is convergent

Eg when $P > 1$.

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots \quad \text{converges.}$$

when $P < 1$.

$$= \frac{1}{1-p} \lim_{a \rightarrow \infty} \left[u^{1-p} \right]_1^a \quad \text{let } p = -1 \quad u^{1+1} = u^2$$

$$= \frac{1}{1-p} \lim_{a \rightarrow \infty} \left[a^{1-p} - 1^{1-p} \right]$$

$$= \frac{1}{1-p} \times \infty$$

$= \infty$ divergent.

$$\begin{array}{l} \infty + 1 = \infty \\ \infty - 1 = \infty \\ \infty \times 1 = \infty \\ \infty / 1 = \infty \end{array}$$

$$\frac{1}{\infty} = 0$$

~~$$\sum_{n=1}^{\infty} \frac{1}{n^{1/2}} = \frac{1}{1^{1/2}} + \frac{1}{2^{1/2}} + \frac{1}{3^{1/2}} + \dots$$~~

when $p = 1$ $p = 1$

$$= \frac{1}{1-p} \lim_{a \rightarrow \infty} \left[u^{1-p} \right]_1^a$$

$$= \frac{1}{1-p} \lim_{a \rightarrow \infty} \left[a^{1-p} - 1^{1-p} \right]$$

$$= \frac{1}{0} \left[a^{1-1} - 1^{1-1} \right]$$

$$= \infty \times 1$$

 $= \infty$ (divergent)

e.g.

$$\sum_{n=1}^{\infty} \frac{1}{n} = \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots$$

∴ Harmonic Series = divergent.

* P-Test by Integral Test

$$\sum_{n=1}^{\infty} \frac{1}{n^p} = \frac{1}{1^p} + \frac{1}{2^p} + \frac{1}{3^p} + \dots$$

when $p > 1$, then series is convergent.when $p \leq 1$, then series is divergentwhen $p = 1$, then series is inconclusive.

$$\sum_{n=1}^{\infty} \frac{1}{n^{p=1}} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots$$

Harmonic Series are always divergent.

$$\sum_{n=1}^{\infty} \frac{1}{n} du = [\log u]_1^{\infty} = \log \infty - \log 1 = \infty$$

Harmonic Series is Divergent.

$$\sum_{n=1}^{\infty} \frac{1}{n}$$

→ Ratio Test, $a_n = \frac{1}{n}$, $a_{n+1} = \frac{1}{n+1}$

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \lim_{n \rightarrow \infty} \left(\frac{1}{n+1} \right) \times \frac{n}{1} = \frac{n}{n+1} = 1$$

$\therefore L = 1$, so test is inconclusive.

example:

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \dots$$

$$\frac{1 + \frac{1}{2}}{2} = 1.5$$

$$\frac{2}{4} = \frac{1}{2} = \frac{1}{3} + \frac{1}{4} = 0.58$$

$$\frac{4}{8} = \frac{1}{2}$$

$$0.58 > \frac{2}{4} = \frac{1}{2}$$

$$= 0.634 > \frac{4}{8} > \frac{1}{2}$$

* Limit Comparison Test:

let $\sum a_n$ and $\sum b_n$ be series then

a) when $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = L > 0$, then series both converge or diverges.

b) when $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \infty$, and $\sum b_n$ diverges then $\sum a_n$ also diverges.

to find b_n compare power and put highest power in.

Questions:

a) $\sum_{n=1}^{\infty} \frac{2n+1}{n^2+2n+1}$

→ Solution,

$$a_n = \frac{2n+1}{n^2+2n+1}, \quad b_n = \frac{2n}{n^2} = \frac{2}{n}$$

$b_n = \frac{2}{n}$, $P = 1$, so by P-test b_n is divergent.

$$\text{Now, } \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{2n+1}{n^2+2n+1} \times \frac{n}{2} = \lim_{n \rightarrow \infty} \frac{n^2(2 + \frac{1}{n})}{n^2(1 + \frac{2}{n} + \frac{1}{n^2})} = 2$$

$$= \frac{2 + \frac{1}{2^2}}{(1 + \frac{2}{2} + \frac{1}{2^2}) \times 2}$$

$$= \frac{2}{2} = 1$$

$\therefore 2n$ is also divergent series.

$$2) \sum_{n=1}^{\infty} \frac{\sqrt{n}}{n^2 + 1}$$

$$\rightarrow \text{Solution, } a_n = \frac{\sqrt{n}}{n^2 + 1}, b_n = \frac{\sqrt{n}}{n^2}$$

Now,

$$b_n = \frac{n^{1/2}}{n^2} = \frac{1}{n^{3/2}}$$

Since $p > 1$, so it is convergent by p-test.

We have,

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{n^2 + 1} \times \frac{n^2}{\sqrt{n}} = \frac{n^2}{n^2(1 + \frac{1}{n^2})} = \frac{1}{1+0} = 1$$

$\therefore L = 1 > 0$, so a_n is also a convergent series.

$$4) \sum_{n=1}^{\infty} \frac{n^4 + 5}{n^5}$$

$$\rightarrow \text{Solution, } a_n = \frac{n^4 + 5}{n^5}, b_n = \frac{n^4}{n^5} = \frac{1}{n}$$

$\therefore b_n = \frac{1}{n}$, $p=1$, so it is divergent by p-test

Now,

$$\frac{a_n}{b_n} = \frac{n^4 + 5}{n^5} \times \frac{n^5}{n^4} = \lim_{n \rightarrow \infty} n^4 \left(1 + \frac{5}{n^4}\right) = 1 + 0 = 1$$

$\therefore L = 1 > 0$, so it is also a divergent series.

$$3) \sum_{n=1}^{\infty} (\sqrt{n^2+1} - n)$$

$$\begin{aligned} \rightarrow a_n &= \sqrt{n^2+1} - n \\ &= \frac{\sqrt{n^2+1} - n}{1} \cdot \frac{\sqrt{n^2+1} + n}{\sqrt{n^2+1} + n} \\ &= \frac{(\sqrt{n^2+1})^2 - (n)^2}{\sqrt{n^2+1} + n} \\ &= \frac{n^2 + 1 - n^2}{\sqrt{n^2+1} + n} \end{aligned}$$

$$\therefore a_n = \frac{1}{\sqrt{n^2+1} + n}, \quad b_n = \frac{1}{n}$$

$\because P=1$, so it is divergent from P-Test.

we know,

$$\frac{a_n}{b_n} = \frac{1}{\sqrt{n^2+1} + n} \times \frac{n}{1}$$

$$\lim_{n \rightarrow \infty} = \frac{n}{n(\sqrt{1+\frac{1}{n^2}} + 1)}$$

$$\lim_{n \rightarrow \infty} = \frac{1}{\sqrt{1+0+1}}$$

$$\therefore L = \frac{1}{2} < 0.5 \neq 0$$

so it is also divergent series.

$$5) \sum_{n=1}^{\infty} \frac{2n-1}{n(n+1)(n+2)}$$

$$\rightarrow \text{solution, } a_n = \frac{2n-1}{n(n+1)(n+2)}$$

$$\begin{aligned} &= \frac{2n-1}{n^2(n+2) + n(n+2)} = \frac{2n-1}{n^3 + 2n^2 + n^2 + 2n} \\ &= \frac{2n-1}{n^3 + 3n^2 + 2n} \end{aligned}$$

$$\therefore b_n = \frac{2n}{n^3} = \frac{2}{n^2}$$

$\because P > 1$, so it is convergent series by P-test.

Now,

$$\frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{(2n-1)}{n^3 + 3n^2 + 2n} \times \frac{n^2}{2}$$

$$= \lim_{n \rightarrow \infty} \frac{n^3(2 - \frac{1}{n^3})}{n^3(1 + \frac{3}{n} + \frac{2}{n^2})} = \lim_{n \rightarrow \infty} \frac{2 - \frac{1}{n^3}}{1 + \frac{3}{n} + \frac{2}{n^2}} = \frac{2}{2} = 1$$

$\therefore L = 1$, so a_n is also convergent series

$$6. \sum_{n=1}^{\infty} \frac{1}{2\sqrt{n+1}}$$

$$\rightarrow a_n = \frac{1}{2\sqrt{n+1}}, \quad b_n = \frac{1}{2\sqrt{n}} = \frac{1}{2n^{1/2}}$$

$\because P < 1$, so it is divergent by P-test.

Now,

$$\frac{a_n}{b_n} = \frac{1}{2\sqrt{n+1}} \times \frac{2\sqrt{n}}{1} = \frac{\sqrt{n}(2)}{\sqrt{n}(2 + \frac{1}{\sqrt{n}})} = \frac{2}{2+0} = 1$$

$\therefore L > 0$, so a_n is also a divergent series.

$$7. \sum_{n=1}^{\infty} \tan\left(\frac{1}{n}\right)$$

$$\rightarrow \text{solution, } a_n = \tan\left(\frac{1}{n}\right), \quad b_n = \frac{1}{n}$$

$b_n = \frac{1}{n} = P = 1$, so it is divergent series.

Now,

$$\frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\tan\left(\frac{1}{n}\right)}{\left(\frac{1}{n}\right)} = \lim_{n \rightarrow \infty} \frac{\sec^2\left(\frac{1}{n}\right)}{\left(-\frac{1}{n^2}\right)}$$

[L-Hospital Rule]

$$= \sec^2(0) = \frac{1}{\cos^2 0} = 1 > 0$$

so it is also divergent series.

8) $\sum_{n=1}^{\infty} \frac{1}{2^{n-1}}$

→ Solution, $a_n = \frac{1}{2^{n-1}}$, $b_n = \frac{1}{2^n}$

For b_n , using root test

$$b_n = \frac{1}{2^n} = \left(\frac{1}{2^n}\right)^{1/n} = \left(\frac{1}{2}\right)^{n \times \frac{1}{n}} = \frac{1}{2} = 0.5$$

$\therefore L = 0.5 < 1$, so it is convergent series.

Now,

$$\frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{2^{n-1}} \times \frac{2^n}{1} = \frac{2^n}{2^{n-1}} = 1$$

so a_n is also convergent series.

n-term test for convergence:

The necessary condition for the convergence of infinite series $\sum a_n$ is,


$$\lim_{n \rightarrow \infty} a_n = 0$$

but this is not sufficient.

Exception:

$$\sum_{n=1}^{\infty} \frac{1}{n} = \text{divergent Series}$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0 \text{ (convergent)}$$



* Questions:-

$$\therefore \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e^1$$

a) $\sum_{n=1}^{\infty} n^2 = \lim_{n \rightarrow \infty} n^2 = \infty$, so it is divergent.

b) $\sum_{n=1}^{\infty} \frac{n+1}{n} = \lim_{n \rightarrow \infty} \frac{\pi(1) + 1/n}{\pi(1)} = \lim_{n \rightarrow \infty} 1 + \frac{1}{n} = 1$, Divergent.

c) $\sum_{n=1}^{\infty} \frac{-n}{2n+5} = \lim_{n \rightarrow \infty} \frac{-n}{\pi(2+5/n)} = -\frac{1}{2}$, Divergent.

d) $\sum_{n=1}^{\infty} \left(1 + \frac{1}{2n}\right)^n = \lim_{n \rightarrow \infty} \left[\left(1 + \frac{1}{2n}\right)^{2n}\right]^{\frac{1}{2}} = e^{1 \cdot \frac{1}{2}} = e^{0.5} \neq 0$
Divergent.

Alternating Series:-

1. $\sum_{n=1}^{\infty} (-1)^{n+1} a_n = a_1 - a_2 + a_3 - a_4 + \dots$ when $n = 1, 2, 3, \dots$

$$\sum_{n=1}^{\infty} (-1)^n a_n = -a_1 + a_2 - a_3 + a_4 - \dots$$

2. $a_1 > a_2 > a_3 > \dots$ decreasing orders.

3. Alternating series test: [Leibniz Test]

* Questions:-

$$\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

i.e. $a > b_2 > b_3 > \dots$

and also 'in (+, -, +, -) alternative sign'

* Leibniz Test:-

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0, \text{ Convergent Series.}$$

* Absolute Convergent:-

A series $\sum_{n=1}^{\infty} a_n$ converges absolutely if the corresponding series $\sum_{n=1}^{\infty} |a_n|$ converges.

$$* 1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^2}$$

Leibniz Test:-

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = 0, \text{ Convergent.}$$

Absolute Convergent:-

$$\sum_{n=1}^{\infty} \left| (-1)^{n+1} \frac{1}{n^2} \right| = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

$$\sum_{n=1}^{\infty} |a_n| = \frac{1}{n^2} \quad (\text{if Leibniz & abel's convergence test/alternating series test is satisfied})$$

\therefore by P-test, $P > 1$, so convergent

So the series is absolutely Convergent.

* Conditional Convergent:-

$$a) \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

absolute value :

$$\sum_{n=1}^{\infty} \left| (-1)^{n+1} \frac{1}{n} \right| = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} + \dots$$

$\therefore \sum |a_n| = \sum \left| \frac{1}{n} \right|$, by P-test, $P = 1$, so series is divergent

Leibniz Test :-

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0, \text{ so convergent}$$

$\textcircled{P=1}$

\therefore Series is conditional convergent.

$$* \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^3}$$

$$\Rightarrow 1 - \frac{1}{2^3} + \frac{1}{3^3} - \frac{1}{4^3} + \dots + \frac{1}{n^3} + \dots$$

Leibniz Test :-

$\lim_{n \rightarrow \infty} \frac{1}{n^3} = 0$, so Series is convergent by This test.

Absolute Value :-

$$\sum_{n=1}^{\infty} \left| (-1)^{n+1} \frac{1}{n^3} \right| = 1 + \frac{1}{2^3} + \frac{1}{3^3} + \frac{1}{4^3} + \dots$$

$$\sum |a_n| = \sum \left| \frac{1}{n^3} \right|,$$

Since $P > 1$, so convergent.

So Series is convergent

$$* \sum_{n=1}^{\infty} (-1)^{n+1} \frac{n}{2^n}$$

$$2^n du = 2^n \ln 2$$


$$\Rightarrow \frac{n}{2^n} = \frac{1}{2} - \frac{1}{2^2} + \frac{3}{2^3} - \frac{4}{2^4} + \dots + \frac{n}{2^n} + \dots$$

Leibniz Test :-

$$\lim_{n \rightarrow \infty} \frac{n}{2^n} = \lim_{n \rightarrow \infty} \frac{1}{2^n \ln 2} \quad \{ \text{L-Hospital} \}$$

$$= 0$$

so convergent



Absolute Value

$$\sum_{n=1}^{\infty} \left| (-1)^{n+1} \frac{n}{2^n} \right| = \frac{1}{2} + \frac{2}{2^2} + \frac{3}{2^3} + \frac{4}{2^4} + \dots$$

$$\therefore \left(\sum_{n=1}^{\infty} \left| \frac{n}{2^n} \right| \right) =$$

* $\sum_{n=1}^{\infty} (-1)^n \frac{1}{2^n}$

$$\rightarrow \text{series} = -\frac{1}{2} + \frac{1}{2^2} - \frac{1}{2^3} + \frac{1}{2^4} - \dots + \frac{1}{2^n} - \dots$$

Leibniz test,

$$\lim_{n \rightarrow \infty} \frac{1}{2^n} = \frac{0}{2n \ln 2} = 0, \text{ Converges.}$$

Absolute Value

$$\sum_{n=1}^{\infty} \left| (-1)^n \frac{1}{2^n} \right| = \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \dots$$

$$\sum_{n=1}^{\infty} \left| \frac{1}{2^n} \right| =$$

Summary

i) Geometric series:-

If $|r| \geq 1$, then series diverges, otherwise it converges.

ii) For non-negative terms:-

Use integral test, ratio test, root test, and limit comparison test.

iii) Alternating series:-

Leibniz's test.

iv) n term test for divergence:-

Unless $a_n \rightarrow 0$

limit $a_n = 0$, convergent, otherwise divergent.
 $n \rightarrow \infty$

v) Series with absolute convergent:-

$\sum |a_n|$ = convergent (absolute convergent)

~~$\sum |a_n| \neq \sum a_n$~~

Power Series:-

A series about $u=0$ in the form of,

$\sum_{n=0}^{\infty} a_n u^n = a_0 + a_1 u + a_2 u^2 + a_3 u^3 + \dots + a_n u^n + \dots$
 is called power series.

at $u=a$

$\sum_{n=0}^{\infty} a_n (u-a)^n = a_0 + a_1 (u-a) + a_2 (u-a)^2 + \dots + a_n (u-a)^n$

where, a = center and $a_0, a_1, a_2, a_3, \dots, a_n$ = coefficient / constant.

* Test the convergence for the given power series.

a) $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{u^n}{n} = u - \frac{u^2}{2} + \frac{u^3}{3} - \dots$

$\rightarrow \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = (-1)^{n-1+1} \frac{u^{n+1}}{n+1} \times \frac{n}{(-1)^{n-1} u^n}$

$$= (-1)^n \frac{u^{n+1}}{n+1} \times \frac{n}{(-1)^{n-1} u^n}$$

$$= (-1)^n \frac{u^n \cdot u}{(n+1)} \times \frac{n}{(-1)^n (-1)^{-1} u^n}$$

$$= \left| \frac{n}{n+1} \cdot u \right| = \frac{n}{n(1+\frac{1}{n})} = \frac{1}{1+\frac{1}{n}} = 1$$

$$= |u|$$

$|u| < 1$, then converges absolutely.

$|u| > 1$, then divergent.

$$\text{b) } \sum_{n=1}^{\infty} (-1)^{n-1} \frac{u^{2n-1}}{2n-1} = u - \frac{u^3}{3} + \frac{u^5}{5} - \dots$$

→ Solution:

$$a_n = (-1)^{n-1} \frac{u^{2n-1}}{2n-1}, a_{n+1} = (-1)^{n-1+1} \frac{u^{2(n+1)-1}}{(2(n+1))-1}$$

Now,


$$\begin{aligned} \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| &= \frac{(-1)^n u^{2n+1}}{(2n+1)} \times \frac{(2n-1)}{(-1)^{n-1} \cdot (-1)^n \cdot u^{2n-1}} \\ &= \frac{u^{2n} \cdot u (2n-1)}{(2n+1) (-1)^{-1} u^{2n} \cdot u^{-1}} \\ &= \frac{u^2 (2n-1)}{(2n+1)} \\ &= |u^2| \end{aligned}$$

$|u^2| < 1$, converges absolutely

$|u^2| > 1$, then divergent.

$$\text{c) } \sum_{n=0}^{\infty} \frac{u^n}{n!} = 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots$$

$$\rightarrow a_{n+1} = 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots = \frac{u^{n+1}}{(n+1)!}$$



Now,

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{x+x^2+x^3}{2x \cdot 3x} + \dots$$

$$= \lim_{n \rightarrow \infty} \frac{u^{n+1}}{(n+1)!} \times \frac{n!}{u^n} = \frac{u^n \cdot u}{(n+1) \cdot n!} \times \frac{n!}{u^n} = \frac{u}{n+1}$$

for all value of u , it is zero. i.e. < 1 , so
converges absolutely.

$$\text{d.r.} \sum_{n=0}^{\infty} n! u^n = 1 + u + 2! u^2 + 3! u^3 + \dots$$

$$\rightarrow a_{n+1} = (n+1)! u^{n+1}, a_n = n! u^n$$

Now,

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1) \cdot n! \cdot u^{n+1}}{n! \cdot u^n} = (n+1) |u|$$

Divergent for all values of u . exception case.

Taylor Series And Maclaurin Series

Let the function 'f' with all derivatives of all orders throughout some interval containing 'a' as an interior point then the ^{Taylor Polynomial as of order n} Taylor Series generated by f at $u=a$ is $\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (u-a)^k$

Taylor series at $u=a$.

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (u-a)^k$$

$$= f(a) + f'(a)(u-a) + f''(a) \frac{(u-a)^2}{2!} + f'''(a) \frac{(u-a)^3}{3!} + \dots + \frac{f^{(n)}(a)}{n!} (u-a)^n + \dots$$

Maclaurin series: or

(Taylor Series at $x=0$): with no center (a).

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(0) \cdot u^k}{k!} = f(0) + f'(0)u + f''(0) \frac{u^2}{2!} + f'''(0) \frac{u^3}{3!} + \dots + \frac{f^{(n)}(0)}{n!} u^n + \dots$$

Q. Find the Taylor series and polynomial for the following at $u=0$.

g) $f(u) = e^u$ at $u=0$

→ Solution.

Given function $f(u) = e^u$

$$f'(u) = e^u$$

$$f''(u) = e^u$$

$$f'''(u) = e^u$$

$$\vdots$$

$$f^n(u) = e^u$$

$$f(0) = e^0 = 1$$


$$f'(0) = e^0 = 1$$

$$f''(0) = e^0 = 1$$

$$f'''(0) = e^0 = 1$$

$$\vdots$$

$$f^n(0) = e^0 = 1$$



Taylor series at $u=0$ (Maclaurin series) is,

$$\sum_{k=0}^{\infty} \frac{f^k(0)}{k!} u^k \quad (\text{upto } \infty)$$

$$= f(0) + f'(0) u + \frac{f''(0)}{2!} u^2 + \frac{f'''(0)}{3!} u^3 + \dots + f^n(0) \frac{u^n}{n!} + \dots$$

$$= 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots + \frac{u^n}{n!} + \dots$$

$$= \sum_{k=0}^{\infty} \frac{u^k}{k!}$$

Taylor Polynomial (only upto n)

$$\sum_{k=0}^n \frac{u^k}{k!} = 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots + \frac{u^n}{n!}$$

~~$$\rightarrow f(u) = e^{-u} \text{ at } u=0$$~~

Given function,

$$f(u) = e^{-u}$$

$$f(0) = 1$$

$$f'(u) = -e^{-u}$$

$$f'(0) = -1$$

$$f''(u) = e^{-u}$$

$$f''(0) = 1$$

$$f'''(u) = -e^{-u}$$

$$f'''(0) = -1$$

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1</p

$$= 1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \dots + \frac{(-u)^n}{n!} + \dots$$

$$= \sum_{k=0}^{\infty} \frac{(-u)^k}{k!}$$

Taylor polynomial :-

$$\sum_{k=0}^n \frac{(-u)^k}{k!} = 1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \dots + \frac{(-u)^n}{n!}$$

$$\Rightarrow \cosh u = \frac{e^u + e^{-u}}{2}$$

→ Solution,

As we know,

$$e^u = 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \frac{u^4}{4!} + \dots$$

$$e^{-u} = 1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \frac{u^4}{4!} - \dots$$

Now,

$$e^u + e^{-u} = \left(1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \frac{u^4}{4!} + \dots \right) +$$


$$\left(1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \frac{u^4}{4!} - \dots \right)$$

$$\therefore e^u + e^{-u} = 2 + 2\frac{u^2}{2!} + 2\frac{u^4}{4!} + \dots$$

$$\frac{e^u + e^{-u}}{2} = 1 + \frac{u^2}{2!} + \frac{u^4}{4!} + \dots + \frac{u^{2n}}{(2n)!} + \dots$$

$$\therefore \text{Taylor Series} = \sum_{k=0}^{\infty} \frac{u^{2k}}{(2k)!}$$

$$\therefore \text{Taylor polynomial} = \sum_{k=0}^n \frac{u^{2k}}{(2k)!}$$



$$\text{d. } \sinh u = \frac{e^u - e^{-u}}{2}$$

→ Solution,

As we know,

$$e^u = 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots$$

$$e^{-u} = 1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \dots$$

$$e^u - e^{-u} = \left(1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots\right) - \left(1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \dots\right)$$

$$\textcircled{a. } e^u - e^{-u} = \left(1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots\right) - \left(1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \dots\right)$$

$$\textcircled{b. } e^u - e^{-u} = 2u + \frac{2u^3}{3!} + \frac{2u^5}{5!} + \dots$$

$$\textcircled{c. } \frac{e^u - e^{-u}}{2} = u + \frac{u^3}{3!} + \frac{u^5}{5!} + \dots + \frac{u^{2n+1}}{(2n+1)!} + \dots$$

$$\therefore \text{Taylor series} = \sum_{k=0}^{\infty} \frac{u^{2k+1}}{(2k+1)!}$$

$$\therefore \text{Taylor polynomial} = \sum_{k=0}^n \frac{u^{2k+1}}{(2k+1)!}$$

$$\text{e. } f(u) = e^{4u} \quad \text{from derivative also you can but alt. method:}$$

→ Solution,

$$e^u = 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots$$

$$f(u) = e^u$$

$$f(0) = e^0 = 1$$

$$f'(u) = e^u$$

$$f'(0) = e^0 = 1$$

$$\vdots$$

$$f^n(u) = e^u$$

$$\vdots$$

$$f^n(0) = e^0 = 1$$

Similarly

for e^{4u} .

$$\text{Taylor Series} = 1 + 4u + \frac{(4u)^2}{2!} + \frac{(4u)^3}{3!} + \dots$$

$$= \sum_{k=0}^{\infty} \frac{(4u)^k}{k!}$$

$$\text{Taylor Polynomial} = \sum_{k=0}^n \frac{(4u)^k}{k!}$$

$$\Rightarrow e^{-u} = f(u)$$

→ Solution

As we know,

$$e^{-u} = 1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \dots$$

$$f(u) = e^{-u}$$

$$f(0) = e^0 = 1$$

$$f'(u) = -e^{-u}$$

$$f'(0) = -e^0 = -1$$

$$f''(u) = e^{-u}$$

$$f''(0) = e^0 = 1$$

⋮

⋮

$$f^n(u) = (-1)^n e^{-u}$$

$$f^n(0) = (-1)^n$$

$$\therefore \text{Taylor Series} = \sum_{k=0}^{\infty} \frac{f^k(0)}{k!} u^k$$

$$= f(0) + f'(0) u + f''(0) \frac{u^2}{2!} + \dots + \frac{u^n}{n!} + \dots$$

$$= 1 - u + \frac{u^2}{2!} - \frac{u^3}{3!} + \dots + \frac{(-u)^n}{n!} + \dots$$

$$= \sum_{k=0}^{\infty} \frac{(-u)^k}{k!}$$

Similarly

for e^{-7u}

$$\text{Taylor series} = \sum_{k=0}^{\infty} \frac{(-7u)^k}{k!} \#$$

$$= 1 - 7u + \frac{(-7u)^2}{2!} - \frac{(-7u)^3}{3!} + \dots + \frac{(-7u)^n}{n!} + \dots$$

$$\text{Taylor polynomial} = \sum_{k=0}^n \frac{(-7u)^k}{k!} \#$$

* Find the Taylor series and polynomial for the following function.

~~Ex~~ $f(u) = \cos u$. even derivative one side } odd one side.

→ Solution,

$$f(u) = \cos u \quad f(0) = 1$$

($\sin u$ doesn't contribute in series)

$$f'(u) = -\sin u \quad f'(0) = 0$$

$$f''(u) = -\cos u \quad f''(0) = -1$$

$$f'''(u) = +\sin u \quad f'''(0) = 0$$

$$f^{2n}(u) = (-1)^n \cos u \quad f^{2n}(0) = (-1)^n$$

$$f^{2n+1}(u) = (-1)^{n+1} \sin u$$

Now,

$$f^{2n+1}(0) = 0$$

Taylor series at $u=0$ (Maclaurin series)

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(0) u^k}{k!} = f(0) + f'(0)u + f''(0) \frac{u^2}{2!} + f'''(0) \frac{u^3}{3!} + \dots + f^{(n)}(0) \frac{u^n}{n!} + \dots$$

$$= 1 + 0 \times u + (-1) \times \frac{u^2}{2!} + 0 \times \frac{u^3}{3!} + \dots + \cancel{f^{2n}(0)} \frac{u^{2n}}{2n!} + \dots$$

+ ...

$$= 1 - \frac{u^2}{2!} + \dots + (-1)^n \frac{u^{2n}}{(2n)!} + \dots$$

$$= \sum_{k=0}^{\infty} (-1)^k \frac{u^{2k}}{(2k)!}$$

$$\therefore \text{Taylor polynomial} = \sum_{k=0}^n (-1)^k \frac{u^{2k}}{(2k)!} \#$$

$$= 1 - \frac{u^2}{2!} + \dots + (-1)^n \frac{u^{2n}}{(2n)!} \#$$

$$b) f(u) = \sin u \text{ at } u=0$$

→ Solution,

$$f(u) = \sin u \quad f(0) = 0 \quad f'(u) = \cos u \quad f'(0) = 1$$

$$f''(u) = -\sin u \quad f''(0) = 0 \quad f'''(u) = -\cos u \quad f'''(0) = -1$$

$$f^{2n}(u) = (-1)^n \sin u \quad f^{2n+1}(u) = (-1)^n \cos u$$

$$f^{2n}(0) = 0 \quad f^{2n+1}(0) = (-1)^n$$

Now,

$$\text{Taylor Series} = \sum_{k=0}^{\infty} \frac{f^{(k)}(0) u^k}{k!} = f(0) + f'(0) u + f''(0) \frac{u^2}{2!} + \dots + f^n(0) \frac{u^n}{n!} + \dots$$

$$= 0 + u - 0 - \frac{1}{3!} u^3 + 0 - \dots + f^{2n+1}(0) \frac{u^{2n+1}}{(2n+1)!} + \dots$$

$$= u - \frac{u^3}{3!} + \dots + (-1)^n \frac{u^{2n+1}}{(2n+1)!} + \dots$$

$$= \sum_{k=0}^{\infty} (-1)^k \frac{u^{2k+1}}{(2k+1)!} \#$$

$$\text{Taylor polynomial} = u - \frac{u^3}{3!} + \dots + (-1)^n \frac{u^{2n+1}}{(2n+1)!}$$

$$= \sum_{k=0}^n \frac{u^{2k+1}}{(2k+1)!} \#$$

$$\Rightarrow f(u) = \sin 4u \text{ at } u=0$$

→ Solution,

for $\sin u$

$$f(u) = \sin u$$

$$f(0) = 0$$

$$f'(u) = \cos u$$

$$f'(0) = 1$$

$$f''(u) = -\cos u \sin u$$

$$f''(0) = 0$$

$$f'''(u) = -\cos u$$

$$f'''(0) = -1$$

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

$$= u - \frac{u^3}{3!} + \dots + (-1)^n \frac{u^{2n+1}}{(2n+1)!}$$

$$= \sum_{k=0}^n (-1)^k \frac{u^{2k+1}}{(2k+1)!} \#.$$

Similarly for $\sin 4u$.

$$\text{Taylor polynomial} = \sum_{k=0}^n (-1)^k \frac{(4u)^{2k+1}}{(2k+1)!}$$

$$= 4u - \frac{(4u)^3}{3!} + \dots + (-1)^n \frac{(4u)^{2n+1}}{(2n+1)!} \#.$$

d) $f(u) = u \sin u$

→ Solution,

$$\begin{aligned} \text{for } f(u) = \sin u & \quad f(0) = 0 & f'(u) = \cos u & \quad f'(0) = 1 \\ f''(u) = -\sin u & \quad f''(0) = 0 & f'''(u) = -\cos u & \quad f'''(0) = -1 \\ & \vdots & \vdots & \vdots \\ f^{2n}(u) = (-1)^n \sin u & \quad f^{2n+1}(u) = (-1)^n \cos u & & \end{aligned}$$

$$f^{2n}(0) = 0$$

$$f^{2n+1}(0) = (-1)^n$$

Now,

$$\text{Taylor series for } \sin u = \sum_{k=1}^{\infty} \frac{f^k(0)}{k!} u^k$$

$$= f(0) + f'(0) u + f''(0) \frac{u^2}{2!} + f'''(0) \frac{u^3}{3!} + \dots + f^n(0) \frac{u^n}{n!}$$

$$= 0 + 1 \times u + 0 \times \frac{u^2}{2!} + (-1) \times \frac{u^3}{3!} + \dots + f^{2n+1}(0) \frac{u^{2n+1}}{(2n+1)!} + \dots$$

$$= u - \frac{u^3}{3!} + \dots + (-1)^n \frac{u^{2n+1}}{(2n+1)!} + \dots$$

$$= \sum_{k=0}^{\infty} (-1)^k \frac{u^{2k+1}}{(2k+1)!} \#.$$

Similarly

$$\begin{aligned}
 \text{Taylor series for } u \sin u &= u \left(u - \frac{u^3}{3!} + \dots + (-1)^n \frac{u^{2n+1}}{(2n+1)!} \right. \\
 &= \sum_{k=0}^{\infty} (-1)^k \frac{u^{2k+1} \cdot u}{(2k+1)!} \left. + \dots \right) \\
 &= \sum_{k=0}^{\infty} (-1)^k \frac{u^{2k+2}}{(2k+1)!} \# .
 \end{aligned}$$

Taylor series polynomial of $\sin u$

$$\begin{aligned}
 &= \left(u - \frac{u^3}{3!} + \dots + (-1)^n \frac{u^{2n+1}}{(2n+1)!} \right) \\
 &= \sum_{k=0}^n (-1)^k \frac{u^{2k+1}}{(2k+1)!}
 \end{aligned}$$

Similarly

Taylor polynomial of $u \sin u$

$$\begin{aligned}
 &= u \left(u - \frac{u^3}{3!} + \dots + (-1)^n \frac{u^{2n+1}}{(2n+1)!} \right) \\
 &= \sum_{k=0}^n (-1)^k \frac{u^{2k+2}}{(2k+1)!} \#
 \end{aligned}$$

$$\Rightarrow f(u) = \frac{1}{u} \text{ at } u=1 \text{ (Taylor series)}$$

→ Solution,

$$f(u) = \frac{1}{u} = u^{-1}$$

$$f(1) = 0$$

$$f'(u) = -u^{-2}$$

$$f'(1) = -1$$

$$f''(u) = 2u^{-3}$$

$$f''(1) = 2$$

$$f'''(u) = -6u^{-4}$$

$$f'''(1) = -6$$

$$f'$$

$$f'$$

$$f'$$

Taylor series at $u=1$ is $a=1$

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (u-a)^k = f(a) + f'(a)(u-a) + \frac{f''(a)}{2!} (u-a)^2 + \dots$$

$$= f(1) + f'(1)(u-1) + \frac{f''(1)}{2!} (u-1)^2 + \dots$$

$$= 1 - (u-1) + 2 \frac{(u-1)^2}{2!} - 6 \frac{(u-1)^3}{3!} + \dots$$

Taylor series $= \sum_{k=0}^{\infty} \frac{f^k(1)}{k!} (u-1)^k$ #

Taylor polynomial $= \sum_{k=0}^n \frac{f^k(1)}{k!} (u-1)^k$ #

* Taylor Polynomial of order n

Let f be a function with derivatives of order k for $k=1, 2, \dots, N$ in some interval containing a as an interior point. Then for any integer n from 0 through N , the Taylor polynomial of order n generated by f at $x=a$ is the polynomial

$$P_n(u) = f(a) + f'(a)(u-a) + \frac{f''(a)}{2!} (u-a)^2 + \dots$$

$$+ \frac{f^{(k)}(a)}{k!} (u-a)^k + \dots + \frac{f^{(n)}(a)}{n!} (u-a)^n$$

Multiple Integral :-

$\int du$ = Single integral

$\iint du dy$ = Double integral

$\iiint du dy dz$ = Triple integral

$$a) \int_0^3 \int_0^2 (4 - y^2) dy dz$$

$$= \int_0^3 \left[4y - \frac{y^3}{3} \right]_0^2 dz$$

$$= \int_0^3 \left[8 - \frac{8}{3} \right] dz$$

$$= \int_0^3 \left[\frac{24}{3} - \frac{8}{3} \right] dz$$

$$= \frac{16}{3} \left[z \right]_0^3 = \frac{16}{3} \times 3 = 16 \#.$$

$$b) \int_0^3 \int_0^2 (u^2 y - 2uy) dy du$$

$$= \int_0^3 \left[\frac{u^2 y^2}{2} - \frac{2uy^2}{2} \right]_0^2 du$$

$$= \int_0^3 \left[\frac{4u^2}{2} - \cancel{\frac{4u^2}{2}} + \frac{2u}{2} \right] du$$

$$= \frac{4}{2} \int_0^3 (u^2 - 2u) du$$

$$= \frac{4}{2} \left[\frac{u^3}{3} - \frac{2u^2}{2} \right]_0^3$$

$$= \frac{4}{2} \left[\frac{27}{3} - 9 \right]$$

$$= \frac{4}{2} \times 0 = 0 \#$$

$$\text{Q: } \int_{-1}^0 \int_{-1}^1 (u+y+1) du dy$$

$$= \int_{-1}^0 \left[\frac{u^2}{2} + uy + u \right]_{-1}^1 dy$$

$$= \int_{-1}^0 \left[\frac{1}{2} - \frac{(-1)^2}{2} + 1 \times y - (-1) \times y + 1 - (-1) \right] dy$$

$$= \int_{-1}^0 \left(\frac{1}{2} - \frac{1}{2} + 2y + 2 \right) dy$$

$$= 2 \times \frac{0}{2} - 2 \times \frac{(-1)^2}{2} + 2 \times 0 - 2 \times (-1) = -1 + 2 = 1 \#.$$

$$\text{d: } \int_{-\pi}^{2\pi} \int_0^{\pi} (\sin u + \cos y) du dy$$

$$= \int_{-\pi}^{2\pi} [-\cos u + u \cos y]_0^{\pi} dy$$

$$= \int_{-\pi}^{2\pi} [u \cos y - \cos u]_0^{\pi} dy$$

$$= \int_{-\pi}^{2\pi} [\pi \cos y - \cos \pi + \cos 0] dy$$

$$= \int_0^{2\pi} [\pi \cos y + 1 + 1] dy$$

$$= [\pi \sin y + 2y]_0^{2\pi}$$

$$= \pi (\sin 2\pi - \sin 0) + 2(2\pi - 0)$$

$$= \pi \times 0 + 2(\pi)$$

$$= 2\pi \# -$$

$$= \cancel{\pi \sin 2\pi + 8}$$

$$\text{e: } \int_1^2 \int_y^2 y^2 du dy$$

$$= \int_1^2 [y^2 - y] dy$$

$$= \left[\frac{y^3}{3} - \frac{y^2}{2} \right]_1^2$$

$$= \left(\frac{2 \times 2 \times 2}{3} - \frac{1}{3} - \frac{2 \times 2}{2} - \frac{1}{2} \right)$$

$$= \frac{4}{3} - \frac{3}{2}$$

$$= \frac{14-9}{6}$$

$$= \frac{5}{6} \#.$$

* Triple integral.

Evaluate

$$a) \int_0^1 \int_u^1 \int_0^{y-u} dz \, dy \, dz$$

$$= \int_0^1 \int_u^1 [z]_0^{y-u} dy \, du = \int_0^1 \int_u^1 [y-u] dy \, du$$

$$= \int_0^1 \left[\frac{y^2}{2} - uy \right]_u^1 du = \int_0^1 \left[\frac{1}{2} - \frac{u^2}{2} - (u - u^2) \right] du$$

$$= \int_0^1 \left[\frac{1}{2} + \frac{u^2}{2} - u \right] du = \left[\frac{1}{2}u + \frac{u^3}{2 \times 3} - \frac{u^2}{2} \right]_0^1$$

$$= \frac{1}{2} + \frac{1}{6} - \frac{1}{2} = \frac{1}{6} \#$$

$$b) \int_0^1 \int_0^{1-z} \int_0^z dz \, dy \, dz$$

$$= \int_0^1 \int_0^{1-z} [u]_0^z dy \, dz = \int_0^1 \int_0^{1-z} (2) dy \, dz$$

$$= 2 \int_0^1 [y]_0^{1-z} dz = 2 \int_0^1 [1-z] dz$$

$$= 2 \left[z - \frac{z^2}{2} \right]_0^1$$

$$= 2 \left[1 - \frac{1}{2} \right]$$

$$= 2 \times \frac{1}{2}$$

$$= 1 \#.$$

$$\begin{aligned}
 & \text{c) } \int_0^1 \int_0^{1-y} \int_0^2 dz dy du = 1 \quad \text{d) } \int_0^2 \int_0^1 \int_0^{1-y} dz dy du = 1 \Rightarrow \text{Some} \\
 & \text{e) } \int_0^1 \int_0^{3-3u} \int_0^{3-3u-y} dz dy du \\
 & = \int_0^1 \int_0^{3-3u} [z]_0^{3-3u-y} dy du \\
 & = \int_0^1 \int_0^{3-3u} [3-3u-y] dy du \\
 & = \int_0^1 \left[3y - 3uy - \frac{y^2}{2} \right]_0^{3-3u} du \\
 & = \int_0^1 \left[\frac{6y - 6uy - y^2}{2} \right]_0^{3-3u} du \\
 & = \frac{1}{2} \int_0^1 (6(3-3u) - 6u(3-3u) - (3-3u)^2) du \\
 & = \frac{1}{2} \int_0^1 (18 - 18u - 18u + 18u^2 - 9 + 18u - 9u^2) du \\
 & = \frac{1}{2} \int_0^1 (9u^2 - 18u + 9) du \\
 & = \frac{1}{2} \left[\frac{9u^3}{3} - \frac{18u^2}{2} + 9u \right]_0^1 \\
 & = \frac{1}{2} [3u^3 - 9u^2 + 9u]_0^1 \\
 & = \frac{1}{2} [3 - 9 + 9] \\
 & = \frac{3}{2} \text{ Ans}
 \end{aligned}$$

$$f = \int_0^{\sqrt{2}} \int_0^{3y} \int_{u^2+3y^2}^{8-u^2-y^2} dz du dy$$

→ Solution:-

$$\begin{aligned}
 &= \int_0^{\sqrt{2}} \int_0^{3y} [z]_{u^2+3y^2}^{8-u^2-y^2} du dy \\
 &= \int_0^{\sqrt{2}} \int_0^{3y} [8-u^2-y^2-u^2-3y^2] du dy \\
 &= \int_0^{\sqrt{2}} \int_0^{3y} [8-2u^2-4y^2] du dy
 \end{aligned}$$

$$\begin{aligned}
 &= \int_0^{\sqrt{2}} [8u - 2\frac{u^3}{3} - 4uy^2]_0^{3y} dy \\
 &= \int_0^{\sqrt{2}} [8 \times 3y - 2 \times \frac{(3y)^3}{3} - 4 \cdot 3y \cdot y^2] dy
 \end{aligned}$$

$$\begin{aligned}
 &= \int_0^{\sqrt{2}} [24y - 18y^3 - 12y^3] dy \\
 &= \left[24\frac{y^2}{2} - \frac{18y^4}{4} - \frac{12y^4}{4} \right]_0^{\sqrt{2}}
 \end{aligned}$$

$$= \left[12y^2 - \frac{30y^4}{4} \right]_0^{\sqrt{2}}$$

$$= 12(\sqrt{2})^2 - \frac{30(\sqrt{2})^4}{4}$$

$$= 12 \times 2 - 30 \times \frac{4}{4}$$

$$= 24 - 30$$

$$= -6 \#.$$

$$g) \int_1^e \int_1^e \int_1^e \frac{1}{xyz} dx dy dz \quad \boxed{\int \frac{1}{u} du = \log u}$$

$$= \int_1^e \int_1^e [\log x]_1^e \frac{1}{yz} dy dz \quad \begin{cases} \log e = 1 \\ \log 1 = 0 \end{cases}$$

$$= \int_1^e \int_1^e [\log e - \log 1] \frac{1}{yz} dy dz$$

$$= \int_1^e \int_1^e [1 - 0] \frac{1}{yz} dy dz$$

$$= \int_1^e [\log y]_1^e \frac{1}{z} dz$$

$$= \int_1^e [\log e - \log 1] \frac{1}{z} dz$$

$$= \int_1^e [1 - 0] \frac{1}{z} dz$$

$$= [\log z]_1^e$$

$$= [\log e - \log 1]$$

$$= [1 - 0]$$

$$= 1 \cancel{\#}$$

$$h) \int_0^{2\pi} \int_0^1 \int_{-\frac{1}{2}}^{\frac{1}{2}} (r^2 \sin^2 \theta + z^2) dz r dr d\theta$$

$$= \int_0^{2\pi} \int_0^1 \left[z r^2 \sin^2 \theta + \frac{z^3}{3} \right]_{-\frac{1}{2}}^{\frac{1}{2}} r dr d\theta$$

$$= \int_0^{2\pi} \int_0^1 \left[\frac{1}{2} r^2 \sin^2 \theta - \left(\frac{1}{2} \right) r^2 \sin^2 \theta + \left(\frac{1}{2} \right)^3 - \left(-\frac{1}{2} \right)^3 \right] r dr d\theta$$

$$\begin{aligned}
 &= \int_0^{2\pi} \int_0^1 \left[\frac{r^2 \sin^2 \theta}{2} + \frac{r^2 \sin^2 \theta}{2} + \frac{1}{24} + \frac{1}{24} \right] r dr d\theta \\
 &= \int_0^{2\pi} \int_0^1 \left[r^2 \sin^2 \theta + \frac{1}{12} \right] r dr d\theta \\
 &= \int_0^{2\pi} \int_0^1 \left[r^3 \sin^2 \theta + \frac{r}{12} \right] dr d\theta \\
 &= \int_0^{2\pi} \left[\frac{r^4 \sin^2 \theta}{4} + \frac{r^2}{12 \times 2} \right]_0^1 d\theta \\
 &= \int_0^{2\pi} \left[\frac{\sin^2 \theta}{4} + \frac{1}{24} \right] d\theta \\
 &= \int_0^{2\pi} \left[\frac{1 - \cos 2\theta}{2 \times 4} + \frac{1}{24} \right] d\theta \\
 &= \left[\frac{[\theta]}{8} - \frac{[\sin 2\theta]}{2 \times 8} + \frac{1}{24} [\cos \theta] \right]_0^{2\pi} \\
 &= \frac{2\pi}{8} - \frac{1}{16} \times (0 - 0) + \frac{1}{24} \times 2\pi \\
 &= \frac{\pi}{4} + \frac{\pi}{12} \\
 &= \frac{3\pi + \pi}{12} \\
 &= \frac{4\pi}{12} \\
 &= \frac{\pi}{3} \# .
 \end{aligned}$$

$$\begin{aligned}
 & \text{Q3} \int \int \int dz \, r \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^{\pi/2} \int_0^{\sqrt{3+24r^2}} [z] \, r \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^{\pi/2} [3+24r^2] \times r \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^{\pi/2} [3r + 24r^3] \, dr \, d\theta \\
 &= \int_0^{2\pi} \left[\frac{3r^2}{2} + \frac{24r^4}{4} \right]_0^{\pi/2} \, d\theta \\
 &= \int_0^{2\pi} \left[\frac{3 \frac{\theta^2}{4\pi^2}}{2} + \frac{24 \frac{\theta^4}{16\pi^4}}{4} \right] \, d\theta \\
 &= \int_0^{2\pi} \left[\frac{3\theta^2}{8\pi^2} + \frac{3\theta^4}{8\pi^4} \right] \, d\theta \\
 &= \frac{3}{8\pi^2} \left[\frac{\theta^3}{3} \right]_0^{2\pi} + \frac{3}{8\pi^4} \left[\frac{\theta^5}{5} \right]_0^{2\pi} \\
 &= \frac{3}{8\pi^2} \times \frac{1}{3} (2\pi)^3 + \frac{3}{8\pi^4} \times \frac{1}{5} (2\pi)^5 \\
 &= \frac{8\pi^3}{8\pi^2} + \frac{32 \times 3 \times \pi^5}{40 \pi^4} \\
 &= \pi + \frac{96}{40} \pi \\
 &= \pi + \frac{12\pi}{5} = \frac{17\pi}{5} \#.
 \end{aligned}$$

$$5. \int_{-1}^1 \int_0^{2\pi} \int_0^{1+\cos\theta} 4r \, dr \, d\theta \, dz.$$

$$= \int_{-1}^1 \int_0^{2\pi} \left[\frac{4r^2}{2} \right]_0^{1+\cos\theta} \, d\theta \, dz$$

$$= \int_{-1}^1 \int_0^{2\pi} \left[2r^2 \right]_0^{1+\cos\theta} \, d\theta \, dz \quad \because \cos^2\theta = \frac{1+\cos 2\theta}{2}$$

$$= \int_{-1}^1 \int_0^{2\pi} 2 \left[1 + \cos\theta \right]^2 \, d\theta \, dz$$

$$= \int_{-1}^1 \int_0^{2\pi} 2 (1 + 2\cos\theta + \cos^2\theta) \, d\theta \, dz$$

$$= \int_{-1}^1 \int_0^{2\pi} 2 \cdot (1 + 2\cos\theta + \frac{1 + \cos 2\theta}{2}) \, d\theta \, dz$$

$$= 2 \int_{-1}^1 \left[[\theta]_0^{2\pi} + 2 [\sin\theta]_0^{2\pi} + \frac{1}{2} \left\{ [\theta]_0^{2\pi} + [\frac{1}{2} \sin 2\theta]_0^{2\pi} \right\} \right] dz$$

$$= 2 \int_{-1}^1 (2\pi - 0) + 2 [\sin 2\pi - \sin 0] + \frac{1}{2} [2\pi - 0] + \frac{1}{4} [\sin 2 \cdot 2\pi - \sin 2 \cdot 0] \, dz$$

$$= 2 \int_{-1}^1 (2\pi + \frac{2\pi}{2} + \frac{1}{4} \times 6) \, dz$$

$$= 2 \int_{-1}^1 3\pi \, dz$$

$$= 6\pi \left[z \right]_{-1}^1$$

$$= 6\pi [1 + 1]$$

$$= 12\pi \quad \#$$

Integration by parts

$$= \int u \cdot v du = u \int v du - \int \left\{ \frac{du}{du} \int v du \right\} du$$

$$= u \int v du - \int \left\{ \frac{du}{du} \left\{ \int v du \right\} \right\} du$$

= first part \times integration of second - Integration $\{$ derivatives
 of ~~second~~ first \times integration of second $\}$

Rule for integration by parts :-

I LATE Rule

I \rightarrow Inverse $\rightarrow \sin^{-1} u \dots$

L \rightarrow Logarithm $\rightarrow \log$

A \rightarrow Algebraic $\rightarrow x, y, z \dots$

T \rightarrow Trigonometric $\rightarrow \sin u, \cos u \dots$

E \rightarrow exponential $\rightarrow e^u$

e.g.

$$\int u \cdot e^u du, = u \int e^u du - \int \left\{ \frac{du}{du} \int e^u du \right\} du$$

\rightarrow here,

$$u = u, v = e^u \Rightarrow u e^u - \int e^u du \\ \therefore u e^u - e^u + C$$

$$\Rightarrow u \int e^u du - \int \left\{ \frac{du}{du} \int e^u du \right\} du$$

$$\Rightarrow u \cdot e^u - \int e^u du$$

$$\Rightarrow u \cdot e^u - e^u + C \#$$

e.g. $\int y \cdot e^y dy$

$$= y \int e^y dy - \int \left\{ \frac{dy}{dy} \int e^y dy \right\} dy$$

$$= y e^y - e^y + C \#$$

* Evaluate:

$$\ln 8 \ln y$$

$$* \int_1^{\ln 8} \int_0^{\ln y} e^{u+y} du dy$$

$$= \int_1^{\ln 8} \int_0^{\ln y} e^u \cdot e^y du dy$$

$$= \int_1^{\ln 8} \left[e^y \cdot [e^u]_0^{\ln y} \right] dy$$

$$= \int_1^{\ln 8} \left[e^y \cdot [e^{\ln y} - e^0] \right] dy$$

$$= \int e^y \cdot y dy$$

$$= \int_1^{\ln 8} \left[e^y \cdot (y-1) \right] dy$$

$$= y \{ e^y dy - \int \frac{dy}{y} \{ e^y dy \} \}$$

$$= \int_1^{\ln 8} [e^y \cdot y - e^y] dy$$

$$= y \cdot e^y - e^y + C$$

$$= \left[ye^y - e^y \right]_1^{\ln 8} - \left[e^y \right]_1^{\ln 8}$$

$$= \left[ye^y - 2e^y \right]_1^{\ln 8}$$

$$e^{\ln 8}(\ln 8 - 1) - e^1 + e^1 = 8 \ln 8$$

$$= \left[\ln 8 e^{\ln 8} - 1 \times e^1 - 2e^{\ln 8} + 2e^1 \right]$$

$$= 8 \ln 8 - e - 2 \times 8 + 2e$$

$$= 8 \ln 8 - 16 + e \quad \underline{\text{Answer}}$$

Fubini's Theorem:-

1) First Form:-

If $f(u, y)$ is continuous on region R :
 $a \leq u \leq b$ and $c \leq y \leq d$, then

$$y=d \quad u=b$$

$$\iint_R f(u, y) dA = \int_{y=c}^d \int_{u=a}^b f(u, y) du dy$$

$$= \int_{x=a}^b \int_{y=c}^d f(u, y) dy du$$

This form gives the volume.

* evaluate:-

$$f(u, y) = 1 - 6u^2y, -1 \leq u \leq 1, 0 \leq y \leq 2$$

→ Solution

$$\int_{y=0}^2 \int_{u=-1}^1 (1 - 6u^2y) du dy = \int_{u=-1}^1 \int_{y=0}^2 (1 - 6u^2y) dy du$$

$$= \int_0^2 \int_{-1}^1 (1 - 6u^2y) du dy$$

$$= \int_0^2 \left[u - \frac{6u^3y}{3} \right]_{-1}^1 dy$$

$$= \int_0^2 \left[1 - (-1) - \left\{ 2y - 2(-1)^3 y^2 \right\} \right] dy$$

$$= \int_0^2 (2 - 2y - 2y^2) dy$$

$$= \left[2y - \frac{4y^2}{2} \right]_0^2$$

$$= 2 \times 2 - 2(2)^2 - 0$$

$$= 4 - 8$$

$$= -4$$

At second,

$$\int_{-1}^1 \int_0^2 (1 - 6u^2y) dy du$$

$$= \int_{-1}^1 \left[y - \frac{6u^2y^2}{2} \right]_0^2 du$$

$$= \int_{-1}^1 \left[2 - 3u^2(2)^2 \right] du$$

$$= \int_{-1}^1 \left[2 - 12u^2 \right] du$$

$$= \left[2u - \frac{12u^3}{3} \right]_{-1}^1$$

$$= \left[2u - 4u^3 \right]_{-1}^1$$

$$= 2 \times 1 - 2 \times (-1) - \{ 4 \times (1)^3 - 4 \times (-1)^3 \}$$

$$= 2 + 2 - (4 + 4)$$

$$= 4 - 8$$

$$= -4$$

$$\iiint_R dxdydz = \text{Volume}$$

$$\iint_R dxdy = \text{Area}$$

\therefore Both $\int dy du$ and $\int dx dy$ have same value.

* Fubini's Theorem (stronger Form)

Function $f(u, y)$ is continuous on region R , then,a. if R is defined by $a \leq u \leq b$, $g_1(u) \leq y \leq g_2(u)$, then,

$$x=b \quad y=g_2(u)$$

$$\iint_R f(u, y) dA = \int_a^b \int_{g_1(u)}^{g_2(u)} f(u, y) dy du$$

b. if R is defined by $a \leq y \leq b$, $g_1(y) \leq u \leq g_2(y)$

$$y=b \quad u=g_2(y)$$

$$\iint_R f(u, y) dA = \int_a^b \int_{g_1(y)}^{g_2(y)} f(u, y) du dy$$

Find the area between $y^2 = 4u$ and $u^2 = 2y$

→ Solution

$$y^2 = 4u \quad \dots \dots \text{(i)}$$

$$u^2 = 2y$$

$$y = \frac{u^2}{2} \quad \dots \dots \text{(ii)}$$

Solving eqn (i) and (ii), we get,

$$\left(\frac{u^2}{2}\right)^2 = 4u$$

$$\frac{u^4}{4} = 4u$$


$$\text{or } u^4 = 16u$$

$$\text{or } u(u^3 - 16) = 0$$

Either $u = 0$

$$u^3 - 16 = 0$$

$$\therefore u = \sqrt[3]{16} = 2.5$$



$$\int_{u=0}^{u=2.5} \int_{y=0}^{y=\sqrt{4u}} dy du$$

from Single integration:-

$$u=2.5 \quad u=2.5$$

$$= \text{Area} = \int_{u=0}^{u=2.5} \sqrt{4u} du - \int_{u=0}^{u=2.5} \frac{u^2}{2} du$$

OR

Double integration:-

$$u=2.5 \quad y=\sqrt{4u}$$

$$\text{Area} = \int_{u=0}^{u=2.5} \int_{y=0}^{y=\frac{u^2}{2}} dy du$$

$$= \int_{u=0}^{u=2.5} \left[y \right]_{u/2}^{\sqrt{4u}} du$$

$$u=2.5 \quad u=2.5$$

$$= \int_{u=0}^{u=2.5} \left(\sqrt{4u} - \frac{u^2}{2} \right) du$$

$$u=2.5 \quad u=2.5$$

$$= \int_{u=0}^{u=2.5} \left(\sqrt{4u} - \frac{u^2}{2} \right) du$$

$$= \sqrt{4} \left[\frac{u^{3/2}}{3/2} - \frac{\sqrt{4}u^3}{2 \times 3} \right]_0^{2.5} = \sqrt{4} \left(\frac{u^{3/2}}{3/2} \right)_0^{2.5} - \left[\frac{u^3}{6} \right]_0^{2.5}$$

$$= 2 \times \frac{2}{3} \left[(2.5)^{3/2} - \frac{\sqrt{4}}{6} (2.5)^3 \right] = 2 \times \frac{2}{3} \times (3.95) - \frac{(2.5)^3}{6}$$

$$= 1.33 \left(3.95 - 2.60 \times \sqrt{4} \right)$$

$$= 1.33 \times \cancel{2.60}$$


=

$$= \frac{4}{3} \times 3.95 - 2.60$$

$$= 5.2 - 2.60$$

$$= 2.66 \cancel{.}$$

Find the Volume of Prism whose base is the triangle in the xy -plane bounded by the x -axis and the line $y=u$ and $u=1$ and whose lies in the plain $Z=f(u,y)=3-u-y$



$$= \text{Volume} = \int_{u=0}^{u=1} \int_{y=0}^{y=u} f(u,y) dy du$$

$$= \int_{u=0}^{u=1} \int_{y=0}^{y=u} (3-u-y) dy du$$

$$= \int_{u=0}^{u=1} \left[3y - uy - \frac{y^2}{2} \right]_0^u du$$

$$= \int_{u=0}^{u=1} \left[3u - u^2 - \frac{u^2}{2} \right] du$$

$$= \left[\frac{3u^2}{2} - \frac{u^3}{3} - \frac{u^3}{2 \times 3} \right]_0^1$$

$$= \frac{3}{2} - \frac{1}{3} - \frac{1}{6}$$

$$= \frac{9-2-1}{6}$$

$$= \frac{6}{6}$$

$$= 1 \text{ cubic units}$$

Required Volume of prism

* Find the Volume of Solid under the surface $z = f(u, y) = u^2 + y^2$ over the triangular region whose vertices are $(0,0)$, $(1,0)$ and $(0,1)$

→ Solution:

$$\text{Point } (u_1, y_1) = (1, 0)$$

$$\text{Point } (u_2, y_2) = (0, 1)$$

equation of a line is given by,

$$y - y_1 = m(u - u_1)$$

$$\textcircled{a.} \quad (y - 0) = \frac{y_2 - y_1}{u_2 - u_1} (u - 1)$$

$$\textcircled{b.} \quad (y - 0) = \frac{(1 - 0)}{(0 - 1)} (u - 1)$$

$$\textcircled{c.} \quad y = -u + 1$$

$$\therefore y = (1 - u)$$

$$\text{Now, } u = 1 \quad y = (1 - u)$$

$$\text{Volume} = \int_{u=0}^1 \int_{y=0}^{1-u} u^2 + y^2 \, dy \, du$$

$$= \int_0^1 \left[8u^2 + \frac{y^3}{3} \right]_0^{1-u} \, du \quad \text{or use } \frac{(1-u)^{3+4}}{3 \times 4 \times (-1)} \, du$$


$$= \int_0^1 \left[(1-u)u^2 + \frac{(1-u)^3}{3} \right] \, du$$

$$= \int_0^1 \left[u^2 - u^3 + \frac{1}{3} (1)^3 - 3(1)^2 \cdot u + 3(1) \cdot u^2 - u^3 \right] \, du$$

$$= \frac{1}{3} \int_0^1 (3u^2 - 3u^3 + 1 - 3u + 3u^2 - u^3) \, du$$

$$= \frac{1}{3} \left[\frac{3u^3}{3} - \frac{3u^4}{4} + u - 3u^2 + \frac{3u^3}{3} - \frac{u^4}{4} \right]_0^1$$

$$= \frac{1}{3} \times \left(1 - \frac{3}{4} + \frac{1}{4} - \frac{3}{2} + 1 - \frac{1}{4} \right) = \frac{1}{3} \times \left[\frac{4-3+4-6+4-1}{4} \right] = \frac{1}{3} \times \frac{2}{4^2} = \frac{1}{3} \times \frac{1}{2} = \frac{1}{6}$$



* Find volume, $z = f(u, v) = uv$
vertices $(1, 1)$, $(4, 1)$ and $(1, 2)$

→ Solution,

for point $(1, 1) = (u_1, v_1)$
 $(4, 1) = (u_2, v_1)$

eqn of line is

$$y - y_1 = m(u - u_1)$$

$$\text{or } (y - 1) = \frac{y_2 - y_1}{u_2 - u_1} (u - 1)$$

$$\text{or } y - 1 = \frac{1 - 1}{4 - 1} (u - 1)$$

$$\text{or } y = 1$$

∴ lower limit $y = 1$

for point $(1, 2) = (u_1, v_2)$ and $(4, 1) = (u_2, v_1)$

eqn of line is

$$(y - 2) = \frac{1 - 2}{4 - 1} (u - 1)$$

$$\text{or } y - 2 = -\frac{1}{3} (u - 1)$$

$$\text{or } 3y - 6 = -u + 1$$


$$\text{or } 3y = 7 - u$$

$$\therefore y = \frac{7-u}{3}$$

$$u=4 \quad y=\frac{7-u}{3}$$

$$\text{Volume} = \int_{u=1}^{u=4} \int_{y=1}^{y=\frac{7-u}{3}} uv \, dy \, du$$

$$= \int_{1}^{4} \left[u \frac{y^2}{2} \right]_{1}^{\frac{7-u}{3}} du$$



$$7-y_1 = \frac{y_1 - y_2}{u_1 - u_2} (u_1 - u_2)$$

$$= \int_1^4 \left[\frac{u \cdot \left(\frac{7-u}{3}\right)^2}{2} - \frac{u}{2} \right] du$$

$$= \int_1^4 \left[\frac{u(49-14u+u^2)}{9 \times 2} - \frac{u}{2} \right] du$$

$$= \int_1^4 \left[\frac{1}{18} (49u - 14u^2 + u^3) - \frac{u}{2} \right] du$$

$$= \frac{1}{18} \int_1^4 [49u - 14u^2 + u^3 - 9u] du$$

$$= \frac{1}{18} \int_1^4 [40u - 14u^2 + u^3] du$$

$$= \frac{1}{18} \left[\frac{40u^2}{2} - \frac{14u^3}{3} + \frac{u^4}{4} \right]_1^4$$

$$= \frac{1}{18} \left[20u^2 - \frac{14u^3}{3} + \frac{u^4}{4} \right]_1^4$$

$$= \frac{1}{18} \left[20 \times (4)^2 - \frac{14 \times (4)^3}{3} + \frac{(4)^4}{4} - 20 \times 1 + \frac{14 \times 1}{3} - \frac{1}{4} \right]$$

$$= \frac{1}{18} [320 - 298.66 + 64 - 20 + 4.66 - 0.25]$$

$$= \frac{69.75}{18}$$

$$= 3.87$$

∴ Required Volume is 3.87 cubic unit

* Find the area of parallelogram $y = u^2$ and $y = u+2$.

→ Solution,

$$y = u^2 \quad \text{--- (i)}$$

$$y = u+2 \quad \text{--- (ii)}$$

Solving eqn ① & ②

$$u^2 = u+2$$

$$\therefore u^2 - u - 2 = 0$$

$$\therefore u^2 - 2u + u - 2 = 0$$

$$\therefore u(u-2) + 1(u-2) = 0$$

$$\therefore (u-2)(u+1) = 0$$

Either,

$$u = -1, y_1 = 1$$

$$u = 2, y_2 = 4$$

eqn of parabola is $y = u^2$

when $y = 0, u = 0$


$$y = 1, u = \pm 1$$

$$y = 4, u = \pm 2$$


$$x = 2, y = u+2$$

$$= \text{Area} = \int_{u=-1}^{u=2} \int_{y=u^2}^{y=u+2} dy du$$

$$\begin{aligned}
 &= \int_{u=-1}^{u=2} \left[y \right]_{u^2}^{u+2} du \\
 &= \int_{-1}^2 (u+2 - u^2) du \\
 &= \left[\frac{u^2}{2} + 2u - \frac{u^3}{3} \right]_{-1}^2
 \end{aligned}
 \quad \left\{
 \begin{aligned}
 &= \frac{4}{2} + 4 - \frac{8}{3} - \frac{1}{2} + 2 - \frac{1}{3} \\
 &= \frac{12 + 24 - 16 - 3 + 12 - 2}{6} \\
 &= \frac{27}{6} \\
 &= 4.5
 \end{aligned}
 \right.$$



* Find the volume of solid in the first octant bounded by the co-ordinate planes, the paraboloid, $z = u^2 + y^2 + 1$, and the plane $2u+y=2$



$$\text{In eq } 2u+y=2$$

when

$$u=0, y=2$$

$$u=1, y=0$$

$$u=2, y=-2$$

$$\therefore \text{Volume (V)} = \int_{u=0}^{u=1} \int_{y=0}^{y=2-2u} f(u, y) dy du$$

$$u=1, y=2-2u$$

$$u=0, y=0$$

$$= \int_{u=0}^{u=1} \int_{y=0}^{y=2-2u} (u^2 + y^2 + 1) dy du$$

$$= \int_{u=0}^{u=1} \left[u^2 y + \frac{y^3}{3} + y \right]_{0}^{2-2u} du$$

$$= \int_{u=0}^{u=1} \left[u^2(2-2u) + \frac{(2-2u)^3}{3} + (2-2u) \right] du$$


$$= \int_{u=0}^{u=1} \left[2u^2 - 2u^3 + (2-3u)^2 \cdot 2u + 3 \cdot 2(2u)^2 - (2u)^3 + (2-2u) \right] du$$

$$= \frac{1}{3} \int_{u=0}^{u=1} \left[6u^2 - 6u^3 + 8 - 24u + 24u^2 - 8u^3 + 6 - 6u \right] du$$

$$= \frac{1}{3} \int_{u=0}^{u=1} \left[14 - 30u + 30u^2 - 14u^3 \right] du$$

$$\begin{aligned}
 &= \frac{1}{3} \left\{ [14u]^{\frac{1}{2}} - 30 \left[\frac{u^2}{2} \right]_0^{\frac{1}{2}} + 30 \left[\frac{u^3}{3} \right]_0^{\frac{1}{2}} - 14 \left[\frac{u^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^{\frac{1}{2}} \right\} \\
 &= \frac{1}{3} \left\{ 14 - 15 + 10 - \frac{7}{2} \right\} \\
 &= \frac{1}{3} \left(9 - \frac{7}{2} \right) \\
 &= \frac{1}{3} \left(\frac{18 - 7}{2} \right) \\
 &= \frac{11}{6} \text{ cubic Unit} \\
 &\therefore \text{Required Volume of Solid is } \frac{11}{6} \text{ cubic Unit}
 \end{aligned}$$

* Find the volume of region bounded by paraboloid $z = u^2 + y^2$ and below by the triangle enclosed by the line $y = u$, $u = 0$ and $u + y = 2$ in the uv plane.



$$\text{for } u + y = 2$$

$$\text{when } u = 0, y = 2$$

$$\text{when } u = 1, y = 1$$

$$\text{when } u = 2, y = 0$$

$$\begin{aligned}
 &X \quad u = 1 \quad y = 2 - u \\
 &Y \quad u = 0 \quad y = u \\
 \text{Volume} &= \int_{u=0}^{u=1} \int_{y=u}^{y=2-u} (u^2 + y^2) dy du
 \end{aligned}$$

$$\begin{aligned}
 &= \int_{u=0}^{u=1} \left[u^2 y + \frac{y^3}{3} \right]_{y=u}^{y=2-u} du
 \end{aligned}$$

$$\begin{aligned}
 &= \int_{u=0}^{u=1} \left[u^2(2-u) - u^2(u) + \frac{(2-u)^3}{3} - \frac{(u)^3}{3} \right] du
 \end{aligned}$$

$$\begin{aligned}
 &= \int_{u=0}^{u=1} \left[2u^2 - u^3 - u^3 + (2)^3 - 3 \cdot (2)^2 \cdot u + 3(2)(u)^2 - (u)^3 - \frac{u^3}{3} \right] du \\
 &= \frac{1}{3} \int_{u=0}^{u=1} \left[6u^2 - 3u^3 - 3u^3 + 8 - 12u + 6u^2 - u^3 - u^3 \right] du \\
 &= \frac{1}{3} \int_{u=0}^{u=1} \left[8 - 12u + 12u^2 - 8u^3 \right] du \\
 &= \frac{1}{3} \left\{ 8[u]_0^1 - 12 \left[\frac{u^2}{2} \right]_0^1 - 8 \left[\frac{u^4}{4} \right]_0^1 + 12 \left[\frac{u^3}{3} \right]_0^1 \right\} \\
 &= \frac{1}{3} \left\{ 8 \times 1 - 12 \times \frac{1}{2} - 8 \times \frac{1}{4} + 12 \times \frac{1}{3} \right\} \\
 &= \frac{1}{3} \left\{ 8 - 6 - 2 + 4 \right\} \\
 &= \frac{1}{3} (12 - 8) \\
 &= \frac{4}{3} \text{ Cubic unit.} \\
 &\therefore \text{Required Volume of Region is } \frac{4}{3} \text{ Cubic unit.}
 \end{aligned}$$

* Find the Volume of Solid in the first octant bounded by coordinate planes, the planes, $u=3$ and the parabolic cylinder, $z = 4 - y^2$.


$$\text{eqn } z = 4 - y^2$$

$$\text{when } z = 0$$

$$y^2 = 4$$

$$u=3 \therefore y = \pm 2$$

$$\rightarrow V = \int_{u=0}^{u=3} \int_{y=-2}^{y=2} (4 - y^2) dy du$$



$$u=3 \\ = \int \left[4y - \frac{y^3}{3} \right]_0^3 du$$

$$u=0$$

$$u=3 \\ = \int \left[8 - \frac{8}{3} \right] du$$

$$u=0$$

$$u=3 \\ = \int \left[\frac{16}{3} \right] du$$

$$u=0$$

$$= \frac{16}{3} \left[u \right]_0^3$$

$$= \frac{16 \times 3}{3}$$

$$= 16 \text{ cubic unit}$$

Hence the required Volume of solid is
cubic unit.

Volume by:

a) Double integral. $= \iint_R f(u, y) dy, du$

b) Triple integral $= \iiint_R dz dy du$

* Find the Volume of region enclosed by Surface
 $z = u^2 + 3y$ and $z = 8 - u^2 - y^2$.

$\rightarrow 601^n$,

STEP 1: First find which is upper and lower limit
in z .

for this, put the value of u and y as 0.

then,

which is greater that is Upper & which value is lower that is lower limit.

i.e.

$$z = 0^2 + 30^2 = 0 \rightarrow \text{lower limit} = u^2 + 3y^2$$

$$z = 8 - 0^2 - 0^2 = 8 \rightarrow \text{Upper limit} = 8 - u^2 - y^2$$

STEP 2: To find limit of y ,

Solve the equation as,

$$u^2 + 3y^2 = 8 - u^2 - y^2$$

$$\therefore 2u^2 + 4y^2 = 8$$

$$\therefore u^2 + 2y^2 = 4$$

$$\therefore 2y^2 = 4 - u^2 \quad \text{which is eqn of ellipse, i.e. } \frac{u^2}{(2)^2} + \frac{y^2}{(\sqrt{2})^2} = 1$$

$$\therefore y^2 = \frac{4 - u^2}{2}$$

$$\therefore a=2, b=\sqrt{2}$$

$$\therefore y = \pm \sqrt{\frac{4-u^2}{2}}$$


$$\text{For } y, \text{ Upper limit} = +\sqrt{\frac{4-u^2}{2}}$$

$$\text{lower limit} = -\sqrt{\frac{4-u^2}{2}}$$

STEP 3: To find limit of u .

by solving,

$$\text{eqn is ellipse, i.e. } \frac{u^2}{(2)^2} + \frac{y^2}{(\sqrt{2})^2} = 1$$



in this eqn

$$a=2$$

$$\therefore \text{Upper limit} = +2$$

$$\therefore \text{lower limit} = -2$$

$$u=+2 \quad y=+\sqrt{\frac{4-u^2}{2}} \quad z=8-u^2-y^2$$

$$u=-2 \quad y=-\sqrt{\frac{4-u^2}{2}} \quad z=u^2+3y^2$$

Now, Volume = $\int_{u=-2}^{u=+2} \int_{y=-\sqrt{\frac{4-u^2}{2}}}^{y=+\sqrt{\frac{4-u^2}{2}}} dz dy du$

$$= \int_{-2}^{+2} \int_{-\sqrt{\frac{4-u^2}{2}}}^{\sqrt{\frac{4-u^2}{2}}} (8-u^2-y^2-u^2-3y^2) dy du$$

$$= \int_{-2}^{+2} \int_{-\sqrt{\frac{4-u^2}{2}}}^{\sqrt{\frac{4-u^2}{2}}} (8-2u^2-4y^2) dy du$$

$$= \int_{-2}^{+2} \left[8y - 2u^2y - \frac{4}{3}y^3 \right]_{-\sqrt{\frac{4-u^2}{2}}}^{\sqrt{\frac{4-u^2}{2}}} du$$

$$= \int_{-2}^{+2} \left[(8-2u^2) \times 2 \sqrt{\frac{4-u^2}{2}} - \frac{4}{3} \times 2 \left(\frac{4-u^2}{2} \right)^{\frac{3}{2}} \right] du$$

$$= 2 \int_{-2}^{+2} \left[2(4-u^2)^{\frac{1}{2}} (4-u^2)^{\frac{1}{2}} - \frac{4}{3} \times 2 \left(\frac{4-u^2}{2} \right)^{\frac{3}{2}} \right] du$$

$$= \int_{-2}^{+2} \left[\frac{4}{\sqrt{2}} (4-u^2)^{\frac{3}{2}} - \frac{4}{3} \times 2 \left(\frac{4-u^2}{2} \right)^{\frac{3}{2}} \right] du$$

(false common)

$$= \int_{-2}^{+2} \frac{4}{\sqrt{2}} (4-u^2)^{\frac{3}{2}} \left(1 - \frac{1}{3} \right) du$$

$$= \frac{3-1}{3} \times \frac{2}{3}$$

$y=+a$
if $\int dy$
 $y=-a$
then,
 $= [y]_{-a}^a$

= 2a
if same lower &
upper limit with +ve
& neg. sign we can
write $2 \times$ that limit

which is in the form of $(a^2 - u^2)$. So $(2^2 - u^2) \therefore a=2$
 put $u=a$ since $u=a$
 $21 = 28 - 7$

differentiating both sides with respect to α .

$$\frac{dn}{dc} = 2 \frac{ds \cos \alpha}{dc}$$

$$\therefore d\mu = 2 \cos \alpha d\alpha$$

when $n=2$

$$2 = 2 \sin \alpha$$

$$\text{as, } 0 = \sin^{-1}(1) = \frac{\pi}{2}$$

when $\kappa = 0$

$$0 = 2 \sin \alpha$$

$$\therefore \varphi = \sin'(0) = 0$$

$$\therefore V = \frac{8\sqrt{2}}{3} \int_{\pi/2}^{3\pi/2} (4 - 4\sin^2\alpha)^{3/2} \cdot 2 \cdot \cos\alpha \, d\alpha$$

$$= \frac{8\sqrt{2}}{3} \int_0^{\pi/2} 4^{\frac{3}{2}} (1 - \sin^2 \alpha)^{\frac{3}{2}} \cdot 2 \cos \alpha \, d\alpha$$

$$= \frac{8\sqrt{2}}{3} \int_0^{\pi/2} 8 \cos^2 \theta \cdot 2 \cos \theta \, d\theta$$

$$= \frac{8\sqrt{2} \times 16}{3} \int_0^{\frac{\pi}{2}} \cos^4 \alpha \, d\alpha \quad \because \cos^2 \alpha = \frac{1 + \cos 2\alpha}{2}$$

$$= \frac{8 \times 16\sqrt{2}}{3} \int_0^{\frac{\pi}{2}} (\cos^2 \alpha)^2 \, d\alpha$$

$$= \frac{8 \times 16\sqrt{2}}{3} \int_0^{\frac{\pi}{2}} \left(\frac{1 + \cos 2\alpha}{2} \right)^2 \, d\alpha$$

$$= \frac{8 \times 16\sqrt{2}}{3} \int_0^{\frac{\pi}{2}} \left\{ \frac{1}{4} (1 + 2\cos 2\alpha + \cos^2 2\alpha) \right\} \, d\alpha$$

$$= \frac{8 \times 16\sqrt{2}}{3} \int_0^{\frac{\pi}{2}} \left\{ \frac{1}{4} (1 + 2\cos 2\alpha + \frac{1 + \cos 4\alpha}{2}) \right\} \, d\alpha \quad \text{L.C.M.}$$

$$= \frac{8 \times 16\sqrt{2}}{3} \int_0^{\frac{\pi}{2}} \frac{1}{8} (2 + 4\cos 2\alpha + 1 + \cos 4\alpha) \, d\alpha$$

$$= \frac{8 \times 16\sqrt{2}}{3} \times \frac{1}{8} \int_0^{\frac{\pi}{2}} (3 + 4\cos 2\alpha + \cos 4\alpha) \, d\alpha$$

$$= \frac{16\sqrt{2}}{3} \left[3 \left[\alpha \right]_0^{\frac{\pi}{2}} + 4 \cdot \left[\frac{\sin 2\alpha}{2} \right]_0^{\frac{\pi}{2}} + \left[\frac{\sin 4\alpha}{4} \right]_0^{\frac{\pi}{2}} \right]$$

$$= \frac{16\sqrt{2}}{3} \left[\frac{3\pi}{2} + 4 \left[\frac{\sin 2 \cdot \frac{\pi}{2}}{2} - \frac{\sin 2 \cdot 0}{2} \right] + \left[\frac{\sin^4 \frac{\pi}{2} - \sin^4 0}{4} \right] \right]$$

$$= \frac{16\sqrt{2}}{3} \left(\frac{3\pi}{2} + 4 \left(\frac{0}{2} - \frac{0}{2} \right) + \left(\frac{0 - 0}{4} \right) \right)$$

$$= \frac{16\sqrt{2}}{3} \times \frac{8\pi}{2} \quad \text{cubic unit}$$

$$= 8\pi\sqrt{2} \quad \text{Answer}$$

∴ Required Volume is $8\pi\sqrt{2}$ cubic unit

* find the Volume bounded above by paraboloid $Z = 5 - u^2 - y^2$ and $Z = 4u^2 + 4y^2$.

→ Solution,

for Upper and lower limit in Z , put $u=0, y=0$

$$Z = 5 - u^2 - y^2 = 5 - 0 - 0 = 5 \Rightarrow \text{Upper limit}$$

$$Z = 4u^2 + 4y^2 = 0 + 0 = 0 \Rightarrow \text{lower limit.}$$

for Upper limit and lower limit in y , solve 2 eqⁿ

$$4u^2 + 4y^2 = 5 - u^2 - y^2$$

$$\text{a. } 5u^2 + 5y^2 = 5$$

$$\text{b. } u^2 + y^2 = 1 \text{ which is eq}^n \text{ of circle}$$

$$\text{c. } y^2 = 1 - u^2 \quad u^2 + y^2 = a^2$$

$$\therefore y = \pm \sqrt{1 - u^2} \quad \therefore a = 1$$

$$\therefore \text{Upper limit in } y = +\sqrt{1 - u^2}$$

$$\therefore \text{lower limit in } y = -\sqrt{1 - u^2}$$


for Upper and lower limit for u .

$$\text{eq}^n \text{ is } u^2 + y^2 = 1$$

∴ eqⁿ of circle

so,

$$u^2 + y^2 = 1 \quad \therefore \text{Upper limit} = +1 \\ \text{lower limit} = -1$$



$$u = \pm 1 \quad y = \pm \sqrt{1-u^2} \quad z = 5 - u^2 - y^2$$

$$\therefore \text{Volume} = \int_{u=-1}^{u=1} \int_{y=-\sqrt{1-u^2}}^{\sqrt{1-u^2}} \int_{z=4u^2+4y^2}^{5-u^2-y^2} dz dy du$$

$$= \int_{-1}^{1} \int_{-\sqrt{1-u^2}}^{\sqrt{1-u^2}} \left[z \right]_{4u^2+4y^2}^{5-u^2-y^2} dy du$$

$$= 5 - u^2 - y^2 - 4u^2 - 4y^2$$

$$= \int_{-1}^{1} \int_{-\sqrt{1-u^2}}^{\sqrt{1-u^2}} \left[5 - 5u^2 - 5y^2 \right] dy du$$

$$= \int_{-1}^{1} \int_{-\sqrt{1-u^2}}^{\sqrt{1-u^2}} 5 \left[(1 - u^2 - y^2) \right] dy du$$

$$= 5 \int_{-1}^{1} \left[y - u^2 y - \frac{y^3}{3} \right]_{-\sqrt{1-u^2}}^{\sqrt{1-u^2}} du$$

$$+ \int_{-1}^{1} f(u) du = 28$$

$$= 5 \int_{-1}^{1} \left[(1 - u^2) y - \frac{y^3}{3} \right]_{-\sqrt{1-u^2}}^{\sqrt{1-u^2}} du$$

$$= 5 \int_{-1}^{1} \left[2(1 - u^2) (1 - u^2)^{\frac{1}{2}} - \frac{2}{3} (1 - u^2)^{\frac{3}{2}} \right] du$$

$$= 5 \int_{-1}^{1} \left[2 \cdot (1 - u^2)^{\frac{1}{2} + \frac{1}{2}} - \frac{2}{3} (1 - u^2)^{\frac{3}{2}} \right] du$$

$$= \frac{5 \times 2}{3} \int_{-1}^{+1} 2 (1-u^2)^{\frac{3}{2}} du$$

when limit is from 0 to 1 we should multiply by 2

$$= \frac{20}{3} \int_{-1}^{+1} (1-u^2)^{\frac{3}{2}} du = \frac{20 \times 2}{3} \int_0^1 (1-u^2)^{\frac{3}{2}} du$$

$(a^2 - u^2)$

which is in the form of ~~$(a^2 - u^2)$~~ , i.e. $(1^2 - u^2)$
 $\therefore a = 1$

$$\text{Put } u = a \sin \theta$$

$$\therefore u = \sin \theta$$

differentiating w.r.t. θ

$$\frac{du}{d\theta} = \frac{d(\sin \theta)}{d\theta}$$

$$\therefore du = \cos \theta d\theta$$

when,

$$u = 1$$

$$\text{when, } u = 0$$

$$1 = \sin \theta$$

$$0 = \sin \theta$$

$$\therefore \theta = \sin^{-1}(1) = \frac{\pi}{2}$$

$$\therefore \theta = \sin^{-1}(0) = 0$$

$$\therefore \text{Volume} = \int_0^{\frac{\pi}{2}} \frac{20 \times 2}{3} (1 - \sin^2 \theta)^{\frac{3}{2}} \cdot \cos \theta d\theta$$

$$= \frac{20 \times 2}{3} \int_0^{\frac{\pi}{2}} \cos^{\frac{3}{2}} \theta \cdot \cos \theta \cdot \cos \theta d\theta$$

$$= \frac{20 \times 2}{3} \int_0^{\frac{\pi}{2}} \cos^4 \theta d\theta$$

$$= \frac{20 \times 2}{3} \int_0^{\frac{\pi}{2}} (\cos^2 \theta)^2 d\theta$$

$$\begin{aligned}
 &= \frac{20 \times 2}{3} \int_0^{\frac{\pi}{2}} \left(\frac{1 + \cos 2\alpha}{2} \right)^2 d\alpha \quad \cos^2 \alpha = \frac{1 + \cos 2\alpha}{2} \\
 &= \frac{20 \times 2}{3} \int_0^{\frac{\pi}{2}} \frac{1}{4} (1 + 2\cos 2\alpha + \cos^2 2\alpha) d\alpha \quad \cos^2 2\alpha = \frac{1 + \cos 4\alpha}{2} \\
 &= \frac{20 \times 2}{3} \int_0^{\frac{\pi}{2}} \frac{1}{4} \left(1 + 2\cos 2\alpha + \frac{1 + \cos 4\alpha}{2} \right) d\alpha \\
 &= \frac{20 \times 2}{3} \int_0^{\frac{\pi}{2}} \frac{1}{4} \left(\frac{2 + 4\cos 2\alpha + 1 + \cos 4\alpha}{2} \right) d\alpha \\
 &= \frac{20 \times 2}{3} \times \frac{1}{8} \int_0^{\frac{\pi}{2}} (3 + 4\cos 2\alpha + \cos 4\alpha) d\alpha \\
 &= \frac{20 \times 2}{3 \times 8} \left[3[\alpha]_0^{\frac{\pi}{2}} + 4 \left[\frac{\sin 2\alpha}{2} \right]_0^{\frac{\pi}{2}} + \left[\frac{\sin 4\alpha}{4} \right]_0^{\frac{\pi}{2}} \right] \\
 &= \frac{20 \times 2}{3 \times 8} \left[3 \cdot \frac{\pi}{2} + 4 \left[\frac{\sin 2 \cdot \frac{\pi}{2} - \sin 0}{2} \right] + \left[\frac{\sin 4 \cdot \frac{\pi}{2} - \sin 0}{4} \right] \right] \\
 &= \frac{20 \times 2}{3 \times 8} \left[\frac{3\pi}{2} + 0 + 0 \right] \\
 &= \frac{20}{3 \times 8} \times \frac{3\pi}{2} \\
 &= \frac{10 \times 2}{8} \\
 &= \frac{5 \times 10 \pi \times 2}{8} \\
 \therefore V &= \frac{2 \times 5 \pi}{4} \text{ Cubic Unit} = \frac{5\pi}{2}
 \end{aligned}$$

Hence Required Volume is $\frac{5\pi}{2}$ Cubic unit.

* find the Volume bounded above by paraboloid $z = 8 - u^2 - y^2$ and $z = u^2 + y^2$.

→ Solution,

for Upper and lower limit of Z , Put $u=0, y=0$

$$Z = 8 - u^2 - y^2 = 8 - 0 - 0 = 8 \rightarrow \text{Upper limit}$$

$$Z = u^2 + y^2 = 0 + 0 = 0 \rightarrow \text{lower limit.}$$

for Upper and lower limit of y , solving eqⁿ $8 - u^2 - y^2$ and $u^2 + y^2$

$$\text{or, } 8 - u^2 - y^2 = u^2 + y^2$$

$$\text{or, } 2u^2 + 2y^2 = 8$$

$$\text{or, } u^2 + y^2 = 4 \text{ eq}^n \text{ of circle}$$

$$\text{or, } y^2 = 4 - u^2$$

$$\therefore y = \pm \sqrt{4 - u^2}$$

$$\therefore \text{lower limit of } y = -\sqrt{4 - u^2}$$

$$\text{Upper limit of } y = +\sqrt{4 - u^2}$$

for Upper & lower limit of Z ,


$u^2 + y^2 = (2)^2$ is the eqⁿ of circle

$$\text{so, } u^2 + y^2 = a^2$$

$$\therefore a = 2$$

$$\text{Upper limit} = 2$$

$$\text{lower limit} = -2$$



$$\therefore \text{Volume } (V) = \int_{u=-2}^{+2} \int_{y=-\sqrt{4-u^2}}^{\sqrt{4-u^2}} \int_{z=u^2+y^2}^{8-u^2-y^2} dz dy du$$

$$= \int_{-2}^{+2} \int_{-\sqrt{4-u^2}}^{\sqrt{4-u^2}} [z]_{u^2+y^2}^{8-u^2-y^2} dy du$$

$$= \int_{-2}^{+2} \int_{-\sqrt{4-u^2}}^{\sqrt{4-u^2}} [8-2u^2-2y^2] dy du$$

$$= \int_{-2}^{+2} \left[8y - 2u^2y - \frac{2y^3}{3} \right]_{-\sqrt{4-u^2}}^{\sqrt{4-u^2}} du$$

$$= \int_{-2}^{+2} \left[(8-2u^2)y - \frac{2y^3}{3} \right]_{-\sqrt{4-u^2}}^{\sqrt{4-u^2}} du \quad \text{if } \int_{-a}^{+a} f(u) du = 2 \int_0^a f(u) du$$

$$= \int_{-2}^{+2} \left[2(4-u^2)y - \frac{2y^3}{3} \right]_{-\sqrt{4-u^2}}^{\sqrt{4-u^2}} du$$

$$= \int_{-2}^{+2} \left[2 \times 2 (4-u^2) (\sqrt{4-u^2}) - \frac{2 \times 2}{3} (\sqrt{4-u^2})^3 \right] du$$

$$= 4 \int_{-2}^{+2} \left[(4-u^2)^{\frac{1}{2}} (4-u^2)^{\frac{1}{2}} - \frac{1}{3} (4-u^2)^{\frac{3}{2}} \right] du$$

$$= \frac{4}{3} \int_{-2}^{+2} \left[3(4-u^2)^{\frac{3}{2}} - (4-u^2)^{\frac{3}{2}} \right] du$$

$$= \frac{4 \times 2}{3} \int_{-2}^{+2} (4-u^2)^{\frac{3}{2}} du$$

$$= \frac{8}{3} \int_{-2}^2 (4-u^2)^{3/2} du$$

if we do \int_{-2}^2 to \int_0^2 we have to $\times 2$.

$$= \frac{8 \times 2}{3} \int_0^2 (4-u^2)^{3/2} du$$

which is in the form of $(a^2 - u^2)^{3/2}$ so, $a=2$
put $2u = 2\sin\theta = 2\sin\theta$

$$\frac{du}{d\theta} = 2\cos\theta$$

$$\therefore du = 2\cos\theta d\theta$$

$$\text{when } u=0$$

$$0 = 2\sin\theta$$

$$\sin\theta = 0$$

$$\therefore \theta = \sin^{-1}(0) = 0^\circ$$

$$\text{when } u=2$$

$$2 = 2\sin\theta$$

$$\sin\theta = 1$$

$$\therefore \theta = \sin^{-1}(1) = \frac{\pi}{2}$$

then,

$$\begin{aligned} \text{Volume (V)} &= \frac{16}{3} \int_0^{\pi/2} (4 - 4\sin^2\theta)^{3/2} \cdot 2\cos\theta d\theta \\ &= \frac{16}{3} \int_0^{\pi/2} (4)^{3/2} \cdot (1 - \sin^2\theta)^{3/2} \cdot 2\cos\theta d\theta \\ &= \frac{16}{3} \int_0^{\pi/2} 2^2 \cdot 2^{3/2} (\cos^2\theta)^{3/2} \cdot 2\cos\theta d\theta \\ &= \frac{16}{3} \times 8 \times 2 \int_0^{\pi/2} \cos^4\theta d\theta \\ &= \frac{16 \times 16}{3} \int_0^{\pi/2} (\cos^2\theta)^2 d\theta \\ &= \frac{16 \times 16}{3} \int_0^{\pi/2} \left(\frac{1 + \cos 2\theta}{2}\right)^2 d\theta \end{aligned}$$

$$\begin{aligned}
 &= \frac{16 \times 16}{3} \int_0^{\frac{\pi}{2}} \frac{1}{4} (1 + 2 \cos 2\alpha + \cos^2 2\alpha) d\alpha \\
 &= \frac{16 \times 16}{3} \int_0^{\frac{\pi}{2}} \frac{1}{4} (1 + 4 \cos 2\alpha + \frac{1}{2} (1 + \cos 4\alpha)) d\alpha \\
 &= \frac{16 \times 16}{3} \int_0^{\frac{\pi}{2}} \frac{1}{4} \left(\frac{2 + 2 \cos 2\alpha + 1 + \cos 4\alpha}{2} \right) d\alpha \\
 &= \frac{16 \times 16}{3} \times \frac{1}{8} \int_0^{\frac{\pi}{2}} (3 + 2 \cos 2\alpha + \cos 4\alpha) d\alpha \\
 &= \frac{32}{3} \left[3[\alpha]_0^{\frac{\pi}{2}} + 2 \left[\frac{\sin 2\alpha}{2} \right]_0^{\frac{\pi}{2}} + \left[\frac{\sin 4\alpha}{4} \right]_0^{\frac{\pi}{2}} \right] \\
 &= \frac{32}{3} \left[3 \times \frac{\pi}{2} + 2 \times 0 + 0 \right] \\
 &= \frac{32}{3} \times \frac{3\pi}{2} \\
 &= 16\pi \text{ cubic unit}
 \end{aligned}$$

Hence the required Volume is 16π (cubic unit).

~~Volume~~ volume nikalda jhukne phao.

$$\Rightarrow \int_{-a}^a x a \, dx \Rightarrow 2a$$

$$\Rightarrow \int_0^a (4 - x^2) \, dx = \text{yeolai put game } x = 2 \sin \theta$$

$$\Rightarrow \int_0^{\frac{\pi}{2}} (4 - 4 \sin^2 \theta) \, d\theta \text{ limit 0 to } \frac{\pi}{2} \text{ mul by } 2a$$

$$\Rightarrow \text{ani } (4 - 4 \sin^2 \theta) \text{ thau ma common } \int_0^{\frac{\pi}{2}} (1 - \sin^2 \theta) \, d\theta = 8 \text{ hundha}$$

In 2 dimension

classmate

Date _____

Page _____

Mass and First Moment formula

Mass :-

$$M = \iint_R \delta(u, y) dA$$

$\therefore \delta(u, y)$ denotes density at (x, y)

First Moments :-

$$Mu = \iint_R y \delta(u, y) dA$$


$$My = \iint_R u \delta(u, y) dA$$

$$\text{Center of mass (centroid)} = \bar{u} = \frac{My}{M}$$

$$\bar{y} = \frac{Mu}{M}$$

- * A thin plate covers the triangular Region bounded by the x-axis and the lines $u=1$, $y=2u$ in the first quadrant. The plate density at the point is $\delta(u, y) = 6u + 6y + 6$. Find the plate's mass, first moments and centroid (Centre of mass) about the co-ordinate axes.

From e.g. $y = 2u$
when $u=0, y=0$
 $u=1, y=2$



$$u=1 \quad y=2u$$

$$= \int_{u=0}^{u=1} \int_{y=0}^{y=2u} s(u, y) dy du$$

$$u=1 \quad y=2u$$

$$\therefore \text{we know mass} = \int_{u=0}^{u=1} \int_{y=0}^{y=2u} (6u+6y+6) dy du$$

$$u=1 \quad y=2u$$

$$= \int_{u=0}^{u=1} \int_{y=0}^{y=2u} 6(u+y+1) dy du$$

$$= 6 \int_0^1 \left[u^2 + \frac{y^2}{2} + y \right]_{y=0}^{y=2u} du$$

$$= 6 \int_0^1 \left[u(2u) + \frac{(2u)^2}{2} + 2u \right] du$$

$$= 6 \int_0^1 \left[2u^2 + \frac{2u^2}{2} + 2u \right] du$$

$$= 6 \int_0^1 \left[4u^2 + 2u \right] du$$

$$= 6 \times 2 \int_0^1 \left[2u^2 + u \right] du$$

$$= 12 \left\{ \left[\frac{2u^3}{3} \right]_0^1 + \left[\frac{u^2}{2} \right]_0^1 \right\}$$

$$= 12 \left\{ \frac{2}{3} + \frac{1}{2} \right\}$$

$$= 12 \times \left(\frac{4+3}{6} \right)$$

$$= 12 \times \frac{7}{6} = \frac{26}{3} \times 7 = 14$$

$$\therefore \text{mass (M)} = 14$$

first moments,

$$M_u = \iint_R y s(u, y) dA$$

$$u=1 \quad y=2u$$

$$= \int_{u=0}^1 \int_{y=0}^{2u} y (6u + 6y + 6) dy du$$

$$u=0 \quad y=0$$

$$= \int_{u=0}^1 \int_{y=0}^{2u} (6uy + 6y^2 + 6y) dy du$$

$$u=1 \quad y=0$$

$$= \int_{u=0}^1 \left[\left[6uy^2 \right]_0^{2u} + \left[26y^3 \right]_0^{2u} + \left[6y^2 \right]_0^{2u} \right] du$$

$$= \int_{u=0}^1 \left[3u \cdot (2u)^2 + 2(2u)^3 + 3(2u)^2 \right] du$$

$$= \int_{u=0}^1 \left[12u^3 + 16u^3 + 12u^2 \right] du$$

$$= \int_{u=0}^1 \left[28u^3 + 12u^2 \right] du$$

$$= 28 \left[\frac{u^4}{4} \right]_0^1 + 12 \left[\frac{u^3}{3} \right]_0^1$$

$$= 28 \times \frac{1}{4} + 12 \times \frac{1}{3}$$

$$= 7 + 4 = 11 \quad \therefore M_u = 11$$

$$\begin{aligned}
 M_y &= \iint_R u \delta(u, y) dA \\
 &= \int_{u=0}^1 \int_{y=0}^{2u} u(6u+6y+6) dy du \\
 &= \int_0^1 \int_0^{2u} (6u^2+6uy+6u) dy du \\
 &= \int_0^1 \left[6u^2y + \frac{3}{2}6uy^2 + 6uy \right]_0^{2u} du \\
 &= \int_0^1 \left[6u^2(2u) + \frac{3}{2}6u(2u)^2 + 6u(2u) \right] du \\
 &= \int_0^1 \left[12u^3 + 12u^3 + 12u^2 \right] du \\
 &= \int_0^1 12 \left[2u^3 + u^2 \right] du \\
 &= 12 \left\{ \left[\frac{2u^4}{4} \right]_0^1 + \left[\frac{u^3}{3} \right]_0^1 \right\} \\
 &= 12 \cdot \left(\frac{1}{2} + \frac{1}{3} \right) \\
 &= 12 \times \left(\frac{3+2}{6} \right) \quad \therefore M_y = 10 \\
 &= 12 \times \frac{5}{6} = 10
 \end{aligned}$$

$$\therefore \text{Centre of mass (centroid)} = \bar{u} = \frac{My}{M}$$

$$\bar{u} = \frac{10}{14} = \frac{5}{7}$$

$$\bar{y} = \frac{Mu}{M} = \frac{11}{14}$$

$$\therefore \text{Centroid} = \left(\frac{5}{7}, \frac{11}{14} \right)$$

Second Moment (Moment of Inertia) :-

1. About X-axis :-

$$I_u = \iint_R y^2 s(u, y) dA$$

2. About Y-axis :-

$$I_y = \iint_R u^2 s(u, y) dA$$

3. About Origin :-

$$I_o = I_u + I_y$$

Radii of Gyration :-

1. About X-axis :-

$$R_u = \sqrt{\frac{I_u}{M}}$$

2. About Y-axis :-

$$R_y = \sqrt{\frac{I_y}{M}}$$

3. About Origin :-

$$R_o = \sqrt{\frac{I_o}{M}}$$

* Find the moment of inertia [Second moment] and radii of Gyration from eqⁿ $\delta(u, y) = 6u + 6y + 6$ about coordinate axes and origin. $x=1, y=2u$ in 1st q

→ Solution,

$$\delta(u, y) = 6u + 6y + 6$$

Moment of Inertia (Second Moment)

About x-axis,

$$I_x = \iint y^2 \delta(u, y) dA$$

$$= \int_0^1 \int_0^{2u} y^2 (6u + 6y + 6) dy du$$

$$= \int_0^1 \int_0^{2u} (6uy^2 + 6y^3 + 6y^2) dy du$$

$$= \int_0^1 \left[\frac{2uy^3}{3} \Big|_0^{2u} + 6 \left[\frac{y^4}{4} \Big|_0^{2u} + 2 \left[\frac{y^3}{3} \Big|_0^{2u} \right] \right] du$$


$$= \int_0^1 \left[2uy^3 + \frac{3}{2} y^4 + 2y^3 \right]_0^{2u} du$$

$$= \int_0^1 \left[2u(2u)^3 + \frac{3}{2} (2u)^4 + 2(2u)^3 \right] du$$

$$= \int_0^1 \left[16u^4 + 3 \times \frac{16}{2} u^4 + 16u^3 \right] du$$

$$= \int_0^1 16 \left[\cancel{u^4} + \frac{3}{2} u^4 + u^3 \right] du$$

$$= \int_0^1 16 \left[\frac{2u^4 + 3u^4 + 2u^3}{2} \right] du$$



$$= \frac{16}{2} \int_0^1 (5u^4 + 2u^3) du$$

$$= \frac{16}{2} \left\{ 5 \left[\frac{u^5}{5} \right]_0^1 + 2 \left[\frac{u^4}{4} \right]_0^1 \right\}$$

$$= \frac{16}{2} \left\{ \frac{5}{5} + \frac{2}{4} \right\}$$

$$= \frac{16}{2} \left(\frac{3}{2} \right)$$

$$= 8 \times \frac{3}{2} = 4 \times 3 = 12$$

About :-

For Y-axis :-

$$I_y = \iint_R \delta(u, y) dA u^2$$

$$a. I_y = \int_0^1 \int_0^{2u} u^2 (6u + 6y + 6) dy du$$

$$= \int_0^1 \int_0^{2u} (6u^3 + 6u^2y + 6u^2) dy du$$

$$= \int_0^1 \left[6u^3y + \frac{6u^2y^2}{2} + 6u^2y \right]_0^{2u} du$$

$$= \int_0^1 [6u^3 \cdot (2u) + 3u^2 (2u)^2 + 6u^2 (2u)] du$$

$$= \int_0^1 (12u^4 + 12u^4 + 12u^3) du$$

$$= \int_0^1 (12u^3 + 12u^4) du$$

$$\begin{aligned}
 &= 12 \int_0^1 (2u^4 + u^3) du \\
 &= 12 \int_0^1 (2u^4 + u^3) du = 12 \left[\frac{2u^5}{5} + \frac{u^4}{4} \right]_0^1 \\
 &= 12 \left(\frac{2}{5} + \frac{1}{4} \right) \\
 &= 12 \times \frac{13}{20} \\
 &= \frac{39}{5}
 \end{aligned}$$

About origin,

$$I_0 = I_x + I_y = 12 + \frac{39}{5} = \frac{99}{5}$$

∴ Radii of Gyration,

$$\text{About } x\text{-axis } (R_x) = \sqrt{\frac{I_x}{M}} = \sqrt{\frac{12}{14}} = \sqrt{\frac{6}{7}}$$

$$\therefore R_x = 0.925$$

$$\text{About } y\text{-axis } (R_y) = \sqrt{\frac{I_y}{M}} = \sqrt{\frac{39}{5} \times \frac{1}{14}} = \sqrt{\frac{39}{70}}$$

$$\therefore R_y = 0.746$$

$$\text{About Origin } (R_0) = \sqrt{\frac{I_0}{M}} = \sqrt{\frac{99}{5} \times \frac{1}{14}} = \sqrt{\frac{99}{70}}$$

$$\therefore R_0 = 1.189$$

* Find the centroid of the region in the first quadrant that is bounded above by the line $y = u$ and below by the parabola $y = u^2$.

$$\rightarrow y = u \quad y = u^2$$

$$u^2 = u$$

$$\text{or, } u^2 - u = 0 \quad u(u-1) \quad \therefore u = 0, u = 1$$

$$y = 0 \quad y = 1$$

when $u=0 \ y=0$ $u=1 \ y=1$ $u=2 \ y=4$ \Rightarrow First Mass is given by

$$\text{Mass} = \iint_R \delta(u, y) dA$$

$$= \int_{u=0}^{u=1} \int_{y=u^2}^{y=u} 1 dy du$$

$$= \int_0^1 [y]_{u^2}^u du = \int_0^1 [u - u^2] du = \int_0^1 u du - \int_0^1 u^2 du$$

$$= \left[\frac{u^2}{2} \right]_0^1 - \left[\frac{u^3}{3} \right]_0^1 = \frac{1}{2} - \frac{1}{3} = \frac{3-2}{6} = \frac{1}{6}$$

$$\therefore \text{Mass (M)} = \frac{1}{6}$$

Now,


first moment,

$$Mx = \iint_R y \delta(u, y) dA$$

$$= \int_0^1 \int_{u^2}^u y dy du = \int_0^1 \int_{u^2}^u y dy du$$

$$= \int_0^1 \left[\frac{y^2}{2} \right]_{u^2}^u du = \frac{1}{2} \int_0^1 [(u)^2 - (u^2)^2] du$$

$$= \frac{1}{2} \int_0^1 (u^2 - u^4) du$$



$$= \frac{1}{2} \left[\frac{u^3}{3} - \frac{u^5}{5} \right]_0^1 = \frac{1}{2} \left(\frac{1}{3} - \frac{1}{5} \right)$$

$$= \frac{1}{2} \left(\frac{2}{15} \right) = \frac{1}{15}$$

$$\therefore M_u = \frac{1}{15}$$

$$M_y = \int_0^1 \int_{u^2}^u u \, dy \, du$$

$$= \int_0^1 [uy]_{u^2}^u \, du = \int_0^1 [u(u) - u(u^2)] \, du$$

$$= \int_0^1 [u^2 - u^3] \, du = \int_0^1 u^2 \, du - \int_0^1 u^3 \, du$$

$$= \left[\frac{u^3}{3} \right]_0^1 - \left[\frac{u^4}{4} \right]_0^1 = \frac{1}{3} - \frac{1}{4} = \frac{1}{12}$$

$$\therefore M_y = \frac{1}{12}$$

Now,

we have, Centroid is given by Ans. for moment of inertia,

$$\bar{u} = \frac{M_y}{M} = \frac{1}{12} \times \frac{6}{1} = \frac{1}{2}$$

$$I_u = \frac{1}{28}, I_y = \frac{1}{20}$$

$$\bar{y} = \frac{M_u}{M} = \frac{1}{15} \times \frac{6}{1} = \frac{6}{15} = \frac{2}{5} \quad I_o = \frac{3}{35}$$

X-axis. R_u = $\frac{I_u}{M} = 0.035$

$$\therefore \text{Centroid} = \left(\frac{1}{2}, \frac{2}{5} \right) \#$$

Y-axis. R_y = $\frac{I_y}{M} = 0.05$

Origin - R_o: $\frac{I_o}{M} = 0.50$

"Formulae in three dimension"

1. Masses And Moments in three dimension

i) Mass (M) = $\iiint_R \delta \, dv$ $\therefore \delta$ = density at x, y, z

2. First Moments in co-ordinate planes.

i) $M_{xy} = \iiint_R z \delta \, dv$ - In about xy -plane
 $\therefore \delta \, dv = dx \cdot dy \cdot dz \delta$

ii) $M_{yz} = \iiint_R x \delta \, dv$ - About yz -plane

iii) $M_{xz} = \iiint_R y \delta \, dv$ - About xz -plane

3. centre of mass (centroid) :

i) $\bar{x} = \frac{M_{yz}}{M}$

ii) $\bar{y} = \frac{M_{xz}}{M}$

iii) $\bar{z} = \frac{M_{xy}}{M}$

4. Motion of Inertia [Second Moment] :

i) $I_x = \iiint_R (y^2 + z^2) \delta \, dv$

ii) $I_y = \iiint_R (x^2 + z^2) \delta \, dv$


iii) $I_z = \iiint_R (x^2 + y^2) \delta \, dv$

* Find I_x, I_y, I_z for rectangular solid of constant density δ as in fig. (Put $M = abc\delta$)

→ Solution, $y = b/2$ $u = a/2$

$$z = +\frac{c}{2} \quad u = \frac{a}{2}$$

$$I_x = \int_{z=-\frac{c}{2}}^{z=\frac{c}{2}} \int_{y=-\frac{b}{2}}^{y=\frac{b}{2}} \int_{u=-\frac{a}{2}}^{u=\frac{a}{2}} \delta(y^2 + z^2) dy dz$$



when we go from lower limit 0 to $+\frac{c}{2}$ we should multiply by 2.

Similarly in 0 to $b/2$ and in 0 to $a/2$

then,

$$I_x = 2 \times 2 \times 2 \int_{z=0}^{z=\frac{c}{2}} \int_{y=0}^{y=\frac{b}{2}} \int_{u=0}^{u=\frac{a}{2}} \delta(y^2 + z^2) dy dz$$

$$= 8\delta \int_{z=0}^{z=\frac{c}{2}} \int_{y=0}^{y=\frac{b}{2}} \int_{u=0}^{u=\frac{a}{2}} [uy^2 + uz^2]_{0}^{\frac{a}{2}} dy dz$$

$$= 8\delta \int_{z=0}^{z=\frac{c}{2}} \int_{y=0}^{y=\frac{b}{2}} \int_{u=0}^{u=\frac{a}{2}} [\frac{a}{2}y^2 + \frac{a}{2}z^2] dy dz$$

$$= 8\delta \cdot \frac{a}{2} \int_{z=0}^{z=\frac{c}{2}} \int_{y=0}^{y=\frac{b}{2}} [y^2 + z^2] dy dz$$

$$= 4a\delta \int_{z=0}^{z=\frac{c}{2}} \left[\frac{y^3}{3} + yz^2 \right]_{0}^{\frac{b}{2}} dz$$

$$= 4a\delta \int_{z=0}^{z=\frac{c}{2}} \left[\frac{b^3}{8 \times 3} + \frac{b}{2}z^2 \right] dz$$

$$= 4ab\delta \frac{b}{2} \int_0^{\frac{b}{2}} \left[\frac{b^2}{12} + z^2 \right] dz$$

$$= 2ab\delta \left[\frac{b^2}{12} z + \frac{z^3}{3} \right]_0^{\frac{b}{2}}$$

$$= 2ab\delta \left[\frac{cb^2}{24} + \frac{c^3}{8 \times 3} \right]$$

$$= 2ab\delta \left[\frac{cb^2}{24} + \frac{c^3}{24} \right]$$

$$= \frac{2abc\delta}{24} (b^2 + c^2)$$

$$= \frac{2abc\delta}{12} (b^2 + c^2)$$

$$\therefore I_u = \frac{M}{12} (b^2 + c^2)$$

Now,

$$I_y = \int_0^{\frac{a}{2}} \int_0^{\frac{b}{2}} \int_0^{\frac{a}{2}} 8\delta(u^2 + z^2) du dy dz$$

$$= 8\delta \int_0^{\frac{a}{2}} \int_0^{\frac{b}{2}} \left[\frac{u^3}{3} + uz^2 \right]_0^{\frac{a}{2}} dy dz$$

$$= 8\delta \int_0^{\frac{a}{2}} \int_0^{\frac{b}{2}} \left[\frac{a^3}{24 \times 3} + \frac{3}{2} z^2 \right] dy dz$$

$$= 8\delta \cdot \frac{a}{2} \int_0^{\frac{a}{2}} \int_0^{\frac{b}{2}} \left[\frac{a^2}{12} + z^2 \right] dy dz$$

$$= 4a\delta \int_0^{\frac{a}{2}} \left[\frac{a^2 y}{12} + z^2 y \right]_0^{\frac{b}{2}} dz$$

$$= 4\pi \delta \int_0^{y_2} \frac{b}{2} \left(\frac{a^2}{12} + z^2 \right) dz$$

$$= 4\pi \delta \frac{b}{2} \left[\frac{a^2 z}{12} + \frac{z^3}{3} \right]_0^{y_2}$$

$$= 2\pi b \delta \left[\frac{a^2 c}{12 \times 2} + \frac{c^3}{24} \right]$$

$$= 2\pi b c \frac{\delta}{24} (a^2 + c^2)$$

$$\therefore I_y = \frac{M}{12} (a^2 + c^2)$$

(Similarly, $\int_0^{y_2} \int_0^{b/2} \int_0^{a/2}$

$$I_z = 8\delta \int_0^{y_2} \int_0^{b/2} \int_0^{a/2} (u^2 + y^2) du dy dz$$

$$= \int_0^{y_2} \int_0^{b/2} 8\delta \left[\frac{u^3}{3} + uy^2 \right]_0^{a/2} dy dz$$

$$= \int_0^{y_2} \int_0^{b/2} 8\delta \left[\frac{a^3}{24} + \frac{a}{2} y^2 \right] dy dz$$

$$= 8\delta \frac{a}{2} \int_0^{y_2} \int_0^{b/2} \left[\frac{a^2}{12} + y^2 \right] dy dz$$

$$= 4\pi \delta \int_0^{y_2} \left[\frac{a^2 y}{12} + \frac{y^3}{3} \right]_0^{b/2} dy$$

$$= 4\pi \delta \int_0^{y_2} \left[\frac{a^2 b}{24} + \frac{b^3}{24} \right] dy$$

$$= \frac{4abc}{24} \left[a^2 z + b^2 z \right]_0^{\frac{c}{2}}$$

$$= \frac{abc}{6} \left[a^2 \frac{c}{2} + b^2 \frac{c}{2} \right]$$

$$= \frac{abc}{6 \times 2} (a^2 + b^2)$$

$$\therefore I_z = \frac{M}{12} (a^2 + b^2) \cancel{\#}$$

$$\therefore I_y = \frac{M}{12} (a^2 + c^2) \cancel{\#}$$

$$\therefore I_x = \frac{M}{12} (b^2 + c^2) \cancel{\#}$$

Double Integral in Polar form :-

Note :-

$$dudv = dydu = r dr d\theta$$

$$u^2 + y^2 = r^2$$

$$u^2 + y^2 = r^2 \text{ eqn of circle}$$

$$* \int_{u=-1}^1 \int_{y=\sqrt{1-u^2}}^{y=-\sqrt{1-u^2}} dy du.$$

$$u = -1 \quad y = -\sqrt{1-u^2}$$

$$= 2 \times 2 \int_0^1 \int_0^{\sqrt{1-u^2}} dy du$$

$$= 4 \int_0^1 \int_0^{\sqrt{1-u^2}} dy du$$

In Polar, when

$$\begin{aligned} \text{always } & \rightarrow y = 0 \\ & \rightarrow r = 0 \end{aligned}$$

$$\begin{aligned} y &= \sqrt{1-u^2} \\ u^2 + y^2 &= 1 \\ \therefore r &= 1 \end{aligned}$$

$$u^2 + y^2 = 1^2$$

$$\begin{aligned} \text{when } y &= \sqrt{1-r^2} \\ \therefore r^2 &= 1-r^2 \\ \therefore r &= 1 \end{aligned}$$

$$u=0$$

$$\alpha = 0$$

Now,

$$\Rightarrow 4 \int_{\alpha_1=0}^{\alpha_2=\pi/2} \int_{r=0}^{r=1} r dr d\alpha$$

$$= 4 \int_0^{\pi/2} \left[\frac{r^2}{2} \right]_0^1 d\alpha = 4 \int_0^{\pi/2} \left[\frac{1}{2} \right] d\alpha$$

$$= 4 \times \frac{1}{2} \left[\alpha \right]_0^{\pi/2}$$

$$= 2 \cdot \frac{\pi}{2}$$

$$= \pi$$

\therefore Area of circle is π sq. unit.

Double integral in Polar form:

$$u = y \quad y = \sqrt{a^2 - u^2}$$


$$\text{Q. } \int_{u=-a}^a \int_{y=-\sqrt{a^2-u^2}}^{\sqrt{a^2-u^2}} dy du$$

$$y = \sqrt{a^2 - u^2}$$

when we go from 0 to a , & 0 to $\sqrt{a^2 - u^2}$ we should do 2×2 .

$$= \int_{-a}^a \int_{-\sqrt{a^2-u^2}}^{\sqrt{a^2-u^2}} dy du$$

$$= 2 \times \int_0^a \times 2 \int_0^{\sqrt{a^2-u^2}} dy du$$



$$y = \sqrt{a^2 - u^2}$$

$$= 4 \int_0^a \int_0^{\sqrt{a^2 - u^2}} dy du$$

to change in polar,
when $y=0$

$$r=0$$

$$\begin{matrix} \text{now} \\ \theta=0 \end{matrix}$$

$$y = \sqrt{a^2 - u^2}$$

then,

$$y^2 = a^2 - u^2$$

$$\therefore u^2 + y^2 = a^2$$

$$\therefore r^2 = a^2 \quad \therefore r = a$$

when,

$$u=0$$

$$u=\pi$$

$$\theta_1 = 0^\circ$$

$$\theta_2 = \frac{\pi}{2}$$

Now,

$$\text{Q: } \pi/2 \quad r=a$$

$$= 4 \int_{\theta=0}^{\pi/2} \int_{r=0}^a r dr d\theta$$

$$[dy du = du dy = r dr d\theta]$$

$$= 4 \int_0^{\pi/2} \left[\frac{r^2}{2} \right]_0^a d\theta = 4 \int_0^{\pi/2} \frac{a^2}{2} d\theta = 2a^2 \int_0^{\pi/2} d\theta$$

$$= 2a^2 \left[\theta \right]_0^{\pi/2} = 2a^2 \times \frac{\pi}{2} = \pi a^2$$

$$\frac{1}{\sqrt{1-u^2}}$$

$$\int_0^1 \int_0^{\sqrt{1-u^2}} (u^2 + y^2) dy du$$

when,

→ solution,

when,

$$y=0$$

$$r=0$$

$$y = \sqrt{1-u^2}$$

then, sq.

$$y^2 = 1 - u^2$$

$$u^2 + y^2 = 1$$

$$\therefore r^2 = 1 \quad \therefore r = 1$$

when, $\begin{cases} u=0 \\ \alpha_1=0^\circ \end{cases}$

when, $\begin{cases} u=1 \\ \alpha_2=\pi/2 \end{cases}$

Now, $\begin{cases} \alpha_2 = \pi/2 \\ \alpha_1 = 0^\circ \end{cases} \quad r=1$

$$\int_{r=0}^{r=1} (u^2 + y^2) r dr d\alpha$$

$$[\because u^2 + y^2 = r^2]$$

$$= \int_0^{\pi/2} \int_0^1 r^2 \cdot r dr d\alpha$$

$$= \int_0^{\pi/2} \left[\frac{r^4}{4} \right]_0^1 d\alpha = \int_0^{\pi/2} \frac{1}{4} d\alpha = \frac{1}{4} [\alpha]_0^{\pi/2}$$

$$= \frac{1}{4} \times \frac{\pi}{2} = \frac{\pi}{8} \neq .$$

$$\text{Q.} \int_{-1}^1 \int_{-\sqrt{1-u^2}}^{\sqrt{1-u^2}} \frac{1}{(1+u^2+y^2)^2} dy du$$

$$\Rightarrow 2 \int_0^1 2 \int_{-\sqrt{1-u^2}}^{\sqrt{1-u^2}} \frac{1}{(1+u^2)^2} dy du$$

$$= 4 \int_0^1 \int_0^{\sqrt{1-u^2}} \frac{1}{(1+u^2)^2} dy du$$

when,

$$\begin{cases} y=0 \\ r=0 \end{cases}$$

$$\begin{cases} y=\sqrt{1-u^2} \\ r=1 \end{cases}$$

Rough

$$y = \int \frac{1}{(1+u^2)^2} u du$$

let

$$y = 1+u^2$$

$$\frac{dy}{du} = 2u$$

$$\therefore \frac{dy}{2} = u du$$

then,

$$y = \int \frac{1}{y^2} \frac{dy}{2}$$

$$= \frac{1}{2} \frac{y^{-2+1}}{-2+1}$$

$$= -\frac{1}{2} y$$

$$= -\frac{1}{2} (1+u^2)$$

when,

$$u=0 \quad u=1$$

$$\alpha_1=0 \quad \alpha_2=\frac{\pi}{2}$$

$$\therefore \int \frac{1}{(1+u^2)^2} u du$$

$$= -\frac{1}{2}(1+u^2)$$

$$= \int_0^{\frac{\pi}{2}} \int_0^1 \frac{1}{(1+r^2)^2} r dr d\alpha$$

$$\leftarrow r dr =$$

$$= \int_0^{\frac{\pi}{2}} \left[-\frac{1}{2}(1+r^2) \right]_0^1 d\alpha = \int_0^{\frac{\pi}{2}} \left[-\frac{1}{2} + \frac{1}{2} \right] d\alpha$$

$$= \int_0^{\frac{\pi}{2}} \left[\frac{1}{2} - \frac{1}{4} \right] d\alpha = \int_0^{\frac{\pi}{2}} \left(\frac{1}{4} \right) d\alpha = \frac{1}{4} \left[u \right]_0^{\frac{\pi}{2}}$$

$$= \frac{\pi}{2} \#.$$

$$\text{Q. } \int_{-1}^1 \int_0^y dy du$$

when,

$$y=0$$

$$\text{when, } y = \sqrt{1-u^2}$$

$$u^2 + y^2 = 1$$

$$= 2 \int_0^1 \int_0^y dy du$$

when $u=0$

$$u=1$$

$$\alpha=0$$

$$\alpha=\frac{\pi}{2}$$

$$= 2 \int_0^{\frac{\pi}{2}} \int_0^1 r dr d\alpha$$

$$= 2 \int_0^{\frac{\pi}{2}} \left[\frac{r^2}{2} \right]_0^1 d\alpha = 2 \int_0^{\frac{\pi}{2}} \frac{1}{2} d\alpha = 2 \times \frac{1}{2} \left[\alpha \right]_0^{\frac{\pi}{2}}$$

$$= \frac{\pi}{2} - 0 = \frac{\pi}{2} \#.$$

Substitutions in Multiple Integrals

Definition of Jacobian:-

- The Jacobian determinant or Jacobian of the co-ordinate transformation $u = g(u, v)$, $y = h(u, v)$ is,

$$J(u, v) = \begin{vmatrix} \frac{\partial u}{\partial u} & \frac{\partial u}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial u}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial u}{\partial v}$$

The Jacobian is denoted by,

$$J(u, v) = \frac{\partial(u, y)}{\partial(u, v)}$$

$$u = g(u, v)$$

$$y = h(u, v)$$

(u, v)

transformation

(u, y)

$$\iint_R f(u, y) dA \xrightarrow{\text{transformation}} \iint_R h(u, v) |J(u, v)| dA$$

* Jacobian determinant in 3-D.

$$u = g(u, v, w)$$

$$y = h(u, v, w)$$

$$z = k(u, v, w)$$

$$J(u, v, w) = \begin{vmatrix} \frac{\partial u}{\partial u} & \frac{\partial u}{\partial v} & \frac{\partial u}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}$$

$$\iiint_R f(u, y, z) du dy dz \xrightarrow{\text{transformation}} \iiint_R J(u, v, w) H(u, v, w) du dv dw$$

* Find the Jacobian of following + determinant.

a) $u = 4v, y = \frac{u}{v}$

→ Solution,

$$\frac{\partial u}{\partial u} = \frac{\partial 4v}{\partial u} = v$$

$$\frac{\partial u}{\partial v} = \frac{\partial 4v}{\partial v} = 4$$

$$\frac{\partial y}{\partial u} = \frac{\partial \frac{u}{v}}{\partial u} = \frac{1}{v}$$

$$\frac{\partial y}{\partial v} = \frac{\partial \frac{u}{v}}{\partial v} = u \frac{\partial v^{-1}}{\partial v}$$

$$= u \cdot \frac{1}{v^2} = \frac{u}{v^2}$$

$$\therefore J(u, v) = \begin{vmatrix} \frac{\partial u}{\partial u} & \frac{\partial u}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} v & 4 \\ \frac{1}{v} & -\frac{u}{v^2} \end{vmatrix}$$

$$= v \cdot \left(-\frac{u}{v^2}\right) - \frac{4}{v} = J(u, v) = \frac{v(-\frac{u}{v^2}) - \frac{4}{v}}{\frac{\partial y}{\partial u}} = \frac{-\frac{u}{v} - \frac{4}{v}}{\frac{\partial y}{\partial u}}$$

$$= -\frac{u+4}{v}$$

b) $u = \frac{u+v}{3}, y = \frac{v-2u}{3}$

→ Solution,

$$\frac{\partial u}{\partial u} = \frac{\partial \frac{u+v}{3}}{\partial u} = \frac{1}{3} \frac{\partial u+v}{\partial u} = \frac{1}{3}$$

$$\frac{\delta u}{\delta v} = \frac{\delta (4+\frac{v}{3})}{\delta v} = \frac{1}{3} \frac{\delta v}{\delta v} = \frac{1}{3}$$

$$\frac{\delta y}{\delta u} = \frac{\delta (v-2u)}{\delta u} = \frac{1}{3} \frac{\delta (2u)}{\delta u} = \frac{2}{3}$$

$$\frac{\delta y}{\delta v} = \frac{\delta (v-2u)}{\delta v} = \frac{1}{3} \frac{\delta v}{\delta v} = \frac{1}{3}$$

Now,

$$J(u, v) = \begin{vmatrix} \frac{1}{3} & \frac{1}{3} \\ -\frac{2}{3} & \frac{1}{3} \end{vmatrix} = \frac{1}{9} + \frac{2}{9} = \frac{3}{9} = \frac{1}{3} \#$$

c) $u = 4\cos v \quad y = 4\sin v$

→ Solution,

$$\frac{\delta u}{\delta u} = \frac{\delta u \cos v}{\delta u} = \cos v$$

$$\frac{\delta u}{\delta v} = \frac{\delta u \cos v}{\delta v} = 4(-\sin v) = -4\sin v$$

$$\frac{\delta y}{\delta u} = \frac{\delta y \sin v}{\delta u} = \sin v$$

$$\frac{\delta y}{\delta v} = \frac{\delta y \sin v}{\delta v} = 4\cos v$$

$$\therefore J(u, v) = \begin{vmatrix} \cos v & -4\sin v \\ \sin v & 4\cos v \end{vmatrix} = 4\cos^2 v + 4\sin^2 v = 4 \#$$

d) $u = 4\sin v \quad y = 4\cos v$

→ Solution,

$$\frac{\delta u}{\delta u} = \frac{\delta u \sin v}{\delta u} = \sin v$$

$$\frac{su}{sv} = \frac{su \sin v}{sv} = 4 \cos v$$

$$\frac{sy}{su} = \frac{su \cos v}{su} = \cos v$$

$$\frac{sy}{sv} = \frac{su \cos v}{sv} = -4 \sin v$$

$$\therefore J(4, v) = \begin{vmatrix} \sin v & 4 \cos v \\ \cos v & -4 \sin v \end{vmatrix} = -4 \sin^2 v - 4 \cos^2 v \\ = -4 (\sin^2 v + \cos^2 v) \\ = -4 \neq 0.$$

* Evaluate the double integral $\int_0^4 \int_0^{\frac{y}{2}} \frac{2u-y}{2} du dy$

by applying transformation,

$u = \frac{2u-y}{2}$, $v = \frac{y}{2}$ and integrating over an appropriate region in the uv -plane.

→ Solution:-

$$u = \frac{2u-y}{2}$$

$$v = \frac{y}{2}$$

$$\therefore 2u = 2u - y$$

$$\therefore y = 2v$$

$$\therefore 2u = 2u + y \quad [\because y = 2v]$$

$$\therefore u = \frac{2u+2v}{2}$$

$$\therefore u = \frac{2(u+v)}{2}$$

$$\therefore u = (u+v) \quad v = \frac{y}{2}$$

$$y = 2v$$

$$u = u + v$$

$$y = 2v$$

Now,

$$\frac{\delta u}{\delta u} = \frac{\delta(u+v)}{\delta u} = 1$$

$$\frac{\delta u}{\delta v} = \frac{\delta(u+v)}{\delta v} = 1$$

$$\frac{\delta y}{\delta u} = \frac{\delta 2v}{\delta u} = 0$$

$$\frac{\delta y}{\delta v} = \frac{\delta 2v}{\delta v} = 2$$

$$\therefore J(u, v) = \begin{vmatrix} 1 & 1 \\ 0 & 2 \end{vmatrix} = 1 \cdot 0 - 0 \cdot 1 = 2$$

Now,

we have to change limit from table.

$$(u = u + v) \quad y = 2v$$

my eqⁿ

$$u = \frac{y}{2} + 1$$

corresp. uv eqⁿ

$$\text{or, } u + v = \frac{2v}{2} + 1$$

simplified uv

$$\therefore u = v - v + 1 = 1$$

$$u = \frac{y}{2}$$

$$\text{or, } u + v = \frac{2v}{2}$$

$$\therefore u = v - v = 0$$

$$y = 4$$

$$2v = 4$$

$$\therefore v = \frac{4}{2} = 2$$

$$y = 0$$

$$2v = 0$$

$$\therefore v = 0$$

$$= \int_{y=0}^{y=4} \int_{u=\frac{y}{2}+1}^{u=\frac{y}{2}+1} \frac{2u-y}{2} du dy$$

$$= \int_{v=0}^{v=2} \int_{u=1}^{u=1} 4 |J(u, v)| du dv$$

$$= \int_{v=0}^{v=2} \int_{u=0}^{u=1} u \times 2 \, du \, dv$$

$$= 2 \int_{v=0}^{v=2} \left[\frac{u^2}{2} \right]_0^1 \, dv$$

$$= \frac{2}{2} \int_0^2 1 \, dv$$

$$= [v]_0^2$$

$$= 2$$

Substitution in Multiple Integrals

Defⁿ of Jacobian:-The Jacobian determinant or Jacobian of the co-ordinate transformation $u = g(u, v)$, $y = h(u, v)$ is,

$$J(u, v) = \begin{vmatrix} \frac{\partial u}{\partial u} & \frac{\partial u}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial u}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial y}{\partial u} \frac{\partial u}{\partial v}$$

* Evaluate:-

$$\int \int \int_{\substack{z=0 \\ y=0 \\ u=\frac{1}{2}}}^{\substack{z=3 \\ y=4 \\ u=\frac{8}{2}+1}} \left(\frac{2u-y}{2} + \frac{z}{3} \right) du \, dy \, dz$$

by applying the transformation,

$$u = \frac{2u-y}{2}, \quad v = \frac{y}{2}, \quad w = \frac{z}{3}$$

→ Solution:-

$$u = \frac{2u-y}{2} \quad y = 2v \quad z = 3w$$

$$\textcircled{1}, \quad 2u = 2u - y$$

$$\textcircled{2}, \quad 2u = 2u + y$$

$$\textcircled{3}, \quad 2v = 2u + 2v$$

$$\textcircled{4}, \quad u = u + v$$

Now,

$$u = u + v \quad y = 2v \quad z = 3w$$

For $J(u, v)$

$$\frac{\partial u}{\partial u} = \frac{\partial(u+v)}{\partial u} = 1$$

$$\frac{\partial y}{\partial u} = \frac{\partial(2v)}{\partial u} = 0 \quad \frac{\partial z}{\partial w} = \frac{\partial(3w)}{\partial w} = 3$$

$$\frac{\partial u}{\partial v} = \frac{\partial(u+v)}{\partial v} = 1$$

$$\frac{\partial y}{\partial v} = \frac{\partial(2v)}{\partial v} = 2 \quad \frac{\partial z}{\partial w} =$$

$$\frac{\delta u}{\delta u} = \frac{\delta(u+v)}{\delta u} = 1 \quad \frac{\delta y}{\delta u} = \frac{\delta 2v}{\delta u} = 0$$

$$\frac{\delta u}{\delta v} = \frac{\delta(u+v)}{\delta v} = 1 \quad \frac{\delta y}{\delta v} = \frac{\delta 2v}{\delta v} = 2$$

$$\frac{\delta u}{\delta w} = \frac{\delta(u+v)}{\delta w} = 0 \quad \frac{\delta y}{\delta w} = \frac{\delta 2v}{\delta w} = 0$$

$$\frac{\delta z}{\delta u} = \frac{\delta 3w}{\delta u} = 0 \quad \frac{\delta z}{\delta v} = \frac{\delta 3w}{\delta v} = 0 \quad \frac{\delta z}{\delta w} = \frac{\delta 3w}{\delta w} = 3$$

Now,

$$J(u, v) = \begin{vmatrix} 1 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{vmatrix} \quad \begin{array}{l} u = u+v \\ v = 2v \\ z = 3w \end{array}$$

$$= 1(6-0) = 6$$

Now,

xyz eq ⁿ	uvw eq ⁿ	Simplified uvw
$u = \frac{y}{2} + 1$	$u+v = \frac{2v}{2} + 1$	$\therefore u = 1$
$u = \frac{y}{2}$	$u+v = \frac{2v}{2}$	$\therefore u = 0$
$y = 0$	$2v = 0$	$\therefore v = 0$
$y = 4$	$2v = 4$	$\therefore v = 2$
$z = 3$	$3w = 3$	$\therefore w = 1$
$z = 0$	$3w = 0$	$\therefore w = 0$

Then,

$$\begin{aligned}
 & \int_{w=0}^1 \int_{v=0}^2 \int_{u=0}^1 (u+w) |J(u,v)| du dv dw \\
 &= \int_0^1 \int_0^2 \int_0^1 (u+w) \times 6 du dv dw \\
 &= 6 \int_0^1 \int_0^2 \left[\frac{u^2}{2} + uw \right]_0^1 dv dw \\
 &= 6 \int_0^1 \int_0^2 \left[\frac{1}{2} + w \right] dv dw \\
 &= 6 \int_0^1 \left[\frac{1}{2}v + vw \right]_0^2 dw \\
 &= 6 \int_0^1 \left[1 + 2w \right] dw \\
 &= 6 \left\{ \left[w + \frac{2w^2}{2} \right]_0^1 \right\} \\
 &= 6 \left\{ 1 + 1 \right\} \\
 &= 6 \times 2 \\
 &= 12
 \end{aligned}$$

∴ Required answer is 12.

* Evaluate:

$$\iint_R (u-y)^4 e^{u+y} du dy$$

applying transformation, $u = \frac{u+v}{2}$ and $y = \frac{u-v}{2}$

[R is square with vertices (1,0) (2,1) (2,2) and (0,1)]

\rightarrow To find eqⁿ (1,2) & (2,1)

$$y - y_1 = \frac{y_2 - y_1}{u_2 - u_1} (u - u_1)$$

$$\text{a), } (y-2) = \frac{1-2}{2-1} (u-1)$$

$$\text{a), } y-2 = -1 (u-1)$$

$$\text{a), } y-2 = -u+1 \quad \therefore u+y = 3 \text{ like other same. eqⁿ given in qust}$$




Table :-

uv eq ⁿ	uv eq ⁿ	simplified uv eq ⁿ
$u+y = 1$	$\frac{u+v}{2} + \frac{u-v}{2} = \frac{u+v+u-v}{2} = \frac{2u}{2} = 1$	$\therefore u = 1$
$u+y = 3$	$\frac{u+v}{2} + \frac{u-v}{2} = 3$	$\therefore u = 3$
$u-y = 1$	$\frac{u+v - u-v}{2} = 1$	$\therefore v = 1$
$u-y = -1$	$\frac{u+v - u+v}{2} = -1$	$\therefore v = -1$

$$u-y = \frac{u+v}{2} - \frac{u-v}{2} =$$

$$= \frac{u+v}{2} + \frac{u-v}{2}$$

$$\text{a), } u-y = \frac{u+v - u+v}{2}$$

$$\text{a), } u+y = \frac{2u}{2}$$

$$\text{a), } v = u-y$$

$$\therefore u = u+y$$

$$\frac{\delta u}{\delta u} = \frac{\delta \left(\frac{u+v}{2}\right)}{\delta u} = \frac{1}{2} \quad \frac{\delta u}{\delta v} = \frac{\delta \left(\frac{u+v}{2}\right)}{\delta v} = \frac{1}{2} \quad \frac{\delta u}{\delta v} \neq \frac{1}{2}$$

$$\frac{\delta y}{\delta u} = \frac{\delta \left(\frac{u-v}{2}\right)}{\delta u} = \frac{1}{2} \quad \frac{\delta y}{\delta v} = \frac{\delta \left(\frac{u-v}{2}\right)}{\delta v} = -\frac{1}{2}$$

then,

$$J(u, v) = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = \frac{1}{4} - \frac{1}{4} = -\frac{2}{4} = -\frac{1}{2}$$

Now,

$$\iint_R (u-y)^4 e^{u+y} du dy \quad [u-y=v] \quad [u+y=v]$$

$$= \int_{v=-1}^{v=1} \int_{u=1}^{u=3} (v)^4 \cdot e^v |J(u, v)| du dv$$

$$= \int_{-1}^1 v^4 [e^v]_1^3 dv \times -\frac{1}{2}$$

$$= -\frac{1}{2} \int_{-1}^1 [e^3 - e^1] v^4 dv$$

$$= -\frac{1}{2} (e^3 - e^1) \times 2 \left[\frac{v^5}{5} \right]_0^1$$

$$= -\frac{1}{5} (e^3 - e^1) \cancel{\#}$$

Jacobian in Polar co-ordinate :-

$$J(r, \theta) = \begin{vmatrix} \frac{\partial u}{\partial r} & \frac{\partial u}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix}$$

* Find Jacobian of :-

$$a) \ u = r \cos \theta \quad y = r \sin \theta$$

$$J(r, \theta) = \begin{vmatrix} \frac{\partial r \cos \theta}{\partial r} & \frac{\partial r \cos \theta}{\partial \theta} \\ \frac{\partial r \sin \theta}{\partial r} & \frac{\partial r \sin \theta}{\partial \theta} \end{vmatrix}$$

$$= \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix}$$

$$= r \cos^2 \theta + r \sin^2 \theta$$

$$= r (\cos^2 \theta + \sin^2 \theta)$$

$$= r \#.$$

$$J(r, \theta, z) = \begin{vmatrix} \frac{\partial u}{\partial r} & \frac{\partial u}{\partial \theta} & \frac{\partial u}{\partial z} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial z} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial z} \end{vmatrix}$$

* find Jacobian of ;

$$x = r \cos \alpha$$

$$y = r \sin \alpha$$

$$z = z$$

→ Solution,

$$\frac{\delta u}{\delta r} = \frac{\delta r \cos \alpha}{\delta r} = \cos \alpha$$

$$\frac{\delta y}{\delta r} = \frac{\delta r \sin \alpha}{\delta r} = \sin \alpha$$

$$\frac{\delta u}{\delta \alpha} = \frac{\delta r \cos \alpha}{\delta \alpha} = -r \sin \alpha$$

$$\frac{\delta y}{\delta \alpha} = \frac{\delta r \sin \alpha}{\delta \alpha} = r \cos \alpha$$

$$\frac{\delta u}{\delta z} = \frac{\delta r \cos \alpha}{\delta z} = 0$$

$$\frac{\delta y}{\delta z} = \frac{\delta r \sin \alpha}{\delta z} = 0$$

$$\frac{\delta z}{\delta r} = \frac{\delta z}{\delta r} = 0$$

$$\frac{\delta z}{\delta \alpha} = \frac{\delta z}{\delta \alpha} = 0$$

$$\frac{\delta z}{\delta z} = 1$$

$$\therefore J(r, \alpha, z) = \begin{vmatrix} \cos \alpha & -r \sin \alpha & 0 \\ \sin \alpha & r \cos \alpha & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= 1 \mid r \cos^2 \alpha + r \sin^2 \alpha \mid$$

$$= r (\cos^2 \alpha + \sin^2 \alpha)$$

$$= r \#.$$

~~U~~ Partial Derivatives :-

$$\frac{\delta f}{\delta u} = f_u$$

$$\frac{\delta f}{\delta y} = f_y$$

$$\frac{\delta^2 f}{\delta u^2} = f_{uu}$$

$$\frac{\delta^2 f}{\delta y^2} = f_{yy}$$

$$\frac{\delta^2 f}{\delta u \delta y} = f_{uy}$$

$$\frac{\delta^2 f}{\delta y \delta u} = f_{uy}$$

* Find value of $\frac{\partial f}{\partial u} = f_u$ and $\frac{\partial f}{\partial y} = f_y$ at $(4, -5)$
if,

$$f(u, y) = u^2 + 3uy + y - 1.$$

→ solution,

$$f(u, y) = u^2 + 3uy + y - 1$$

$$\therefore \frac{\partial f}{\partial u} = f_u = 2u + 3y \text{ at } (4, -5)$$

$$\begin{aligned} f(u) &= 2 \times 4 + 3 \times (-5) \\ &= 8 - 15 \\ &= -7 \# \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{\partial f}{\partial y} &= f_y = 3u + 1 \text{ at } (4, -5) \\ &= 3 \times 4 + 1 \\ &= 13 \# \end{aligned}$$

* Find $\frac{\partial f}{\partial y}$ if $f(u, y) = u \cos ny$. $\frac{\partial f}{\partial u} = ?$

→ solution,

$$\frac{\partial f}{\partial y} = \frac{\partial u \cos ny}{\partial y}$$


$$= \cancel{u} \frac{\partial \cos ny}{\cancel{\partial y}} \times \frac{\partial ny}{\partial y}$$

$$= u^2 (-\sin ny)$$

$$= -u^2 \sin ny$$

$$\text{Product rule} = \frac{d}{du}(uv) = u \frac{dv}{du} + v \frac{du}{du}$$

$$\text{Quotient rule} = \frac{d}{du}\left(\frac{u}{v}\right) = v \frac{du}{du} - u \frac{dv}{du} / v^2$$



$$\begin{aligned}
 \frac{\delta f}{\delta u} &= fy = \frac{\delta u \cdot \cos ny}{\delta u} \\
 &= u \frac{\delta \cos ny}{\delta u} + \cos ny \frac{\delta u}{\delta u} \\
 &= u \frac{\cos ny \times \sin y}{\sin y} + \cos ny \times 1 \\
 &= ny(-\sin ny) + \cos ny \\
 &= -ny \sin ny + \cos ny
 \end{aligned}$$

* Find f_u and f_y if $f(u, y) = \frac{2y}{y + \cos u} \Rightarrow$ quotient

$$f_u = \frac{\delta}{\delta u} \left(\frac{2y}{y + \cos u} \right)$$

$$= (y + \cos u) \frac{\delta 2y}{\delta u} - 2y \frac{\delta(y + \cos u)}{\delta u}$$

$$= \frac{(y + \cos u) \times 0 - 2y (0 + (-\sin u))}{(y + \cos u)^2}$$

$$= \frac{(y + \cos u) \times 0 - 2y (0 + (-\sin u))}{(y + \cos u)^2}$$

$$\therefore f_u = \frac{2y \sin u}{(y + \cos u)^2}$$

$$\begin{aligned}
 f_y &= \frac{\delta}{\delta y} \left(\frac{2y}{y + \cos u} \right) = (y + \cos u) \frac{\delta 2y}{\delta y} - 2y \frac{\delta(y + \cos u)}{\delta y} \\
 &= (y + \cos u) \times 2 - 2y (1 + 0) / v^2 \\
 &= 2(y + \cos u - y) / v^2
 \end{aligned}$$

$$\therefore f_y = \frac{2 \cos u}{(y + \cos u)^2} \neq .$$

* find $\frac{\delta f}{\delta u} = f_u$ and $\frac{\delta f}{\delta y} = f_y$.

$$\text{a)} f(u, y) = \sqrt{u^2 + y^2}$$

$$\begin{aligned} \rightarrow f(u) &= \frac{\delta (u^2 + y^2)^{1/2}}{\delta u} = \frac{1}{2} (u^2 + y^2)^{1/2-1} \times 2u \\ &= \frac{\delta (u^2 + y^2)^{1/2}}{\delta u} \times \frac{\delta (u^2 + y^2)}{\delta u} = \frac{u}{\sqrt{u^2 + y^2}} \end{aligned}$$

$$\therefore f(u) = \frac{u}{\sqrt{u^2 + y^2}} \neq .$$

$$\begin{aligned} f_y &= \frac{\delta (u^2 + y^2)^{1/2}}{\delta y} \\ &= \frac{\delta (u^2 + y^2)^{1/2}}{\delta (u^2 + y^2)} \times \frac{\delta (u^2 + y^2)}{\delta y} \\ &= \frac{1}{2\sqrt{u^2 + y^2}} \times 2y \end{aligned}$$

$$\therefore f_y = \frac{y}{\sqrt{u^2 + y^2}} \neq .$$

$$\text{b)} f(u, y) = e^{u+y+1}$$

$$\therefore f_u = \frac{\delta e^{u+y+1}}{\delta (u+y+1)} \times \frac{\delta (u+y+1)}{\delta u}$$

$$= e^{u+y+1} \times 1$$

$$f_u = e^{u+y+1} \quad \#$$

$$f_y = \frac{\delta(e^{u+y+1})}{\delta(u+y+1)} \times \frac{\delta(u+y+1)}{\delta y}$$

$$\therefore f_y = e^{u+y+1} \quad \#$$

$$\therefore f(u, y) = \ln(u+y)$$

→ Solution,

$$f_u = \frac{\delta \ln(u+y)}{\delta(u+y)} \times \frac{\delta(u+y)}{\delta u}$$

$$\frac{d \ln u}{du} = \frac{1}{u}$$

$$= \frac{1}{u+y} \times \frac{1}{1}$$

$$\therefore f(u) = \frac{1}{u+y} \quad \#$$

$$f_y = \frac{\delta \ln(u+y)}{\delta(u+y)} \times \frac{\delta(u+y)}{\delta y}$$

$$\therefore f_y = \frac{1}{u+y} \quad \#$$

Implicit Differentiation :

* Find $\frac{\delta z}{\delta u}$ if the $yz - \ln z = u+y$

→ Solution,

$$\frac{\delta(yz - \ln z)}{\delta u} = \frac{\delta(u+y)}{\delta u}$$

$$\text{a. } \frac{\delta yz}{\delta u} - \frac{\delta \ln z}{\delta u} = 1$$

$$a, y \frac{\partial z}{\partial u} - \frac{\partial(\ln z)}{\partial z} \times \frac{\partial z}{\partial u} = 1$$

$$a, \frac{\partial z}{\partial u} \left(y - \frac{1}{z} \right) = 1$$

$$\therefore \frac{\partial z}{\partial u} = \left(\frac{z}{yz-1} \right) \#.$$

* If $f(x, y) = x \cos y + y e^x$

→ find, $\frac{\partial^2 f}{\partial x^2} = f_{xx}, f_{yy}, f_{xy}, f_{yx}$

→ Solution,

for f_{xx}

we have to do first f_x then do f_x again with first for answer.

i.e.,

$$f_x = \frac{\partial(x \cos y + y e^x)}{\partial x}$$

$$= \frac{\partial x \cos y}{\partial x} + \frac{\partial y e^x}{\partial x}$$

$$\therefore f_x = \cos y + y e^x$$

$$\therefore f_{xx} = \frac{\partial(\cos y + y e^x)}{\partial x}$$

$$= \frac{\partial \cos y}{\partial x} + \frac{\partial(y e^x)}{\partial x}$$

$$= 0 + y e^x$$

$$\therefore f_{xx} = y e^x$$

$$f_{yy} = ?$$

$$\begin{aligned} f_y &= \frac{\delta(u \cos y + y e^u)}{\delta y} \\ &= \frac{\delta u \cos y}{\delta y} + \frac{\delta(y e^u)}{\delta y} \\ &= -u \sin y + e^u \end{aligned}$$

$$f_{yy} = \frac{\delta(-u \sin y + e^u)}{\delta y}$$

$$= -u \cos y$$

$$f_{yy} = \frac{\delta^2(f_{xy})}{\delta y \delta u} = f_{xy} - \text{second do } f_y$$

$$\therefore f_{xy} = \cos y + y e^u$$

$$\therefore f_{xy} = \frac{\delta(\cos y + y e^u)}{\delta y}$$

$$= -\sin y + e^u$$

$$f_y = -u \sin y + e^u$$

$$f_{yy} = \frac{\delta(-u \sin y + e^u)}{\delta u}$$

$$= -\sin y + e^u$$

$$\boxed{\therefore f_{xy} = f_{yy} = -\sin y + e^u}$$

✓ This is called Euler's theorem / mixed derivative Theorem.

* Euler's Theorem (Mixed Derivative Theorem):

$$f_{uy} = f_{yu}$$

$$\frac{\delta^2 f}{\delta y \delta u} = \frac{\delta^2 f}{\delta u \delta y}$$

* Find $f_{uyyz} \xrightarrow{4^{\text{th}}} \xrightarrow{3^{\text{rd}}} f$,
 $\xrightarrow{1^{\text{st}}} \xrightarrow{2^{\text{nd}}} f$

$$f(u, y, z) = 1 - 2uy^2z + u^2y$$

→ Solution,

first do from first y , then $u \dots z$.

$$\begin{aligned} f_y &= \frac{\delta (1 - 2uy^2z + u^2y)}{\delta y} \\ &= -2 \times 2uyz + u^2 \end{aligned}$$

$$\begin{aligned} f_{yu} &= \frac{\delta (-4uyz + u^2)}{\delta u} \\ &= -4yz + 2u \end{aligned}$$

$$\begin{aligned} &\& f_{uyy} = \frac{\delta (-4yz + 2u)}{\delta y} \\ &= -4z + 0 \\ &= -4z \end{aligned}$$

$$f_{uyyz} = \frac{\delta (-4z)}{\delta z}$$

$$= -4$$

$$\left\{ \because f_{uyyz} = -4 \right\} \text{✓}$$

* Verify Euler's Theorem for the following:

a) $f(x, y) = x^2 + 5xy + \sin x + 7e^x$

→ Solution,

To verify Euler's theorem,
 $f_{xy} = f_{yx}$

$$\therefore f_x = \frac{\partial (x^2 + 5xy + \sin x + 7e^x)}{\partial x}$$

$$= 2x + 5y + \cos x + 7e^x$$

$$\therefore f_{xy} = \frac{\partial (2x + 5y + \cos x + 7e^x)}{\partial y}$$

$$= 0 + 5 + 0 + 0$$

$$= 5$$

$$f_y = \frac{\partial (x^2 + 5xy + \sin x + 7e^x)}{\partial y}$$

$$= 0 + 5x + 0 + 0$$

$$= 5x$$

$$\therefore f_{yx} = \frac{\partial (5x)}{\partial x}$$

$$= 5$$

$\therefore f_{xy} = f_{yx} = 5$ Hence Euler's Theorem is Verified.

b) $f(x, y) = y + x^2y + 4y^3 - \ln(y^2 + 1)$

→ Solution,

$$f_x = \frac{\partial (y + x^2y + 4y^3 - \ln(y^2 + 1))}{\partial x}$$

$$= 0 + 2xy + 0 - 0 = 2xy$$

$$\therefore f_{xy} = \frac{\partial (2xy)}{\partial y} = 2x$$

Similarly,

$$f_y = \frac{\delta}{\delta y} (y + u^2 y + 4y^3 - \ln(y^2 + 1))$$

$$= \frac{\delta y}{\delta y} + \frac{\delta(u^2 y)}{\delta y} + \frac{\delta(4y^3)}{\delta y} - \frac{\delta(\ln(y^2 + 1))}{\delta(y^2 + 1)} \times \frac{\delta(y^2 + 1)}{\delta y}$$

$$= 1 + u^2 + 4 \times 3y^2 - \frac{1}{(y^2 + 1)} \times 2y$$

$$\therefore f_{yu} = \frac{\delta}{\delta u} \left(1 + u^2 + 12y^2 - \frac{1}{(y^2 + 1)} \times 2y \right)$$

$$= 0 + 2u + 0 - 0$$

$$= 2u$$

$$\therefore f_{uy} = f_{yu} = 2u$$

Hence Euler's theorem is verified.

c) $f(u, y) = u \ln u y$

→ Solution,

$$f_u = \frac{\delta(u \ln u y)}{\delta u} \quad \frac{d(u \cdot v)}{du} = u \frac{dv}{du} + v \frac{du}{du}$$

$$= u \frac{\delta \ln u y}{\delta u} \times \frac{\delta u y}{\delta u} + \ln u y \frac{\delta u}{\delta u}$$

$$= \frac{u}{u y} \times y + \ln u y$$

$$= 1 + \ln u y$$

$$\therefore f_{uy} = \frac{\delta(1 + \ln u y)}{\delta y}$$

$$= 0 + \frac{\delta(1 + \ln u y)}{\delta u y} \times \frac{\delta u y}{\delta y}$$

$$= \frac{1}{u y} \times u \quad \therefore f_{uy} = \frac{1}{y}$$

(Similarly,

$$f_y = \frac{\delta(u \ln ny)}{\delta y}$$

$$= u \frac{\delta(\ln ny)}{\delta(u)} \times \frac{\delta u}{\delta y} + \ln ny \frac{\delta u}{\delta y}$$

$$= u \frac{1}{u} \times 0 + 0$$

$$= \frac{u}{y}$$

$$\therefore f_{yu} = \frac{\delta(\frac{u}{y})}{\delta u}$$

$$\frac{d}{du} \left(\frac{u}{v} \right) = v \frac{du}{du} - u \frac{dv}{du} \underbrace{v^2}_{v^2}$$

$$= y \frac{\delta u}{\delta u} - u \frac{\delta y}{\delta u}$$

$$y^2$$

$$= \frac{y \times 1}{y^2} - u \times 0$$

$$= \frac{y}{y^2} = \frac{1}{y}$$

$$\therefore f_{yu} = f_{yu} = \frac{1}{y}$$

Hence Euler's theorem is verified.

$$\text{d}f(u, y) = u \sin y + y \sin u + ny$$

→ solution,

$$f_u = \frac{\delta(u \sin y + y \sin u + ny)}{\delta u}$$

$$= \sin y + y \cos u + y$$

$$\therefore f_{uy} = \frac{\delta(\sin y + y \cos u + y)}{\delta y}$$

$$\therefore f_{uy} = \cos y + \cos u + 1$$

$$f_y = \frac{\partial (u \sin y + y \sin u + uy)}{\partial y}$$

$$= u \cos y + \sin u + u$$

$$\therefore f_{yu} = \frac{\partial (u \cos y + \sin u + u)}{\partial u}$$

$$= \cos y + \cos u + 1$$

$$\therefore f_{uy} = f_{yu} = \cos y + \cos u + 1$$

Hence Euler's theorem is verified.

$$\Rightarrow f(u, y) = e^u + u \ln y + y \ln u$$

→ Solution,

$$f_u = \frac{\partial (e^u + u \ln y + y \ln u)}{\partial u}$$

$$= e^u + \ln y + y \times \frac{1}{u}$$

$$f_{uy} = \frac{\partial (e^u + \ln y + \frac{y}{u})}{\partial y}$$

$$= 0 + \frac{1}{y} + \frac{1}{u}$$

$$\therefore f_{uy} = \frac{1}{u} + \frac{1}{y}$$


$$f_y = \frac{\partial (e^u + u \ln y + y \ln u)}{\partial y} = 0 + \frac{u}{y} + \ln u$$

$$f_{yu} = \frac{\partial (\frac{u}{y} + \ln u)}{\partial u}$$

$$= \frac{1}{y} + \frac{1}{u} = \frac{1}{u} + \frac{1}{y}$$

$$\therefore f_{uy} = f_{yu} = \frac{1}{u} + \frac{1}{y}$$

Hence Euler's theorem is verified.



* Find the value of $\frac{\partial z}{\partial u}$ at point $(1, 1, 1)$ if eqⁿ $uy + z^3u - 2yz = 0$.

→ Solution:-

z is function,

so we cannot put z as constant.

$$uy + z^3u - 2yz = 0 \quad (\text{Product rule})$$

Differentiating on both side, we get

$$\frac{\partial (uy)}{\partial u} + \frac{\partial (z^3u)}{\partial u} - 2 \frac{\partial (yz)}{\partial u} = 0$$

$$\textcircled{1}, \quad y + z^3 \frac{\partial u}{\partial u} + u \frac{\partial z^3}{\partial z} \times \frac{\partial z}{\partial u} - 2 \left(\frac{\partial yz}{\partial u} + z \frac{\partial y}{\partial u} \right) = 0$$

$$\textcircled{2}, \quad y + z^3 + 3uz^2 \frac{\partial z}{\partial u} - 2y \frac{\partial z}{\partial u} + z \times 0 = 0$$

$$\textcircled{3}, \quad y + z^3 + \frac{\partial z}{\partial u} (3uz^2 - 2y) = 0$$

$$\textcircled{4}, \quad \frac{\partial z}{\partial u} (3uz^2 - 2y) = -y - z^3$$

$$\textcircled{5}, \quad \frac{\partial z}{\partial u} = \frac{-y - z^3}{3uz^2 - 2y} \quad \text{At } (1, 1, 1)$$

$$\therefore \frac{\partial z}{\partial u} = \frac{-1 - 1}{3 \times 1 \times 1 - 2 \times 1} = \frac{-2}{3 - 2} = -2 \quad \#.$$

* Find the value of $\frac{\partial u}{\partial z}$ at $(1, -1, -3)$ if eqⁿ $uz + y \ln u - u^2 + 4 = 0$

→ Solution,

$$uz + y \ln u - u^2 + 4 = 0$$

Differentiating both side, we get,

$$\frac{\delta(uz)}{\delta z} \frac{\delta(ylnu)}{\delta z} \xrightarrow{\text{Product}} \frac{\delta(uz)}{\delta z} + \frac{\delta(ylnu)}{\delta z} - \frac{\delta u^2}{\delta z} + \frac{\delta u}{\delta z} = 0$$

$$\text{a. } u \frac{\delta u}{\delta z} + z \frac{\delta u}{\delta z} + y \frac{\delta ln u}{\delta z} \times \frac{\delta u}{\delta z} - \frac{\delta u^2}{\delta z} + \frac{\delta u}{\delta z} = 0$$

$$\text{a. } u + z \frac{\delta u}{\delta z} + \frac{y}{u} \frac{\delta u}{\delta z} - 2u \frac{\delta u}{\delta z} = 0$$

$$\text{a. } u + \frac{\delta u}{\delta z} (z + \frac{y}{u} - 2u) = 0$$

at point $(1, -1, -3)$

$$\frac{\delta u}{\delta z} \left(-3 + -\frac{1}{1} - 2 \times 1 \right) = -1$$

$$\text{a. } \frac{\delta u}{\delta z} = -\frac{1}{(-3 - 1 - 2)} = \frac{1}{-6} = \frac{1}{6}$$

Hence the required value of $\frac{\delta u}{\delta z}$ is $\frac{1}{6}$.

One Dimension Wave Equation :-

$$\frac{\delta^2}{\delta t^2} = c^2 \frac{\delta^2 u}{\delta x^2}$$

$$\text{i.e. } w_{tt} = c^2 w_{xx}$$

where,

w = wave height

t = time Variable

x = distance variable

c = Velocity with which waves propagated.

are.

* Show that the following functions are solution of the following wave eqⁿ:

$$9.7 \quad w = \sin(u+ct)$$

→ Solution,

To be the solution of wave eqⁿ,

$$w_{tt} = c^2 w_{uu}$$

Taking L.H.S

$$w_{tt} = \frac{\delta(\sin(u+ct))}{\delta t} = \frac{\delta(\sin(u+ct)) \times \delta(u+ct)}{\delta(u+ct)} \times \frac{1}{\delta t}$$

$$= \cos(u+ct) \times c$$

$$= c \cos(u+ct)$$

$$w_{tt} = \frac{\delta c \cos(u+ct)}{\delta t} = c \frac{\delta(\cos(u+ct))}{\delta(u+ct)} \times \frac{\delta(u+ct)}{\delta t}$$

$$= c(-\sin(u+ct)) \times c$$

$$\therefore w_{tt} = -c^2 \sin(u+ct) \#.$$

Taking R.H.S.

$$= c^2 w_{uu}$$

$$w_{uu} = \frac{\delta(\sin(u+ct))}{\delta u} = \frac{\delta(\sin(u+ct)) \times \delta(u+ct)}{\delta(u+ct)} \times \frac{1}{\delta u}$$

$$= \cos(u+ct) \times 1 = \cos(u+ct)$$

$$w_{uu} = \frac{\delta(\cos(u+ct))}{\delta u} = \frac{\delta(\cos(u+ct)) \times \delta(u+ct)}{\delta(u+ct)} \times \frac{1}{\delta u}$$

$$= -\sin(u+ct) =$$

$$\therefore c^2 w_{uu} = -c^2 \sin(u+ct) \#.$$

$$\boxed{\therefore L.H.S = R.H.S = -c^2 \sin(u+ct)}$$

Hence the given function is the solⁿ of wave eqⁿ.

$$\text{by } w = \cos(2u + 2ct)$$

→ Solution,

To be solⁿ of wave eqⁿ, it should satisfy,

$$w_{tt} = c^2 w_{uu}$$

Taking L.H.S = w_{tt}

$$\begin{aligned} w_{tt} &= \frac{\delta(\cos(2u + 2ct)) \times \delta(2u + 2ct)}{\delta t} \\ &= -\sin(2u + 2ct) \times 2c \\ &= -2c \sin(2u + 2ct) \end{aligned}$$

$$\begin{aligned} w_{tt} &= \frac{\delta(-2c \sin(2u + 2ct))}{\delta t} \\ &= -2c \frac{\delta(\sin(2u + 2ct)) \times \delta(2u + 2ct)}{\delta t} \\ &= -2c \cos(2u + 2ct) \times 2c \\ &= -4c^2 \cos(2u + 2ct) \end{aligned}$$

Taking R.H.S = $w_{uu} c^2$

$$\begin{aligned} w_{uu} &= \frac{\delta(\cos(2u + 2ct))}{\delta u} = \frac{\delta(\cos(2u + 2ct)) \times \delta(2u + 2ct)}{\delta(2u + 2ct) \delta u} \\ &= -2 \sin(2u + 2ct) \times 1 \\ w_{uu} &= \frac{\delta(-2 \sin(2u + 2ct))}{\delta u} \\ &= \frac{\delta(-2 \sin(2u + 2ct)) \times \delta(2u + 2ct)}{\delta(2u + 2ct) \delta u} \\ &= -2 \cos(2u + 2ct) \times 2 \\ &= -4 \cos(2u + 2ct) \end{aligned}$$

$$\therefore c^2 w_{uu} = -4c^2 \cos(2u + 2ct)$$

$$\therefore \text{L.H.S} = \text{R.H.S} = -4c^2 \cos(2u + 2ct)$$

Hence the given function is the solⁿ of wave eqⁿ.

Verify Euler's theorem for

a) $f(u, y) = y + \left(\frac{u}{y}\right)$

→ Solution,

$$f_u = \frac{\partial}{\partial u} \left(y + \frac{u}{y} \right)$$

$$= 0 + \frac{1}{y} \times 1 = \frac{1}{y}$$

$$\therefore f_{uy} = \frac{\partial}{\partial y} \left(\frac{1}{y} \right)$$

$$= \frac{\partial (y^{-1})}{\partial y} = -\frac{1}{y^2}$$

$$f_y = \frac{\partial}{\partial y} \left(y + \frac{u}{y} \right)$$

$$= 1 + u(-y^{-2})$$

$$= 1 + \left(-\frac{u}{y^2}\right)$$

$$\therefore f_{yu} = \frac{\partial}{\partial u} \left(1 - \frac{u}{y^2} \right)$$

$$= 0 - \frac{1}{y^2} \times 1$$

$$= -\frac{1}{y^2}$$

$$\therefore f_{uy} = f_{yu} = -\frac{1}{y^2}$$

Hence, Euler's theorem is verified.

b) $f(u, y) = u \sin y + e^y$

→ Solution,

$$f_u = \frac{\partial}{\partial u} (u \sin y + e^y) = \sin y$$

$$f_y = \frac{\partial}{\partial y} (u \sin y + e^y)$$

$$\therefore f_{uy} = \frac{\partial}{\partial y} (\sin y) = \cos y$$

$$\therefore f_{yu} = \frac{\partial}{\partial u} (\sin y + e^y) = 1 \times \cos y = \cos y$$

$$\therefore f_{uy} = f_{yu} = \cos y$$

Hence Euler's theorem is verified.

classmate
Date _____
Page _____

$$\Leftrightarrow f(x,y) = \tan^{-1}(\frac{y}{x}) \quad [\text{Note: } \frac{d}{du}(\tan^{-1}u) = \frac{1}{1+u^2}]$$

→ Solution,

$$f_u = \frac{\partial}{\partial u} \left(\tan^{-1} \left(\frac{y}{u} \right) \right) \times \frac{\partial}{\partial u} \left(\frac{y}{u} \right)$$

$$= \frac{1}{1 + \left(\frac{y}{u} \right)^2} \times -\frac{y}{u^2} = \frac{1}{u^2 + y^2} \times -\frac{y}{u^2}$$

$$f_{uy} // f_u = \frac{1}{1} \times \frac{u^2}{u^2 + y^2} \times \frac{(-y)}{u^2}$$

$$\therefore f_u = -\frac{y}{u^2 + y^2}$$

$$f_{uy} = \frac{\partial}{\partial y} \left(-\frac{y}{u^2 + y^2} \right) \quad [\text{quotient rule}]$$

$$= (u^2 + y^2) \frac{\partial}{\partial y} \left(-\frac{y}{u^2 + y^2} \right) - (-y) \frac{\partial}{\partial y} \left(\frac{u^2 + y^2}{u^2 + y^2} \right)$$

$$= (u^2 + y^2) \times (-1) + y \left(0 + 2y \right) / (u^2 + y^2)^2$$

$$= -u^2 - y^2 + 2y^2 / (u^2 + y^2)^2$$

$$= -u^2 + y^2 / (u^2 + y^2)^2$$

$$f_y = \frac{\partial}{\partial y} \left(\tan^{-1} \left(\frac{y}{u} \right) \right) \times \frac{\partial}{\partial y} \left(\frac{y}{u} \right)$$

$$= \frac{1}{1 + \left(\frac{y}{u} \right)^2} \times \frac{1}{u}$$

$$= \frac{u^2}{u^2 + y^2} \times \frac{1}{u} = \frac{u}{u^2 + y^2}$$

$$\therefore f_{yu} = \frac{\partial}{\partial u} \left(\frac{u}{u^2+y^2} \right)$$

$$= (u^2+y^2) \frac{\partial u}{\partial u} + u \frac{\partial (u^2+y^2)}{\partial u} / (u^2+y^2)^2$$

$$= (u^2+y^2) \times 1 + u (2u+0) / (u^2+y^2)^2$$

$$= \frac{u^2+y^2 - 2u^2}{(u^2+y^2)^2}$$

$$= \frac{-u^2+y^2}{(u^2+y^2)}$$

$$\therefore f_{ny} = f_{yu} = \boxed{-\frac{u^2+y^2}{(u^2+y^2)}}$$

Hence Euler's theorem is verified.

$$\text{d) } f(u, y) = y + u^2 y + 4y^3 - \ln(y^2 + 1)$$

\rightarrow Solution,

$$f_u = \frac{\partial (y + u^2 y + 4y^3 - \ln(y^2 + 1))}{\partial u}$$

$$= 0 + 2uy + 0 - 0$$

$$= 2uy$$

$$f_{ny} = \frac{\partial (2uy)}{\partial y} = 2u$$

$$f_y = \frac{\partial (y + u^2 y + 4y^3 - \ln(y^2 + 1))}{\partial y} = \frac{\partial (\ln(y^2 + 1))}{\partial (y^2 + 1)} \times \frac{\partial (y^2 + 1)}{\partial y}$$

$$= 1 + u^2 + 12y^2 - \frac{1}{(y^2 + 1)} \times 2y$$

$$= \frac{1}{(y^2 + 1)} \times 2y$$

$$8y f_{yu} = \frac{\partial}{\partial u} \left(1 + u^2 + 12y^2 - \frac{1}{(y^2 + 1)} \right) = 8 \times 0 + 2u - \frac{1}{(y^2 + 1)} \times 2y$$

$$= 0 + 2u + 0 - 0 = 2u$$

$$\therefore f_{ny} = f_{yu} = 2u \quad \text{Hence Euler's theorem is verified.}$$

partial deriv. \rightarrow we take constant which is not w.r.t.
 Derivatives \rightarrow we do chain rule or.... No constant is taken

classmate
Date _____
Page _____

chain Rule's (2-D)

$$\frac{d\omega}{dt} = \frac{\partial f}{\partial u} \times \frac{du}{dt} + \frac{\partial f}{\partial y} \times \frac{dy}{dt}$$

where $\omega = f(u, y)$ = function.

$$\frac{d\omega}{dt} = \frac{\partial f}{\partial u} \times \frac{du}{dt} + \frac{\partial f}{\partial y} \times \frac{dy}{dt} + \frac{\partial f}{\partial z} \times \frac{dz}{dt}$$

where $\omega = f(u, y, z)$ = function.

* Find $\frac{d\omega}{dt}$

$$1. \omega = u^2 + y^2, u = \cos t, y = \sin t$$

\rightarrow Solution,

$$\frac{d\omega}{dt} = \frac{\partial \omega}{\partial u} \times \frac{du}{dt} + \frac{\partial \omega}{\partial y} \times \frac{dy}{dt}$$

$$= \frac{\partial (u^2 + y^2)}{\partial u} \times \frac{d(\cos t)}{dt} + \frac{\partial (u^2 + y^2)}{\partial y} \times \frac{d(\sin t)}{dt}$$

$$= 2u \times (-\sin t) + 2y \cos t$$

Put value of u & y

$$= 2 \cos t \cdot (-\sin t) + 2 \sin t \cdot \cos t$$

$$= -2 \sin t \cdot \cos t + 2 \sin t \cdot \cos t$$

$$= 0 \#.$$

$$2. \omega = u^2 + y^2, u = \cos t + \sin t, y = \cos t - \sin t$$

\rightarrow Solution,

$$\frac{d\omega}{dt} = \frac{\partial \omega}{\partial u} \times \frac{du}{dt} + \frac{\partial \omega}{\partial y} \times \frac{dy}{dt}$$

$$= \frac{\partial (u^2 + y^2)}{\partial u} \times \frac{d(\cos t + \sin t)}{dt} + \frac{\partial (u^2 + y^2)}{\partial y} \times \frac{d(\cos t - \sin t)}{dt}$$

$$= 2u \times ((-\sin t) + \cos t) + 2y \times ((-\cancel{\cos t}) - \sin t)$$

$$= 2(\cos t + \sin t) \times (\cos t - \sin t) + 2(\cos t - \sin t) (-\cos t - \sin t)$$

$$= 2(\cos^2 t - \sin^2 t) + 2(\cos t \cancel{+ \cos t} - \sin t \cancel{- \sin t}) \rightarrow \text{common}$$

$$+ 2(\cos t - \sin t) - (\cos t + \sin t)$$

$$= 2(\cos^2 t - \sin^2 t) - 2(\cos^2 t - \sin^2 t)$$

$$= 0 \quad \#$$

$$3.7 \omega = u^2 y - y^2, u = \sin t, y = e^t \text{ at } t=0$$

→ Solution,

$$\frac{du}{dt} = \frac{\partial \omega}{\partial u} \times \frac{du}{dt} + \frac{\partial \omega}{\partial y} \times \frac{dy}{dt}$$

$$= \frac{\partial (u^2 y - y^2)}{\partial u} \times \frac{d(\sin t)}{dt} + \frac{\partial (u^2 y - y^2)}{\partial y} \times \frac{d(e^t)}{dt}$$

$$= 2uy \times \cos t + (u^2 - 2y)e^t$$

$$= 2\sin t \cdot e^t \times \cos t + (\sin^2 t - 2(e^t)) \cdot e^t$$

$$\text{at } t=0$$

$$= 2\sin 0^\circ \cdot e^0 \times \cos 0^\circ + (\sin 0^\circ)^2 - 2(e^0) \cdot e^0$$

$$= 2 \times 0 + 0 - 2 \times 1 \times 1$$

$$= -2 \quad \#$$

$$4.7 \omega = uy + z, u = \cos t, y = \sin t, z = t, \text{ at } t=0$$

→ Solution

$$\frac{d\omega}{dt} = \frac{\partial (\omega)}{\partial u} \times \frac{du}{dt} + \frac{\partial (\omega)}{\partial y} \times \frac{dy}{dt} + \frac{\partial (\omega)}{\partial z} \times \frac{dz}{dt}$$

$$\begin{aligned}
 &= \frac{\partial w}{\partial u} (uy+z) \times \frac{d(\cos t)}{dt} + \frac{\partial w}{\partial y} (uy+z) \times \frac{d(\sin t)}{dt} \\
 &\quad + \frac{\partial w}{\partial z} (uy+z) \times \frac{d(t)}{dt}
 \end{aligned}$$

$$= y \times -\sin t + u \times \cos t + 1 \times 1$$

$$= \sin t \times (-\sin t) + \cos t \times \cos t + 1$$

$$= -\sin^2 t + \cos^2 t + 1$$

$$\text{at } t=0$$

$$= (-\sin 0)^2 + (\cos 0)^2 + 1$$

$$= 0 + 1 + 1$$

$$= 2 \#.$$

chain Rule :- (3-D) all partial

For two independent variables (r,s)

and three intermediate (u,y,z) variables

$w = f(u,y,z)$, $u = g(r,s)$, $y = h(r,s)$, $z = k(r,s)$

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial u} \times \frac{\partial u}{\partial r} + \frac{\partial w}{\partial y} \times \frac{\partial y}{\partial r} + \frac{\partial w}{\partial z} \times \frac{\partial z}{\partial r}$$

$$\frac{\partial w}{\partial s} = \frac{\partial w}{\partial u} \times \frac{\partial u}{\partial s} + \frac{\partial w}{\partial y} \times \frac{\partial y}{\partial s} + \frac{\partial w}{\partial z} \times \frac{\partial z}{\partial s}$$

* Find $\frac{\partial w}{\partial r}$ and $\frac{\partial w}{\partial s}$ if,

$$a) \omega = u + 2y + z^2, u = \frac{r}{s}, y = r^2 + \ln s, z = 2r$$

→ Solution:

$$\begin{aligned}
 \frac{\delta \omega}{\delta r} &= \frac{\delta \omega}{\delta u} \times \frac{\delta u}{\delta r} + \frac{\delta \omega}{\delta y} \times \frac{\delta y}{\delta r} + \frac{\delta \omega}{\delta z} \times \frac{\delta z}{\delta r} \\
 &= \frac{\delta(u+2y+z^2)}{\delta u} \times \frac{\delta(\frac{r}{s})}{\delta r} + \frac{\delta(u+2y+z^2)}{\delta y} \times \frac{\delta(r^2+\ln s)}{\delta r} \\
 &\quad + \frac{\delta(u+2y+z^2)}{\delta z} \times \frac{\delta(2r)}{\delta r} \quad \begin{array}{l} \frac{\delta u}{\delta r} = 0 \\ \frac{\delta y}{\delta r} = 2 \\ \frac{\delta z}{\delta r} = 2 \end{array} \\
 &= 1 * \frac{1}{s} + 2 \times 2r + 2z \times 2 \\
 &= \frac{1}{s} + 4r + 4z \quad [z = 2r] \\
 &= \frac{1}{s} + 4r + 4 \times 2r
 \end{aligned}$$

$$\boxed{\frac{\delta \omega}{\delta r} = \frac{1}{s} + 12r}$$

$$\begin{aligned}
 \frac{\delta \omega}{\delta s} &= \frac{\delta \omega}{\delta u} \times \frac{\delta u}{\delta s} + \frac{\delta \omega}{\delta y} \times \frac{\delta y}{\delta s} + \frac{\delta \omega}{\delta z} \times \frac{\delta z}{\delta s} \\
 &= \frac{\delta(u+2y+z^2)}{\delta u} \times \frac{\delta(\frac{r}{s})}{\delta s} + \frac{\delta(u+2y+z^2)}{\delta y} \times \frac{\delta(r^2+\ln s)}{\delta s} \\
 &\quad + \frac{\delta(u+2y+z^2)}{\delta z} \times \frac{\delta(2r)}{\delta s}
 \end{aligned}$$

$$= 1 \times \left(-\frac{r}{s^2}\right) + 2 \times \frac{1}{s} + 2z \times 0$$

$$= -\frac{r}{s^2} + \frac{2}{s}$$

$$\boxed{\therefore \frac{\delta \omega}{\delta s} = -\frac{r}{s^2} + \frac{2}{s}}$$

* Define partial differential eqⁿ of second order with suitable e.g.

→ Partial differential equation

if a dependent variable is a function of two or more than two independent variable, then an equation involving with partial coefficient it is known as partial differential eqⁿ. This is the relation of dependent variable, independent variable and partial differential coefficient.

$\frac{d^2z}{du^2} + \frac{d^2}{dudy} + \frac{2d^2}{dy^2} = 0$ is second order partial

$$\boxed{\frac{d^2z}{du^2} + \frac{d^2}{dudy} + \frac{2d^2}{dy^2} = 0}$$

* find $\frac{\delta w}{\delta r}$ and $\frac{\delta w}{\delta s}$ if :-

$$\Rightarrow w = u^2 + y^2, u = r-s, y = r+s$$

→ Solution,

$$\begin{aligned} \frac{\delta w}{\delta r} &= \frac{\delta w}{\delta u} \times \frac{\delta u}{\delta r} + \frac{\delta w}{\delta y} \times \frac{\delta y}{\delta r} \\ &= \frac{\delta(u^2+y^2)}{\delta u} \times \frac{\delta(r-s)}{\delta r} + \frac{\delta(u^2+y^2)}{\delta y} \times \frac{\delta(r+s)}{\delta r} \\ &= 2u \times 1 + 2y \times 1 \\ &= 2(r-s) + 2(r+s) \\ &= 2r - 2s + 2r + 2s \\ &= 4r \end{aligned}$$

$$\begin{aligned} \frac{\delta w}{\delta s} &= \frac{\delta w}{\delta u} \times \frac{\delta u}{\delta s} + \frac{\delta w}{\delta y} \times \frac{\delta y}{\delta s} \\ &= \frac{\delta(u^2+y^2)}{\delta u} \times \frac{\delta(r-s)}{\delta s} + \frac{\delta(u^2+y^2)}{\delta y} \times \frac{\delta(r+s)}{\delta s} \end{aligned}$$

$$\begin{aligned}
 &= 2u \times (-1) + 2y \times 1 \\
 &= -2(r-s) + 2(r+s) \\
 &= -2r + 2s + 2r + 2s \\
 &= 4s
 \end{aligned}$$

$$\therefore \frac{\delta \omega}{\delta r} = 4r$$

$$\therefore \frac{\delta \omega}{\delta s} = 4s \quad \#$$

$$b) \omega = (u+y+z)^2, u = r-s, y = \cos(r+s), z = \sin(r+s)$$

→ Solution, when $r=1, s=-1$

$$\frac{d\omega}{dr} = \frac{\delta(\omega)}{\delta u} \times \frac{\delta u}{\delta r} + \frac{\delta \omega}{\delta y} \times \frac{\delta y}{\delta r} + \frac{\delta \omega}{\delta z} \times \frac{\delta z}{\delta r}$$

$$\begin{aligned}
 &= \frac{\delta(u+y+z)^2}{\delta u} \times \frac{\delta(u+y+z)}{\delta r} + \frac{\delta(u+y+z)^2}{\delta y} \times \frac{\delta(r-s)}{\delta r} + \frac{\delta(u+y+z)^2}{\delta z} \times \frac{\delta(u+y+z)}{\delta r} \\
 &\quad \times \frac{\delta \cos(r+s)}{\delta r} \times \frac{\delta(r+s)}{\delta r} + \frac{\delta(u+y+z)^2}{\delta u} \times \frac{\delta(u+y+z)}{\delta z} \\
 &\quad \times \frac{\delta \sin(r+s)}{\delta r} \times \frac{\delta(r+s)}{\delta r}
 \end{aligned}$$

$$= 2(u+y+z) \times 1 \times 1 + 2(u+y+z) \times 1 \times (-\sin(r+s)) \times 1 +$$

$$2(u+y+z) \times 1 \times \cos(r+s) \times 1$$

$$= 2(u+y+z) [1 + (-\sin(r+s)) + \cos(r+s)]$$

Putting the value of u, y, z

$$= 2(r-s + \cos(r+s) + \sin(r+s)) [1 + (-\sin(1-1)) + \cos(1-1)]$$

$$= 2(1+1+\cos(1-1)+\sin(1-1)) [1+(-\sin(0))+\cos(0)]$$

$$= 2(2+1+0)[1+0+1]$$

$$= 2 \times 3 [2] = 6 \times 2 = 12 \#.$$

$$\begin{aligned}
 \frac{\delta w}{\delta s} &= \frac{\delta w}{\delta u} \times \frac{\delta u}{\delta s} + \frac{\delta w}{\delta y} \times \frac{\delta y}{\delta s} + \frac{\delta w}{\delta z} \times \frac{\delta z}{\delta s} \\
 &= \frac{\delta(u+y+z)^2}{\delta(u+y+z)} \times \frac{\delta(u+y+z)}{\delta u} \times \frac{\delta(r-s)}{\delta s} \\
 &\quad + \frac{\delta(u+y+z)^2}{\delta(u+y+z)} \times \frac{\delta(u+y+z)}{\delta y} \times \frac{\delta(\cos(r+s))}{\delta(r+s)} \times \frac{\delta(r+s)}{\delta s} \\
 &\quad + \frac{\delta(u+y+z)^2}{\delta(u+y+z)} \times \frac{\delta(u+y+z)}{\delta z} \times \frac{\delta(\sin(r+s))}{\delta(r+s)} \times \frac{\delta(r+s)}{\delta s} \\
 &= 2(u+y+z) \times 1 \times (-1) + 2(u+y+z) \times 1 \times -\sin(r+s) \times 1 \\
 &\quad + 2(u+y+z) \times 1 \times \cos(r+s) \times 1 \\
 &= 2(u+y+z) [-1 + (-\sin(r+s) + \cos(r+s))] \\
 &\quad \text{when } r=1, s=-1 \\
 &= 2(1+1+1) [-1 + (0+1)] \\
 &= 2 \times 3 \times 0 \\
 &= 0 \# \\
 \therefore \frac{\delta w}{\delta s} &= 0 \quad \frac{\delta w}{\delta r} = 12 \#
 \end{aligned}$$

Implicit Differentiation:-

$f(u, y)$ is differentiable and eq^n is in the form $\frac{\partial}{\partial u} f(u, y) = 0$ and $f_y \neq 0$, then,

$$\frac{dy}{du} = -\frac{f_{yu}}{f_y}$$

* Find $\frac{dy}{du}$ if, $2u^2 - u^2 - \sin u y = 0$

→ Solution,

to find $\frac{dy}{du}$ we have to find f_u and f_y

$$\therefore f_u = \frac{\partial (u^2 - u^2 - \sin u y)}{\partial u}$$

$$= 0 - 2u - \frac{\partial \sin u y}{\partial u} \times \frac{\partial u y}{\partial u}$$

$$= -2u - y \cos u y$$

$$\therefore f_y = \frac{\partial (u^2 - u^2 - \sin u y)}{\partial y}$$

$$= 2u - 0 - \cos u y \times y$$

$$= 2u - y \cos u y$$

~~$$\therefore \frac{dy}{du} = -\frac{f_u}{f_y} = -\frac{(-2u - y \cos u y)}{2u - y \cos u y} = \frac{2u + y \cos u y}{2u - y \cos u y} \#$$~~

b) $y^3 + y^2 - 5y - u^2 + 4 = 0$

→ Solution,

$$f_u = \frac{\partial (y^3 + y^2 - 5y - u^2 + 4)}{\partial u} = -2u$$

$$f_y = \frac{\partial (y^3 + y^2 - 5y - u^2 + 4)}{\partial y} = 3y^2 + 2y - 5$$

$$\therefore \frac{dy}{du} = -\frac{f_u}{f_y} = \frac{-2u}{(3y^2 + 2y - 5)} \#$$

$$c) xy + y^2 - 3u - 3 = 0 \text{ at } (-1, 1)$$

→ Solution,

$$f_u = y + 0 - 3 = 0$$

$$f_y = u + 2y$$

$$\therefore \frac{dy}{du} = -\frac{u+3}{u+2y}$$

at $(-1, 1)$

$$\frac{dy}{du} = \frac{-1+3}{-1+2 \times 1} = \frac{2}{1} = 2 \neq$$

$$d) ue^y + \sin y + y - \ln 2 = 0 \text{ at } (0, \ln 2)$$

→ Solution

$$f_u = \frac{\partial (ue^y + \sin y + y - \ln 2)}{\partial u}$$

$$= e^y + \cos y \times y$$

$$= e^y + y \cos y$$

$$f_y = \frac{\partial (ue^y + \sin y + y - \ln 2)}{\partial y}$$

$$= ue^y + \cos y \times u + 1$$

$$= ue^y + u \cos y + 1$$

$$\therefore \frac{dy}{du} = -\frac{(e^y + y \cos y)}{ue^y + u \cos y + 1} \text{ at } (0, \ln 2)$$

$$= -\frac{(e^{\ln 2} + \ln 2 \cos(0 \times \ln 2))}{0 \times e^y + 0 \times \cos(0 \times \ln 2) + 1}$$

$$= -(2 + \ln 2)$$

$$= -2.69 \neq$$

Directional Derivative And Gradient Vector

Gradient Vector (∇f): ∇f at point $P(x, y)$

$$\begin{aligned}\nabla f &= \frac{\partial f}{\partial x} \vec{i} + \frac{\partial f}{\partial y} \vec{j} \\ &= f_x \vec{i} + f_y \vec{j}\end{aligned}$$

$$\therefore \nabla f \text{ (Gradient Vector)} = f_x \vec{i} + f_y \vec{j}$$

 ∇f at point $P(x, y, z)$ is,

$$\nabla f = f_x \vec{i} + f_y \vec{j} + f_z \vec{k}$$

* Find Gradient Vector (∇f).

$$a) f(x, y) = \ln(x^2 + y^2) \text{ at } (1, 1)$$

→ Solution,

$$f_x = \frac{\partial}{\partial x} \ln(x^2 + y^2) = \frac{2x}{x^2 + y^2}$$

$$= \frac{1}{(1^2 + 1^2)} \times 2 \cdot 1 \text{ at } (1, 1) = \frac{2}{2} = 1.$$

$$f_y = \frac{\partial}{\partial y} \ln(x^2 + y^2) = \frac{2y}{x^2 + y^2}$$

$$= \frac{1}{(1^2 + 1^2)} \times 2 \cdot 1 \text{ at } (1, 1) = \frac{2 \times 1}{2} = 1$$

$$\therefore \nabla f = f_x \vec{i} + f_y \vec{j} = \vec{i} + \vec{j}$$

$$b) f(x, y, z) = x^3 - xy^2 - z \text{ at } (1, 1, 0)$$

→ Solution,

$$f(u) = \frac{\partial(u^3 - uy^2 - z)}{\partial u} = 3u^2 - y^2 \text{ at } (1, 1, 0)$$

$$f(u) = 3 - 1 = 2$$

$$f(y) = \frac{\partial(u^3 - uy^2 - z)}{\partial y} = -2uy \text{ at } (1, 1, 0)$$

$$f(y) = -2 \times 1 \times 1 = -2$$

$$f(z) = \frac{\partial(u^3 - uy^2 - z)}{\partial z} = -1$$

$$\therefore \vec{V}f = 2\vec{i} - 2\vec{j} - \vec{k}$$

* Directional Derivatives:-

If $f(u, y)$ is differentiable at point $p(u, y)$ then it is denoted by $D_u f$ is given by

$$D_u f = \vec{V}f \cdot \vec{u} \text{ at that point } p(u, y).$$

where, $\vec{u} = \frac{\vec{v}}{|\vec{v}|}$ = unit vector

Directional derivative is a dot product of gradient vector and unit vector.

$$D_u f = \vec{V}f \cdot \vec{u} = |\vec{V}f| |\vec{u}| \cos \theta$$

$$= |\vec{V}f| \cos \theta$$

Properties:-

- $D_u f$ increase when $\cos \theta = 1$ i.e. $\theta = 0^\circ$
- $D_u f$ decrease when $\cos \theta = -1$, i.e. $\theta = 180^\circ$

* find the directional derivatives of $f(u, y) = u^2 \sin 2y$

at point $(1, \pi/2)$ in the direction of $v = 3\vec{i} - 4\vec{j}$.

→ Solution,

given vector $(\vec{v}) = 3\vec{i} - 4\vec{j}$

$$\text{Unit vector } (u) = \frac{v}{|v|} = \frac{3\vec{i} - 4\vec{j}}{\sqrt{9+16}} = \frac{3}{5}\vec{i} - \frac{4}{5}\vec{j}$$

given function $= f(u, y) = u^2 \sin 2y$

$$f(u) = \frac{\delta(u^2 \sin 2y)}{\delta u} = 2u \sin 2y \times \frac{\delta 2y}{\delta u}$$

at point $(1, \pi/2)$

$$f(u) = 2 \times 1 \sin 2 \times \frac{\pi}{2} = 0$$

$$f(y) = \frac{\delta(u^2 \sin 2y)}{\delta y} = u^2 \sin 2y (-\cos 2y) = -u^2 y \cos 2y$$

at point $(1, \pi/2)$

~~$$\frac{\delta(u^2 \sin 2y)}{\delta y} f(y) = -1^2 \cos 2 \cdot \frac{\pi}{2} = -1$$~~

~~$$\therefore \text{Gradient vector } (\vec{v}_f) = 0\vec{i} - \vec{j} = -\vec{j}$$~~

~~$$\text{Unit vector } (u) = \frac{3}{5}\vec{i} - \frac{4}{5}\vec{j}$$~~

∴ directional derivatives = $\vec{v}_f \cdot u$

$$= -\vec{j} \cdot \left(\frac{3}{5}\vec{i} - \frac{4}{5}\vec{j} \right)$$

$$= 0 + \frac{4}{5}$$

~~$$= \frac{4}{5}$$~~

(b) $f(u, y) = u \cdot e^y + \cos(uy)$ at $(2, 0)$ in direction of
 $v = 3\vec{i} - 4\vec{j}$

→ Solution,

given vector $(\vec{v}) = 3\vec{i} - 4\vec{j}$

$$\therefore \text{Unit Vector } (\mathbf{u}) = \frac{\vec{v}}{\|v\|} = \frac{3\vec{i} - 4\vec{j}}{\sqrt{9+16}} = \frac{3}{5}\vec{i} - \frac{4}{5}\vec{j}$$

$$\text{given function } = f(u, y) = u \cdot e^y + \cos(uy)$$

$$f(u) = \frac{\partial (u \cdot e^y + \cos(uy))}{\partial u}$$

$$= e^y + \frac{\partial \cos(uy)}{\partial u} \times \frac{\partial (uy)}{\partial u}$$

$$= e^y + (-\sin uy) \times y$$

at (2, 0)

$$= e^0 + (-\sin 0 \times 2) \times 0$$

$$f(u) = 1$$

$$f(y) = \frac{\partial (u \cdot e^y + \cos uy)}{\partial y}$$

$$= ue^y + (-\sin uy) \times u$$

at (2, 0)

$$f(y) = 2 \times e^0 + (-\sin 0 \times 2) \times 0$$

$$= 2$$

Now,

$$\text{gradient Vector } (\vec{\nabla}f) = \vec{i} + 2\vec{j}$$


$$\therefore \text{directional derivatives } (D_{\mathbf{u}}f) = \vec{\nabla}f \cdot \mathbf{u}$$

$$= (\vec{i} + 2\vec{j}) \cdot \left(\frac{3}{5}\vec{i} - \frac{4}{5}\vec{j} \right)$$

$$= \frac{3}{5} - \frac{8}{5}$$

$$= -\frac{5}{5} = -1 \quad \#.$$

$$* f(u, y, z) = -u^3 - ny^2 - z \text{ at } (1, 1, 0) \text{ in the direction of } \mathbf{v} = 2\vec{i} - 3\vec{j} + 6\vec{k}$$



→ Solution,

$$\text{Unit Vector } (\mathbf{u}) = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{2\mathbf{i} - 3\mathbf{j} + 6\mathbf{k}}{\sqrt{4+9+36}} = \frac{2}{7}\mathbf{i} - \frac{3}{7}\mathbf{j} + \frac{6}{7}\mathbf{k}$$

then,

$$f(u) = \frac{\delta(u^3 - uy^2 - z)}{\delta u} = 3u^2 - y^2 \text{ at } (1, 1, 0)$$

$$\therefore f(u) = 3 \times 1 - 1 = 2 \#.$$

$$f(y) = \frac{\delta(u^3 - uy^2 - z)}{\delta y} = -2uy \text{ at } (1, 1, 0)$$

$$\therefore f(y) = -2 \times 1 \times 1 = -2$$

$$f_z = \frac{\delta(u^3 - uy^2 - z)}{\delta z} = -1$$

$$\therefore \text{gradient vector } (\nabla f) = 2\mathbf{i} - 2\mathbf{j} - \mathbf{k}$$

$$\begin{aligned} \therefore \text{directional derivatives } (\text{D}_u f) &= \nabla f \cdot \mathbf{u} \\ &= (2\mathbf{i} - 2\mathbf{j} - \mathbf{k}) \cdot \left(\frac{2}{7}\mathbf{i} - \frac{3}{7}\mathbf{j} + \frac{6}{7}\mathbf{k} \right) \\ &= \frac{4}{7} + \frac{6}{7} - \frac{6}{7} \\ &= \frac{4}{7} \# \end{aligned}$$

* Equation of Tangent at points (u_1, y_1, z_1) :

when,

$$\nabla f = A\mathbf{i} + B\mathbf{j} + C\mathbf{k} \text{ at } (u_1, y_1, z_1)$$

$$\text{Eqn of tangent} = A(u - u_1) + B(y - y_1) + C(z - z_1)$$

* Equation of normal line at (u_1, y_1, z_1)
 eq^n are,

$$u = u_1 + At$$

$$y = y_1 + Bt$$

$$z = z_1 + Ct$$

* Find the eqⁿ of tangent and normal line of the Surface. $f(u, y, z) = u^2 + y^2 + z^2 - 9 = 0$ at point $P(1, 2, 4)$.

$$\nabla f = A\vec{i} + B\vec{j} + C\vec{k} \text{ at } (1, 2, 4)$$

→ Solution,

$$\text{given function} = f(u, y, z) = u^2 + y^2 + z^2 - 9 = 0$$

$$f_u = \frac{\delta(u^2 + y^2 + z^2 - 9)}{\delta u} = 2u \text{ at } (1, 2, 4)$$

$$f_u = 2 \times 1 = 2$$

$$f_y = \frac{\delta(u^2 + y^2 + z^2 - 9)}{\delta y} = 2y \text{ at } (1, 2, 4)$$

$$f_y = 2 \times 2 = 4$$

$$f_z = \frac{\delta(u^2 + y^2 + z^2 - 9)}{\delta z} = 1$$

$$\therefore \text{gradient Vector } (\nabla f) = 2\vec{i} + 4\vec{j} + \vec{k} \dots (1)$$

Comparing eqⁿ (1) with $A\vec{i} + B\vec{j} + C\vec{k}$

$$\therefore A = 2, B = 4, C = 1$$

Now, Point $(u_1, y_1, z_1) = (1, 2, 4)$

∴ Required eqⁿ of tangent is,

$$= A(u - u_1) + B(y - y_1) + C(z - z_1)$$

$$= 2(u - 1) + 4(y - 2) + 1(z - 4)$$

$$= 2u - 2 + 4y - 8 + z - 4$$

$$\therefore \text{eq}^n \text{ of tangent} = 2u + 4y + z - 14$$

$2u + 4y + z = 14$ } eqⁿ of tangent.

eqⁿ of normal are,

$$u = u_1 + At = 1 + 2t$$

$$\therefore u = 1 + 2t \#$$

$$y = y_1 + Bt$$

$$\therefore y = 2 + 4t \#$$

$$z = z_1 + Ct$$

$$\therefore z = 4 + t \#.$$

* Find the eqⁿ of tangent to surface $z = 1 - \frac{1}{10}(u^2 + 4y^2)$ at $(1, 1, 1/2)$.

→ Solution:- For eqⁿ of T to S = $A(u - u_1) + B(y - y_1) - (z - z_1)$

given function $= z = f(u, y) = 1 - \frac{1}{10}(u^2 + 4y^2)$

$$f_u = \frac{\partial}{\partial u} \left(1 - \frac{1}{10}(u^2 + 4y^2) \right)$$

$$= 0 - \frac{1}{10} \times 2u \quad \text{at } (1, 1)$$

$$f_u = -\frac{1}{5}$$

$$f_y = \frac{\partial}{\partial y} \left(1 - \frac{1}{10}(u^2 + 4y^2) \right)$$

$$= 0 - \frac{1}{10} (0 + 8y)$$

$$\text{at } (1, 1)$$

$$f_y = -\frac{1}{10} \times 8 \times 1 = -\frac{4}{5}$$

Plane tangent to surface $z = f(u, y)$

$$A(u-u_1) + B(y-y_1) - (z-z_1) = 0$$

classmate

Date _____

Page _____

$$f_z = \frac{\delta}{\delta z} \left(1 - \frac{1}{10}(u^2 + 4y^2) \right) = 0$$

Now,

$$\text{Vector gradient } (\vec{\nabla} f) = -\frac{1}{5}\vec{i} - \frac{4}{5}\vec{j}$$

Comparing $\vec{\nabla} f$ with $A\vec{i} + B\vec{j} + C\vec{k}$

$$A = -\frac{1}{5}, B = -\frac{4}{5}, C = 0$$

$$\therefore \text{eqn of tangent} = A(u-u_1) + B(y-y_1) - (z-z_1) = 0$$

$$a, -\frac{1}{5}(u-1) - \frac{4}{5}(y-1) - (z-\frac{1}{2}) = 0$$

$$a, -\frac{1}{5}u + \frac{1}{5} - \frac{4}{5}y + \frac{4}{5} - z + \frac{1}{2} = 0$$

$$a, -\frac{1}{5}u - \frac{4}{5}y - z + \frac{3}{2} = 0 \quad \#$$

$$\frac{1}{5}u + \frac{4}{5}y + z = \frac{3}{2}$$

Req. eqn

~~z~~

Extreme Values And Saddle Points.

1. First derivative test :-

If a function $f(x,y)$ is continuous and differentiable.

$$\begin{aligned} f_x &= 0 \\ f_y &= 0 \end{aligned} \quad \left. \begin{array}{l} \text{maximum or minimum.} \\ \text{maximum or minimum.} \end{array} \right\}$$

2. Second derivative test:

If a function $f(x,y)$ is continuous and its first and second derivative is differentiable at point (a,b) then;

- a) If $f_{xx} > 0$ and $f_{xx} \cdot f_{yy} - f_{xy}^2 > 0$
Then function is minimum.
- b) If $f_{xx} < 0$ and $f_{xx} \cdot f_{yy} - f_{xy}^2 > 0$, then
the function is maximum.
- c) If $f_{xx} \cdot f_{yy} - f_{xy}^2 < 0$, then there is saddle point.
- d) If $f_{xx} \cdot f_{yy} - f_{xy}^2 = 0$, then function is inconclusive.

* Find the extreme value of the following function :-

a) $f(x,y) = xy - x^2 - y^2 - 2x - 2y + 4$.

→ Solution :

The function is defined and differentiable for all x and y and its domain has no boundary points. The function therefore has extreme values only at the points

where f_u and f_y are simultaneously zero. This leads to,

$$f_u = \frac{\delta (uy - u^2 - y^2 - 2u - 2y + 4)}{\delta u} \text{ or,}$$

for Stationary point

$$\therefore f_u = y - 2u - 2 = 0 \quad \text{--- (i)}$$

$$f_u = y - 2u - 2$$

$$f_y = \frac{\delta (uy - u^2 - y^2 - 2u - 2y + 4)}{\delta y}$$

$$f_y = u - 2y - 2$$

$$f_y = u - 2y - 2 = 0 \quad \text{--- (ii)}$$

$$f_u = f_y = 0$$

$$\therefore f_u = y - 2u - 2 = 0$$

$$\therefore y = 2u + 2$$

Putting $y = 2u + 2$ in eq (ii)

$$u - 2(2u + 2) - 2 = 0$$

$$\therefore u - 4u - 4 - 2 = 0$$

$$\therefore -3u = 6$$

$$\therefore u = -2$$

$$\therefore y = 2 \times (-2) + 2 = \cancel{-4} - 2$$

$$\therefore (u, y) = (-2, +2)$$

Now,

$$f_{uu} = \frac{\delta (y - 2u - 2)}{\delta u} = -2$$

$$f_{yy} = \frac{\delta (y - 2u - 2)}{\delta y} = -2$$

$$f_{uy} = \frac{\delta (y - 2u - 2)}{\delta y} = 1$$

$$\therefore f_{uu} = -2 < 0$$

$$f_{uu} \cdot f_{yy} - f_{uy}^2 = (-2) \times (-2) - (-1)^2 = 4 - 1 = 3 > 0$$

Thus the function is maximum at $(-2, -2)$. The value of f at $f(-2, -2)$ is

$$\begin{aligned} f(-2, -2) &= (-2) \times (-2) - (-2)^2 - (-2)^2 - 2 \times (-2) - 2 \times (-2) + 4 \\ &= 4 - 4 - 4 + 4 + 4 + 4 \\ &= 8 \end{aligned}$$

b) $f(u, y) = 2uy - 5u^2 - 2y^2 + 4u + 4y - 4$
 → Solution :

The function is defined and differentiable for all u and y and its domain has no boundary point. The function therefore has extreme values only at the point where f_u and f_y are simultaneously 0. This leads

$$f_u = 2y - 10u + 4 = 0 \quad \text{--- (i)}$$

$$f_y = 2u - 4y + 4 = 0 \quad \text{--- (ii)}$$

$$2y = 10u - 4$$

$$\therefore y = 5u - 2 \quad \text{--- (iii)}$$

put $y = 5u - 2$ in eqn (ii)

$$2u - 4(5u - 2) + 4 = 0$$

$$\therefore 2u - 20u + 8 + 4 = 0$$

$$\therefore -18u = -12$$

$$\therefore u = \frac{12}{18} = \frac{2}{3}$$

Put $u = \frac{2}{3}$ in eqn (iii)

$$y = 5 \times \frac{2}{3} - 2 = \frac{10}{3} - 2 = \frac{4}{3}$$

$$\therefore (u, y) = \left(\frac{2}{3}, \frac{4}{3}\right)$$

Now,

$$f_{uu} = -10$$

$$f_{yy} = -4$$

$$f_{uy} = 2$$

Here,

$$f_{uu} = -10 < 0$$

and,

$$\begin{aligned} f_{uu} \cdot f_{yy} - f_{uy}^2 &= -10 \times -4 - (2)^2 \\ &= 40 - 4 \\ &= 36 > 0 \end{aligned}$$

So the function f has local maximum at $(\frac{2}{3}, \frac{4}{3})$ and its value is,

$$\begin{aligned} f\left(\frac{2}{3}, \frac{4}{3}\right) &= 2 \times \frac{2}{3} \times \frac{4}{3} - 5 \times \left(\frac{2}{3}\right)^2 - 2 \times \left(\frac{4}{3}\right)^2 + 4 \times \frac{2}{3} + 4 \times \frac{4}{3} - 5 \\ &= \frac{16}{9} - \frac{20}{9} - \frac{32}{9} + \frac{8}{3} \times \frac{3}{3} + \frac{16}{3} + \frac{3}{3} - \frac{4}{1} \times \frac{4}{3} \\ &= \frac{16 - 20 - 32 + 24 + 48 - 36}{9} \\ &= \frac{88 - 88}{9} = 0 \end{aligned}$$

∴ The value of f at $f\left(\frac{2}{3}, \frac{4}{3}\right)$ is 0.

b) $f(u, y) = 5uy - 7u^2 + 3u - 6y + 2$

→ Solution,

The function is defined and differentiable for all u and y . and its domain has no boundary point. The function therefore has the extreme value only at the point where f_u and f_y are simultaneously zero (0). This leads,

$$f_u = 5y - 14u + 3 = 0 \quad (i)$$

$$f_y = 5u - 6 = 0 \quad (ii) \quad \therefore u = \frac{6}{5}$$

$$5y - 14 \times \frac{6}{5} + 3 = 0$$

$$\therefore 5y = \frac{84}{5} - 3$$

$$\therefore 5y = \frac{69}{5}$$

$$\therefore y = \frac{69}{25}$$

$$f_{uu} = 8(5y - 14u + 3) = -14$$

$$f_{yy} = \frac{8(15u - 6)}{8y} = 0$$

$$f_{uy} = \frac{8(5y - 14u + 3)}{8y} = 5$$

$$f_{uu} = -14 < 0$$

$$f_{uu} \cdot f_{yy} - f_{uy}^2 = -14 \times 0 - 5^2 = 0 - 25 = -25 < 0$$

Since $f_{uu} \cdot f_{yy} - f_{uy}^2 = -25 < 0$, so the function f_{uy} has saddle point at $(\frac{6}{5}, \frac{69}{25})$

$$\Rightarrow f(u, y) = u^2 + ny + 3u + 20y + 5$$

→ Solution,

Given function is defined and differentiable for all u and y and its domain has no boundary point. The function therefore has extreme value only at point where f_u and f_y are simultaneously 0. This leads.

$$f_u = \frac{\partial f}{\partial u} = 8(u^2 + uy + 3u + 2y + 5)$$

$$f_u = 2u + y + 3 = 0$$

$$f_y = \frac{\partial f}{\partial y} = 5(u^2 + uy + 3u + 2y + 5)$$

$$f_y = u + 2 = 0 \quad \therefore u = -2$$

then,

$$2u + y + 3 = 0$$

$$\therefore 2(-2) + y + 3 = 0$$

$$\therefore -4 + y + 3 = 0$$

$$\therefore y = 1 \quad \therefore (u, y) = (-2, 1)$$

Now,

$$f_{uu} = \frac{\partial^2 f}{\partial u^2} = 2$$

$$f_{yy} = \frac{\partial^2 f}{\partial y^2} = 0$$

$$f_{uy} = \frac{\partial^2 f}{\partial u \partial y} = 1$$

Now,

$$\therefore f_{uu} = 2 > 0$$

$$f_{uu} \cdot f_{yy} - f_{uy}^2 = 2 \times 0 - 1 = -1$$


$$\therefore -1 < 0$$

\therefore The function has Saddle point at $(-2, 1)$.

$$\text{d} \Rightarrow f(u, y) = u^2 + uy + y^2 + 3u - 3y + 4$$

\rightarrow Solution,

given function is defined and differentiable for all u and y and its domain has no boundary point. The function therefore has extreme value only at point where f_u and f_y are simultaneously 0.



then,

$$f_{uu} = \frac{\partial^2 f}{\partial u^2} = 2(u^2 + 2u + 3) = 2(u^2 + 2u + 1 + 2) = 2(u+1)^2 + 4$$

$$\therefore f_{uu} = 2u + 2 + 3 = 2u + 5 = 0$$

$$f_{uy} = \frac{\partial^2 f}{\partial u \partial y} = u + 2y - 3$$

$$\therefore f_{uy} = u + 2y - 3 = 0$$

$$\therefore u = 3 - 2y$$

Now,

$$2u + 2y + 3 = 0$$

$$\therefore 2(3 - 2y) + 2y + 3 = 0$$

$$\text{or, } 6 - 4y + 2y + 3 = 0 \quad \text{or, } 6 - 2y + 3 = 0$$

$$\therefore -2y + 9 = 0$$

$$\therefore 2y = 9$$

$$\therefore y = 3$$

Put $y = 3$ in $u = 3 - 2y$

$$\therefore u = 3 - 2 \times 3 = 3 - 6 = -3$$

$$\therefore (u, y) = (-3, 3)$$

we know,

$$f_{uu} = \frac{\partial^2 f}{\partial u^2} = 2$$

$$f_{yy} = \frac{\partial^2 f}{\partial y^2} = 2$$

$$f_{uy} = \frac{\partial^2 f}{\partial u \partial y} = 1$$

$$\therefore f_{uy} = 1$$

Now,

$$f_{uu} = 2 > 0$$

$$f_{uu} \cdot f_{yy} - f_{uy}^2 = 2 \times 2 - 1 = 3 > 0$$

so the function f has local minimum at $(-3, 3)$ and its value at $f(-3, 3)$ is

$$f(-3, 3) = (-3)^2 + 2(-3) + 2 + 3(-3)^2 + 4 = 9 + 2(-3) + 2 + 27 + 4 = 38$$

$$f(-3, 3) = 4 + 4 + 4 + 4 = 16 \quad \text{not } -13$$

e) $f(u, y) = u^3 + y^3 + 3u^2 - 3y^2 - 8$

→ Solution,

given function is defined and differentiable for all u and y and its domain has no boundary point. The function therefore has extreme value only at point where f_u and f_y ^{are} simultaneously zero. This leads,

$$f_u = \frac{\partial (u^3 + y^3 + 3u^2 - 3y^2 - 8)}{\partial u}$$

$$= 3u^2 + 6u = 0$$

$$f_y = \frac{\partial (u^3 + y^3 + 3u^2 - 3y^2 - 8)}{\partial y}$$

$$= 3y^2 - 6y = 0$$

$$f_u = 3u^2 + 6u = 0 \quad \dots \textcircled{1}$$

$$f_y = 3y^2 - 6y = 0 \quad \dots \textcircled{11}$$

$$3u^2 + 6u = 0$$

$$\text{or, } 3u(u+2) = 0$$

$$\text{Either } u = 0, -2$$

$$3y^2 - 6y = 0$$

$$\text{or, } 3y(y-2) = 0$$

$$\text{with } y = 0, y = 2$$

$$\therefore u = (0, -2) \quad (y = 0, 2)$$

∴ The points are $(0, 0), (0, 2), (-2, 0), (-2, 2)$

Now,

$$f_{uu} = 6u + 6 \quad f_{uy} = 0$$

$$f_{yy} = 6y - 6$$

at point $(0, 0)$

$$f_{uu} \cdot f_{yy} - f_{uy}^2 = 6 \times (-6) - 0 = -36 < 0$$

$$f_{uu} = 6 > 0$$

@

then the function has saddle point at $(0, 0)$.

(b) at point $(0, 2)$

$$f_{xx} = 6x + 6 = 6 \times 0 + 6 = 6 \quad f_{yy} = 0$$

$$f_{yy} = 6y - 6 = 6 \times 2 - 6 = 12 - 6 = 6$$

$$f_{xx} = 6 > 0$$

$$f_{xx} \cdot f_{yy} - f_{xy}^2 = 6 \times 6 - 0 = 36 > 0$$

Hence the function have local minimum at $(0, 2)$ and its value is

$$\begin{aligned} f(0, 2) &= 0^3 + 2^3 + 3 \times 0^2 - 3 \times (2)^2 - 8 \\ &= 8 - 12 - 8 \\ &= -12 \# \end{aligned}$$

(c) at point $(-2, 0)$

$$f_{xx} = 6 \times (-2) + 6 = -12 + 6 = -6$$

$$f_{yy} = 6 \times 0 - 6 = -6$$

$$f_{xy} = 0$$

$$f_{xx} = -6 < 0$$

$$f_{xx} \cdot f_{yy} - f_{xy}^2 = -6 \times -6 - 0 = +36 > 0$$

Hence the function have local maximum at

$(-2, 0)$ and its value is,

$$\begin{aligned} f(-2, 0) &= (-2)^3 + (0)^3 + 3(-2)^2 - 3 \times 0^2 - 8 \\ &= -8 + 12 - 8 \\ &= -4 \# \end{aligned}$$

(d) at point $(2, 2)$

$$f_{xx} = 6 \times 2 + 6 = -6, \quad f_{yy} = 6 \times 2 - 6 = -6 \quad f_{xy} = 0$$

$$f_{xx} < 0 \quad f_{xx} = 18 > 0, \quad f_{xx} \cdot f_{yy} - f_{xy}^2 = 18 \times -6 - 0 = -108 < 0$$

Hence the function have local minimum at point

$(2, 2)$ and its value at $f(2, 2)$ is,

$$f(2, 2) = 2^3 + 2^3 + 3 \cdot 2^2 - 3 \cdot 2^2 - 8 = 8 + 8 - 8 = 8 \#$$

saddle point at $(2, 2)$

Absolute Maximum And Minimum"

STEPS:

1. List the interior points of the region R , where function may have local maxima and minima and evaluate function at this point. These points are critical /stationary point.
2. List the boundary point of the Region R , where the function have local maxima and local minima and evaluate function at this point.
3. Look for maximum and minimum values of function.

* Find the absolute maximum and minimum values of $f(u, y) = 2 + 2u + 2y - u^2 - y^2$ on the triangular region in the first quadrant bounded by line $u=0$, $y=0$ and $y=9-u$.

→ solution,


Solving eqⁿ $y = g - u$

when $n=0$

$$y = 9$$

when $y = 0$

$$v = 9$$



→ Solution :-

Given function,

$$f(u, y) = 2 + 2u + 2y - u^2 - y^2$$

$$\frac{\partial f}{\partial u} = \cancel{2} (2 + 2u + 2y - u^2 - y^2) = 2 - 2u = 0$$

$$\therefore 2u = 2$$

$$\therefore u = 1$$

$$\frac{\partial f}{\partial y} = \cancel{2} (2 + 2u + 2y - u^2 - y^2) = 2 - 2y = 0$$

$$\therefore y = 1$$

$$\therefore 2 - 2y = 0 \quad \therefore y = 1$$

$$\therefore f(1, 1) = 2 + 2 \times 1 + 2 \times 1 - 1 - 1 \rightarrow \text{In given function}$$

$$= 2 + 2 + 2 - 2$$

$$= 4$$

Boundary points

a) line segment OA, $y=0$

$$y=0$$

$$f(u, 0) = 2 + 2u + 2 \times 0 - u^2 - 0 = 2u - u^2 + 2$$

$$\therefore f(u) = 2 - 2u = 0$$

$$\therefore 2u = 2 \quad \text{so } u = 1, y = 0$$

Critical point is $(1, 0)$

$$f(1, 0) = 2 + 2 \times 1 + 2 \times 0 - 1^2 - 0^2$$

$$= 2 + 2 - 1 = 3$$

At point A(9, 0)


$$f(9, 0) = 2 + 2 \times 9 + 2 \times 0 - 9^2 - 0^2 = 2 + 18 - 81 = -61$$

At point O(0, 0)

$$f(0, 0) = 2 + 2 \times 0 + 2 \times 0 - 0^2 - 0^2 = 2$$

b) line segment OB, $u=0$

$$u=0$$



$$f(0, y) = 2 + 2 \times 0 + 2y - 0^2 - y^2 = 2y - y^2 + 2$$

$$fy = 2y - 2y = 0$$

$$2y - 2y \therefore y = 1$$

critical point $(0, 1)$

$$\therefore f(0, 1) = 2 + 2 \times 0 + 2 \times 1 - 0^2 - 1^2 = 2 + 2 - 1 = 3$$

✓ at point $B(0, 9)$

$$f(0, 9) = 2 + 2 \times 0 + 2 \times 9 - 0^2 - 9^2 = 2 + 18 - 81 = -61$$

⇒ line segment AB , $y = 9 - u$

$$\begin{aligned} f(u, 9-u) &= 2 + 2u + 2(9-u) - u^2 - (9-u)^2 \\ &= 2 + 2u + 18 - 2u - u^2 - (81 - 2 \cdot 9 \cdot u + u^2) \\ &= 20 + 2u - 2u - u^2 - 81 + 18u - u^2 \\ &= -61 - 2u^2 + 18u \end{aligned}$$

$$\therefore f(u) = -4u + 18$$

$$f(u) = 0$$

$$\therefore 4u = 18 \therefore u = \frac{18}{4} = \frac{9}{2}$$

Put $u = \frac{9}{2}$ in eqⁿ $y = 9 - u$

$$\text{critical point} = \left(\frac{9}{2}, \frac{9}{2}\right) \therefore y = 9 - \frac{9}{2} = \frac{9}{2}$$

$$\therefore f\left(\frac{9}{2}, \frac{9}{2}\right) = 2 + 2 \times \frac{9}{2} + 2 \times \frac{9}{2} - \left(\frac{9}{2}\right)^2 - \left(\frac{9}{2}\right)^2$$

$$= 2 + \frac{18}{2} + \frac{18}{2} - \frac{81}{4} - \frac{81}{4} = \frac{8 + 36 + 36 - 81 - 81}{4}$$

$$= -\frac{41}{2} = -20.5$$

AF/BS/1

$$\therefore f(1,1) = 4, f(1,0) = 3, f(0,1) = 3, f(9,9) = -61$$

at A $f(9,0) = -61$ -20.5

at B $f(0,9) = -61$

at O $f(0,0) = 2$

The absolute maximum is 4 at (1,1) and absolute minimum is -61 at (0,9) and (9,0).

* Find absolute max. and min. values of $f(u,y) = 2u^2 - 4u + y^2 - 4y + 1$ on closed triangular plate bounded by line $u=0, y=2, y=2u$ in first quadrant.

$$\rightarrow \text{eqn } y=2u$$

when

$$u=0 \quad y=0$$

$$u=1 \quad y=2$$

$$u=2 \quad y=4$$

\rightarrow Solution,

given function,

$$f(u,y) = 2u^2 - 4u + y^2 - 4y + 1$$

here,

$$f_u = \frac{\partial}{\partial u} (2u^2 - 4u + y^2 - 4y + 1) = 4u - 4$$

$$\therefore f_u = 0$$


$$4u - 4 = 0$$

$$\therefore u = 1$$

$$f_y = 2y - 4 = 0$$

$$2y = 4$$

$$\therefore y = 2$$



$$\therefore f(1,2) = 2 \times 1^2 - 4 \times 1 + 2^2 - 4 \times 2 + 1$$

$$= 2 - 4 + 4 - 8 + 1$$

$$= -5 \quad \text{--- (1)}$$

boundary points

a) line segment OA, $y = 2u$

$$y = 2u$$

$$\begin{aligned} f(u, 2u) &= 2u^2 - 4u + (2u)^2 - 4(2u) + 1 \\ &= 2u^2 - 4u + 4u^2 - 8u + 1 \\ &= 6u^2 - 12u + 1 \end{aligned}$$

$$f(u) = \frac{6(6u^2 - 12u + 1)}{6u}$$

$$= 12u - 12$$

$$f_u = 0$$

$$12u - 12 = 0$$

$$\therefore u = 1$$

Put $u = 1$ in $y = 2u$

$$\therefore y = 2 \times 1 = 2$$

\therefore (critical) point is $(1, 2)$

$$\textcircled{2} - f(1, 2) = 2 \times 1^2 - 4 \times 1 + 2^2 - 4 \times 2 + 1 = 2 - 4 + 4 - 8 + 1 = -5$$

at point $O(0, 0)$

$$\textcircled{3} - f(0, 0) = 2 \times 0^2 - 4 \times 0 + 0^2 - 4 \times 0 + 1 = 1$$

at point $A(1, 2)$

$$\textcircled{4} - f(1, 2) = -5$$

b) line segment $OB, u = 0$

$$u = 0$$

$$f(0, y) = 2 \times 0^2 - 4 \times 0 + y^2 - 4y + 1 = y^2 - 4y + 1$$

$$fy = 2y - 4 = 0$$

$$2y = 4$$

$$\therefore y = 2$$

$$\textcircled{5} - f(0, 2) = 2 \times 0^2 - 4 \times 0 + 2^2 - 4 \times 2 + 1 = 4 - 8 + 1 = -3$$

⑥- at point B(0, 2)

$$f(0, 2) = -3$$

c.7 line segment AB $y=2$

$$\cancel{f(0, 2)} \quad f(u, y) = f(u, 2) = 2u^2 - 4u + 2^2 - 4 \times 2 + 1$$

$$f(u, 2) = 2u^2 - 4u + 4 - 8 + 1 = 2u^2 - 4u - 3$$

$$f_u = 4u - 4 = 0$$

$$\therefore u = 1$$

put

⑦ $\therefore f(1, 2) = -5$

Hence,

$$f(1, 2) = -5$$

$$f(0, 0) = 1$$

$$f(0, 2) = -3$$

Hence absolute max. is 1 at (0, 0) and absolute minimum is -5 at (1, 2).


Q. Find absolute maximum and minimum for $f(u, y) = u^2 + my + y^2 - 6u + 2$ on rectangular plate $0 \leq u \leq 5, -3 \leq y \leq 0, m=0, u=5, y=-3, y=0$

→ Solution,

points are $u=0, 5$

$y= -3, 0$

∴ Rectangular points are $(0, -3), (0, 0), (5, -3), (5, 0)$



→ Solution,

given function,

$$f(u, y) = u^2 + uy + y^2 - 6u + 2$$

here,

$$f_u = 2u + y - 6$$

$$f_y = u + 2y$$

$$\text{since, } f_u = f_y = 0$$

$$2u + y = 6, \quad u = -2y$$

Now,

$$2(-2y) + y = 6$$

$$\therefore -4y + y = 6$$

$$\therefore -3y = 6$$

$$\therefore y = -\frac{6}{3} = -2$$

$$\text{Put } y \text{ in } u = -2y \quad \therefore u = -2 \times -2 = 4$$

$$\therefore \text{critical point} = (4, -2)$$

$$\begin{aligned} f(4, -2) &= (-4)^2 + (4)(-2) + (-2)^2 - 6 \times (4) + 2 \\ &= 16 - 8 + 4 - 24 + 2 = 8 - 24 = -16 \end{aligned}$$

Boundary points,

a) at line segment OA

at point O(0, 0)

$$f(0, 0) = 0^2 + 0 \times 0 + 0^2 - 6 \times 0 + 2 = 2$$

at point A (0, -3)

$$f(0, -3) = 0^2 + 0 \times -3 + (-3)^2 - 6 \times 0 + 2 = 9 + 2 = 11$$

at point B(5, -3)

$$\begin{aligned} f(5, -3) &= 5^2 + 5 \times (-3) + (-3)^2 - 6 \times 5 + 2 \\ &= 25 - 15 + 9 - 30 + 2 = -9 \end{aligned}$$

at point C (5, 0)

$$\begin{aligned} f(5, 0) &= 5^2 + 5 \times 0 + 0^2 - 6 \times 5 + 2 \\ &= 25 - 30 + 2 = -3 \end{aligned}$$

b) at line segment OA

$$u = 0$$

$$f(0, y) = 0^2 + 0 \times y + y^2 - 6 \times 0 + 2 = y^2 = y^2$$

$$\therefore fy = 2y = 0$$

$$\therefore y = 0 \quad \therefore \text{critical point} = (0, 0)$$

$$f(0, 0) = 0^2 + 0 \times 0 + 0^2 - 6 \times 0 + 2 = 2$$

b) at line segment AB, $y = -3$

$$f(u, -3) = u^2 + uy + y^2 - 6u + 2$$

$$= u^2 + u \times (-3) + (-3)^2 - 6u + 2$$

$$= u^2 - 3u + 9 - 6u + 2$$

$$= u^2 - 9u + 11$$

$$\therefore f(u) = 2u - 9 = 0$$

$$\therefore u = \frac{9}{2}$$

$$f\left(\frac{9}{2}, -3\right) = \left(\frac{9}{2}\right)^2 + \frac{9}{2} \times (-3) + (-3)^2 - 6 \times \frac{9}{2} + 2$$

$$f\left(\frac{9}{2}, -3\right) = \frac{81}{4} - \frac{27}{2} + 9 - \frac{54}{2} + 2 = -\frac{37}{4}$$

c) at line segment BC,

$$u=5$$

$$\begin{aligned} \therefore f(5, y) &= 25 + 5y + y^2 - 6 \times 5 + 2 \\ &= y^2 + 5y - 3 \end{aligned}$$

$$\therefore fy = 2y + 5 = 0$$

$$\therefore 2y = -5$$

$$y = -\frac{5}{2}$$

$$\therefore \text{critical point} = (5, -\frac{5}{2})$$

$$\begin{aligned} \therefore f(5, -\frac{5}{2}) &= (5)^2 + 5 \times (-\frac{5}{2}) + (-\frac{5}{2})^2 - 6 \times 5 + 2 \\ &= 25 - \frac{25}{2} + \frac{25}{4} - 30 + 2 \\ &= -\frac{37}{4} \end{aligned}$$

$$\therefore f(5, -\frac{5}{2}) = -\frac{37}{4}$$

d) at line segment OC, $y=0$

$$f(u, 0) = u^2 + ux0 + 0^2 - 6u + 2 = u^2 - 6u + 2$$

$$\therefore fu = 2u - 6 = 0$$

$$\therefore u = \frac{6}{2} = 3$$

$$\therefore f(3, 0) = 3^2 * 3 \times 0 + 0^2 - 6 \times 3 + 2 = 9 - 18 + 2 = -7$$

\therefore Hence absolute maximum is 12 at $(0, -3)$,
and absolute minimum is -10 at $(4, -2)$.

Linearization :-

$$L(u, y) = f(u_1, y_1) + f_u(u_1, y_1)(u - u_1) + f_y(u_1, y_1)(y - y_1)$$

$$\therefore f(u, y) \approx L(u, y)$$

* Find the linearization of :

q) $f(u, y) = u^2 - ny + \frac{1}{2} y^2 + 3$ at point $(3, 2)$.
 → Solution :-

given function $f(u, y) = u^2 - ny + \frac{1}{2} y^2 + 3$

and point $(u_1, y_1) = (3, 2)$

Now,

$$f_u = \frac{\partial}{\partial u}(u^2 - ny + \frac{1}{2} y^2 + 3) = 2u - y$$

at $(3, 2)$

$$f_u = 2 \times 3 - 2 = 6 - 2 = 4$$

$$f_y = \frac{\partial}{\partial y}(u^2 - ny + \frac{1}{2} y^2 + 3) = -u + \frac{1}{2} \times 2y$$

$$f_y = -u + y \text{ at } (3, 2)$$

$$f_y = -3 + 2 = -1$$

$$\begin{aligned}
 f(3, 2) &= f(u_1, y_1) = 3^2 - 3 \times 2 + \frac{1}{2} \times 2^2 + 3 \\
 &= 3^2 - 3 \times 2 + \frac{1}{2} \times 2^2 + 3 = 9 - 6 + \cancel{\frac{2}{2} \times 2} + 3 \\
 &= 9 - 6 + 2 + 3 = \cancel{18 - 12} + \cancel{2} + 3 \\
 &= 9 - 1 = 8 = \cancel{18 - 12} + 2 + 3
 \end{aligned}$$

$$L(3, 2) = f(u_1, y_1) + f_u(u_1, y_1)(u - u_1) + f_y(u_1, y_1)(y - y_1)$$

$$= 8 + 4(u - 3) + (-1)(y - 2)$$

$$= 8 + 4u - 12 - y + 2$$

$$\therefore L(3, 2) = 4u - y - 2 = 0$$

Method of Lagrange Multiplier:

Suppose that $f(u, y, z)$ and $g(u, y, z)$ are differentiable and to find the local maximum and local minimum values of the function subject to constraints $g(u, y, z) = 0$, we need to find the value of u, y, z and λ that simultaneously satisfy the equation,

$$\vec{\nabla}f = \lambda \vec{\nabla}g \quad \text{--- (i), and}$$

$$g(u, y, z) = 0 \quad \text{--- (ii)}$$

where

λ = Lagrange Multiplier

$$\vec{\nabla}f = f_u \vec{i} + f_y \vec{j} + f_z \vec{k}$$

$$\vec{\nabla}g = g_u \vec{i} + g_y \vec{j} + g_z \vec{k}$$

* Find the greatest and smallest values that the function $f(u, y) = uy$ takes on the ellipse $\frac{u^2}{8} + \frac{y^2}{2} - 1 = 0$

→ Solution,

given function, $f(u, y) = uy$

Subject to constraint $g(u, y) = \frac{u^2}{8} + \frac{y^2}{2} - 1 = 0$

Then, Vector gradient $\vec{\nabla}f$ of $f(u, y)$

$$f_u = y \quad f_y = u$$

$$\therefore \vec{\nabla}f = y \vec{i} + u \vec{j}$$

Then, Vector gradient $\vec{\nabla}f$ of $g(u, y)$

$$gu = \frac{\delta}{\delta u} \left(\frac{u^2}{8} + \frac{y^2}{2} - 1 \right)$$

$$= \frac{2u}{8} = \frac{u}{4}$$

$$gy = \frac{\delta}{\delta y} \left(\frac{u^2}{8} + \frac{y^2}{2} - 1 \right)$$

$$= \frac{2y}{2} = y$$

$$\therefore \bar{V}g = gu\vec{i} + gy\vec{j}$$

$$\bar{V}g = \frac{u}{4}\vec{i} + y\vec{j}$$

Now,

$$\bar{V}f = \lambda \bar{V}g$$

$$y\vec{i} + u\vec{j} = \lambda \left(\frac{u}{4}\vec{i} + y\vec{j} \right)$$

Equating both sides we get,

$$y = \frac{\lambda u}{4} \quad \text{and} \quad u = \lambda y$$

$$\therefore y = \frac{\lambda \cdot \lambda y}{4}$$

$$\therefore 4y - \lambda^2 y = 0$$

$$\therefore y(4 - \lambda^2) = 0$$

$$\therefore y = 0$$

$$\lambda = \pm 2$$

then,

$$g(u, y) = 0$$

$$\frac{u^2}{8} + \frac{y^2}{2} - 1 = 0$$

when $y = 0$,

$$\frac{u^2}{8} + \frac{0^2}{2} = 1 \quad u^2 = 8 \quad \therefore u = \pm\sqrt{8}$$

$$\therefore u = \pm 2\sqrt{2} \quad \text{why?}$$

$$f(2\sqrt{2}, 0) = f(u, y) = 0$$

$$f(-2\sqrt{2}, 0) = f(u, y) = 0$$

$$\therefore y = 0, u = \pm 2\sqrt{2}, \lambda = \pm 2$$

$u = \lambda y$ ✓ by equating $\lambda = \pm 2 \rightarrow$ by song.

$$\text{Put } u = \pm 2y \text{ in } g(u, y)$$

Now

$$\frac{u^2}{8} + \frac{y^2}{2} = 1$$

$$\text{or, } \frac{(\pm 2y)^2}{8} + \frac{y^2}{2} = 1$$

$$\text{or, } \frac{4y^2}{8} + \frac{y^2}{2} = 1$$

$$\text{or, } 2y^2 = 2$$

$$\therefore y = \pm 1, u = \pm 2y$$

when $y = +1$

$$u = \pm 2 \times (+1) = \pm 2$$

$$\therefore (+2, +1) (-2, +1)$$

$$f(2, 1) = 2 \times 1 = 2$$

$$f(-2, 1) = -2 \times 1 = -2$$

when $y = -1$

$$u = \pm 2 \times (-1) = \pm 2$$

$$\therefore (2, -1) (-2, -1)$$

$$f(2, 1) = -1 \times 2 = -2$$

$$f(-2, -1) = 1 \times -2 = -2$$

∴ The greatest value is 2 at $(-2, -2)$ and $(2, 1)$
and the smallest value is -2 at $(-2, 1)$ and $(2, -2)$.

* Find the maximum and minimum values of function $f(u, y) = 3u + 4y$ on circle $u^2 + y^2 = 1$.

→ Solution,

we have to find the value of u, y and λ which satisfy the condition,

$$\vec{V}f = \lambda \vec{V}g \text{ and } g(u, y) = 0$$

Now,

$$fu = 3, fy = 4$$

$$\therefore \vec{\lambda}f = 3\vec{i} + 4\vec{j}$$

$$gu = 2u, gy = 2y$$

$$\therefore \vec{V}g = 2u\vec{i} + 2y\vec{j}$$

$$\therefore 3\vec{i} + 4\vec{j} = (2u\vec{i} + 2y\vec{j})\lambda$$

$$3\vec{i} + 4\vec{j} = 2u\lambda\vec{i} + 2y\lambda\vec{j}$$

Equating on both sides

we get,

$$2u\lambda = 3, 4 = 2y\lambda$$

$$\therefore u = \frac{3}{2\lambda}, \therefore y = \frac{4}{2\lambda}$$

Substituting these values in $g(u, y) = 0$, we get

$$u^2 + y^2 = 1$$

$$\left(\frac{3}{2\lambda}\right)^2 + \left(\frac{4}{2\lambda}\right)^2 = 1$$

$$\therefore \frac{9}{4\lambda^2} + \frac{16}{4\lambda^2} = 1$$

$$\therefore 9 + 16 = 4\lambda^2$$

$$25 = 4\lambda^2$$

$$\therefore \lambda^2 = \frac{25}{4}$$

$$\therefore \lambda = \sqrt{\frac{25}{4}} = \pm \frac{5}{2}$$

Since $u = \frac{3}{2\lambda}$ and $y = \frac{4}{2\lambda}$, u and y have

different signs at $\lambda = 5/2$

$$\text{For } u = \frac{3}{2\lambda} \quad \therefore u = \frac{3}{2 \times \frac{5}{2}} = \frac{3}{5}$$

$$\text{at } \lambda = -5/2$$

$$u = \frac{3}{2 \times -\frac{5}{2}} = -\frac{3}{5}$$

$$\therefore u = \left(\frac{3}{5}, -\frac{3}{5} \right)$$

$$\text{For } y = \frac{4}{2\lambda}, \quad y = \frac{4}{2 \times \frac{5}{2}} = \frac{4}{5} \text{ at } \lambda = 5/2$$

$$\text{at } \lambda = -5/2$$

$$u(1, 2)$$

$$y(4, 5)$$

$$y = \frac{4}{2 \times -\frac{5}{2}} = -\frac{4}{5}$$

then

point are $(1, 4)$, $(1, 5)$, $(2, 4)$, $(2, 5)$

$$\therefore y = \left(\frac{4}{5}, -\frac{4}{5} \right)$$

as same

then points are $(\frac{3}{5}, \frac{4}{5})$, $(\frac{3}{5}, -\frac{4}{5})$, $(-\frac{3}{5}, \frac{4}{5})$

$$(-\frac{3}{5}, -\frac{4}{5})$$

$$u + f_{xy} = 3u + 4y$$

$$\therefore f\left(\frac{3}{5}, \frac{4}{5}\right) = 3 \times \frac{9}{25} + 4 \times \frac{4}{5} = \frac{9}{5} + \frac{16}{5} = \frac{25}{5} = 5$$

$$\therefore f\left(\frac{3}{5}, -\frac{4}{5}\right) = 3 \times \frac{3}{5} + 4 \times \left(-\frac{4}{5}\right) = \frac{9}{5} - \frac{16}{5} = -\frac{7}{5}$$

$$\therefore f\left(-\frac{3}{5}, \frac{4}{5}\right) = 3 \times \left(-\frac{3}{5}\right) + 4 \times \frac{4}{5} = -\frac{9}{5} + \frac{16}{5} = \frac{7}{5}$$

$$\therefore f\left(-\frac{3}{5}, -\frac{4}{5}\right) = 3 \times \left(-\frac{3}{5}\right) + 4 \times \left(-\frac{4}{5}\right) = -\frac{9}{5} - \frac{16}{5} = -\frac{25}{5} = -5$$

\therefore The maximum value of function is 5 at $f\left(\frac{3}{5}, \frac{4}{5}\right)$

The minimum value of function is -5 at $f\left(-\frac{3}{5}, -\frac{4}{5}\right)$.

* Partial Differentiable Equation (PDE):

Review of Ordinary differentiable equation

$$\frac{dy}{du} + ny = u^2 \quad \text{--- ODE}$$

$$\frac{dy}{du} + \frac{y^2}{u^2} = 1 \quad \text{--- PDE}$$

i) Separate Variable

ii) Homogeneous Equation

iii) Linear Equation

* Separate Variable:

$$\text{a. } \sqrt{1-u^2} dy + \sqrt{1-y^2} du = 0$$

\rightarrow Solution,

$$\sqrt{1-u^2} dy = -\sqrt{1-y^2} du$$

$$\frac{dy}{\sqrt{1-y^2}} = -\frac{du}{\sqrt{1-u^2}}$$

∴ Integrating both sides, we get,

$$\int \frac{dy}{\sqrt{1-y^2}} = - \int \frac{du}{\sqrt{1-u^2}} \quad \left(\frac{du}{\sqrt{1-u^2}} = \sin^{-1} u \right)$$

a. $\sin^{-1} y = -\sin^{-1} u + C$

$\therefore \sin^{-1} u + \sin^{-1} y = C$ is the required equation.

b) $\frac{dy}{du} + y = 1$

→ Solution,

$$\frac{dy}{du} = 1-y$$

a. $\frac{dy}{(1-y)} = du$

Integrating on both sides, we get,

$$\int \frac{dy}{(1-y)} = \int du \quad \left(\frac{du}{1-u} = -\log(1-u) \right)$$

a. $-\log(1-y) = u + C$

a. $u + \log(1-y) = C$ is required equation.

c) $dy = e^{u-y} du + u \cdot e^{-y} du$

a. $dy = e^u \cdot e^{-y} du + u \cdot e^{-y} du$ take common e^{-y}

$$\textcircled{a}, dy = e^{-y} (e^u du + u du)$$

$$\textcircled{a}, \frac{dy}{e^{-y}} = e^u du + u du$$

Integrating both sides,

$$\int \frac{dy}{e^{-y}} = \int e^u du + \int u du$$

$$\textcircled{a}, \int e^y dy = e^u + \frac{u^2}{2}$$

$$\textcircled{a}, e^y = e^u + \frac{u^2}{2} + C$$

$$\therefore e^u + \frac{u^2}{2} - e^y = C$$

is the required equation.

$$\textcircled{d}, (1+u^2)dy = (1+y^2)du$$

→ Solution,

$$\frac{dy}{(1+y^2)} = \frac{du}{(1+u^2)}$$

Integrating both sides,

$$\int \frac{dy}{(1+y^2)} = \int \frac{du}{(1+u^2)}$$

$$\left[\therefore \int \frac{dy}{(1+y^2)} = \tan^{-1} y \right]$$

$$\textcircled{a}, \tan^{-1} y = \tan^{-1} u + C$$

$$\left[\therefore \tan^{-1} y - \tan^{-1} u = C \right]$$

is the required equation.

* Homogeneous eq'

* Homogeneous Equation:

- equation in the form of $\Phi(\frac{y}{u})$

$$\text{put } v = \frac{y}{u}$$

$$\therefore y = vu \rightarrow \text{Product rule}$$

differentiating both side w.r.t u

$$\frac{dy}{du} = \frac{d(v \cdot u)}{du}$$

$$\frac{dy}{du} = v \frac{du}{du} + u \frac{dv}{du}$$

$$\therefore \frac{dy}{du} = v + u \frac{dv}{du}$$

$$\frac{y}{u} = v$$

$$9.7 \quad \frac{dy}{du} - \frac{y}{u} = -\frac{y^2}{u^2}$$

change $\frac{y}{u}$ to v . and put $\frac{dy}{du} = v + u \frac{dv}{du}$

$$9. \quad v + u \frac{dv}{du} - v = -\frac{v^2}{u^2}$$

$$9. \quad \frac{dv}{u^2} = -\frac{du}{v}$$

$$9. \quad v^{-2} dv = -u^{-1} du$$

Integrating both side

$$9. \quad \int v^{-2} dv = - \int u^{-1} du \quad \frac{1}{u}$$

$$9. \quad \frac{v^{-2+1}}{-2+1} = - \frac{u^{-1+1}}{1} \ln u$$

$$9. \quad -\frac{1}{v} = -\ln u + C$$

$$[\ln a + \ln b = \ln(ab)]$$

$$[\ln a - \ln b = \ln(\frac{a}{b})]$$

$$\text{a, } -\frac{1}{y/x} = -\ln u + \ln c$$

$$\text{a, } -\frac{1}{x} \times \frac{x}{y} = \ln\left(\frac{c}{u}\right)$$

$$\text{a, } -\frac{u}{y/x} = \ln\left(\frac{c}{u}\right)$$

$$\text{a, } e^{-u/y} = \frac{c}{u}$$

$$\boxed{\therefore u e^{-u/y} = c} \text{ is required eq? } \checkmark$$

$$\text{b)} \frac{dy}{du} = \frac{y-2u}{u}$$

$$y = v u$$

→ solution,

$$\frac{dy}{du} = \frac{y}{u} - \frac{2u}{u}$$

$$\frac{dy}{du} = v + u \frac{dv}{du}$$

$$\text{a, } v + u \frac{dv}{du} = v - 2$$

$$\text{a, } u \frac{dv}{du} = -2$$

$$\text{a, } \frac{dv}{du} = -2 \frac{1}{u}$$

Integrating on both sides,

$$\int dv = -2 \int \frac{du}{u}$$

$$\boxed{\int \frac{1}{u} du = \ln u}$$

$$\text{a, } v = -2 \ln u$$

$$\text{a, } \frac{y}{u} = -2 \ln u + c \quad \checkmark \quad \left\{ \text{a, } \frac{y}{u} = \ln(c/u) \right.$$

OR

$$\text{a, } \frac{y}{u} = -2 \ln u + \ln c$$

$$\text{a, } \frac{y}{u} = \ln(c u) \quad \checkmark$$

* Linear Equation:

$$\frac{dy}{du} + py = Q$$

where P and Q are function of u (not y)

I.F = integrating factor

i.e. $I.F = e^{\int P du}$

Multiplying the linear eqⁿ with I.F

$$\frac{d(y \cdot I.F)}{du} = Q \cdot I.F$$

or, $\int d(y \cdot I.F) = \int Q du \times I.F$

$$y \cdot I.F = \int Q du \times I.F + C$$

a) $\tan u \frac{dy}{du} + y = \sec u$

→ changing this eqⁿ to $\frac{dy}{du} + py = Q$

dividing both sides by $\tan u$, we get

$$\frac{dy}{du} + \frac{1}{\tan u} y = \frac{\sec u}{\tan u}$$

or $\frac{dy}{du} + \cot u \cdot y = \frac{1}{\cos u} \times \frac{\cos u}{\sin u} = \frac{1}{\sin u}$ - (i)

Comparing eqⁿ ① with $\frac{dy}{du} + py = Q$

$$\therefore P = \cot u$$

$$Q = \frac{1}{\sin u}$$

Now,

$$\text{Integrating Factor (I.F)} = e^{\int pdv}$$

$$= e^{\int \cot u du} \quad \left[\because \int \cot u du = \ln \sin u \right]$$

$$= e^{\ln \sin u}$$

$$= \sin u$$

Multiplying both sides by I.F, we get,

$$\sin u \left(\frac{dy}{du} + (vtu \cdot y) \right) = \frac{1}{\sin u} \times \sin u$$

$$\text{a, } d(y \cdot \sin u) = du$$

always $\frac{d(y \cdot I.F)}{du}$

Integrating both sides, we get

$$\int d(y \cdot \sin u) = \int du$$

$$\text{a, } \boxed{y \cdot \sin u = u + C} \quad \text{which is required eqn.}$$

$$\text{b, } \frac{dy}{du} + y = e^u$$

→ Comparing $\frac{dy}{du} + y = e^u$ with $\frac{dy}{du} + py = 0$

$$\therefore p = -1 \quad C = e^u$$

Now,

$$= \int P du = \int du = u$$

$$\text{Integrating Factor} = e^{\int pdv} = e^u$$

Multiplying both sides by e^u

$$e^u \left(\frac{dy}{du} + y \right) = e^u \cdot e^u$$

$$\text{or, } d(y \cdot e^u) = e^{2u} du$$

Integrating, we get

$$\text{or, } \int d(y \cdot e^u) = \int e^{2u} du$$

$$\text{or, } y \cdot e^u = \frac{1}{2} e^{2u} + C$$

is the required equation.

* Linear differentiable equation of Second Order

$$\frac{d^2y}{du^2} + \frac{dy}{du} + py = Q$$

$\frac{d}{du}$ = D = Differential operation

$$D^2y + D + Dy = Q \Rightarrow m^2 + m + p = 0 \Rightarrow$$

STEPS for solution :-

⇒ Auxillary Equation

$$\frac{d}{du} = D = m$$

$$y = 1$$

$m = \text{two values roots}$

② Solution :-

① $m = \text{roots and are real and different.}$

$$\therefore y = C_1 e^{m_1 u} + C_2 e^{m_2 u}$$

C_1 and C_2 are constant.

b) When roots are real and equal.

$$y = (C_1 + C_2 u) e^{mu}$$

$$m = m_1 = m_2$$

c) When roots are imaginary and distinct.

$$m = a + ib$$

$$y = e^{au} (C_1 \cos bu + C_2 \sin bu)$$

* Solve :-

$$a) \frac{d^2y}{du^2} + 5 \frac{dy}{du} + 6y = 0$$

→ Solution,

Auxiliary equation is,

$$m^2 + 5m + 6 = 0$$

$$\text{or } m^2 + 2m + 3m + 6 = 0$$

$$\text{or } m(m+2) + 3(m+2) = 0$$

$$\text{either } m_1 = -2 \text{ and } m_2 = -3$$

∴ Required eqⁿ is,

Roots m_1 and m_2 are real and different.

So,

$$y = C_1 e^{-2u} + C_2 e^{-3u}$$

$$b) \frac{d^2y}{du^2} + \omega^2 y = 0$$

$$\omega = a + ib = \text{Imaginary}$$

Auxiliary eqⁿ is,

$$\text{or } m^2 + \omega^2 = 0$$

$$\therefore m = \pm \omega i \quad 0 + \omega i$$

$$m = a \pm ib$$

$\therefore a = 0$ Roots are imaginary & distinct.

$$b = \omega$$

$$\therefore y = e^{\omega t} (C_1 \cos \omega n + C_2 \sin \omega n) \#.$$

$$37) \frac{d^2y}{du^2} + 4 \frac{dy}{du} + 12y = 0 \quad ???$$

→ solution,

given auxiliary eqn 13

$$m^2 + 4m + 12 = 0 \quad \text{---} \textcircled{1}$$

Comparing eqn 13 with

$$am^2 + bm + c = 0$$

$$\therefore a = 1$$

$$b = 4$$

$$c = 12$$

Now,

$$m = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad -b \pm \sqrt{b^2 - 4ac}$$

$$\therefore 4ac = \pm \sqrt{16 - 4 \times 1 \times 12} = -4 \pm \sqrt{16 - 4 \times 1 \times 12}$$

$$= \pm \sqrt{\frac{-32}{2}} \quad = -4 \pm \frac{\sqrt{-32}}{2}$$

$$\therefore \pm 2\sqrt{32}$$

$$m^2 + 6m - 2m + 12 = 0$$

$$m(m+6) - 2(m+6) = 0$$

$$(m-2)(m+6)$$

impossible!!!

so

use quadratic eqn

$$d) (D+3)^2 y = 0$$

$$D = \frac{dy}{du} \quad \frac{d^2y}{du^2} = D = m$$

→ Auxilliary Equation is

$$(m+3)^2 = 0$$

roots are,

$$(m+3)(m+3) = 0$$

real

Either, $m = -3, -3$

and

Required eqⁿ is,

equal,

∴

so

$$y = (C_1 + C_2 u) e^{-3u}$$

H

Partial Differential Equation:

→ The equation in the form of $f(u, y, z, p, q) = 0$ is called partial differential equation of first order. (u, y, z) are independent variable

$$p = \frac{\delta z}{\delta u}, \quad q = \frac{\delta z}{\delta y}$$

Second order:

$$P(u, y, z, p, q, r, s, t, \dots) = 0$$

$$r = \frac{\delta^2 z}{\delta u^2}, \quad s = \frac{\delta^2 z}{\delta u \delta y}, \quad t = \frac{\delta^2 z}{\delta y^2}, \quad \dots \text{etc.}$$

Origin of first order PDE

- eliminating constants
- eliminating arbitrary function.

1) Form PDE by eliminating constants a and b from:

$$(u-a)^2 + (y-b)^2 + z^2 = f = 1 \quad \text{--- (A)}$$

2) Form PDE by eliminating arbitrary function f from the eqⁿ,

$$z = ny + f(u^2 + y^2).$$

~~1st step~~ w.r.t to u

$$\Rightarrow \frac{\delta(u-a)^2}{\delta u} \times \frac{\delta(u-a)}{\delta u} + \frac{\delta(y-b)^2}{\delta u} + \frac{\delta z^2}{\delta u} \times \frac{\delta z}{\delta u} = 0 \quad (\because \frac{\delta z}{\delta u} = p)$$

$$\text{a, } 2(u-a) \times 1 + 0 + 2z \times p = 0$$

$$\text{a, } 2(u-a) = -2zp$$

$$\text{a, } u-a = -zp \quad \text{--- (i)}$$

$$(\because \frac{\delta z}{\delta y} = q)$$

Partial derivative w.r.t to y

$$\frac{\delta(u-a)^2}{\delta y} \times \frac{\delta(u-a)}{\delta y} + \frac{\delta(y-b)^2}{\delta y} \times \frac{\delta(y-b)}{\delta y} + \frac{\delta z^2}{\delta y} \times \frac{\delta z}{\delta y} = 0$$

$$\text{a, } 2(u-a) \times 0 + 2(y-b) \times 1 + 2z \times q = 0$$

$$y-b = -2q \quad \text{--- (ii)}$$

from eqⁿ (i) and (ii) we get, in eqⁿ - (A)

$$(u-a)^2 + (y-b)^2 + z^2$$

$$(-zp)^2 + (-2q)^2 + z^2 = 1$$

$$z^2 (p^2 + q^2 + 1) = 1$$

P.D.E ~~✓~~

2nd Gf

$$z = uy + f(u^2 + y^2)$$

Partial derivative w.r.t u_i

$$\frac{S_z}{S_n} = y + \frac{S f(u^2+y^2)}{S(u^2+y^2)} \times \frac{S(u^2+y^2)}{S_n}$$

$$P = y + f'(u^2 + y^2) \times 2u \quad \text{---(i)}$$

partial derivatives w.r.t. y.

$$\frac{S_x}{S_y} = u + \frac{S_f(u^2+y^2)}{S(u^2+y^2)} \times \frac{d(u^2+y^2)}{S_y}$$

$$q = u + f'(u^2 + y^2) \times 2y \quad \dots \dots \dots \quad (i)$$

from eqⁿ (i) and (ii))

from eqⁿ (i)

$$f'(u^2+y) = \frac{p-y}{2} \quad - \textcircled{111}$$

from 11

$$f'(u^2+y^2) = \frac{9-u}{2y} \quad \text{--- (14)}$$

equating eqⁿ 11 & 12

$$\frac{P-Y}{2u} = \frac{q-u}{2u}$$

$$\infty, \quad \boxed{py - y^2 = q^u - u^2}$$

$$\frac{P-g}{x^m} = \frac{g-u}{xy}$$

$$P - g - \frac{g+u}{4} = 0$$

PDE ~~4~~

$$\cancel{E} \cancel{M} \quad \cancel{p} y^2 - q u^2 = q u^2$$

РУ-9

Q. Form PDE by eliminating ϕ from

$$lu + my + nz = \phi(u^2 + y^2 + z^2)$$

→ Solution

differentiating both sides w.r.t u

$$\frac{\delta(lu)}{\delta u} + \frac{\delta(my)}{\delta u} + \frac{\delta(nz)}{\delta u} = \frac{\delta \phi(u^2 + y^2 + z^2)}{\delta u}$$

$$\text{or } l + 0 + np = \frac{\delta \phi(u^2 + y^2 + z^2)}{\delta(u^2 + y^2 + z^2)} \times \frac{\delta(u^2 + y^2 + z^2)}{\delta u} = \frac{\delta z^2}{\delta z} \times \frac{\delta z}{\delta u}$$

$$\text{or } l + np = \phi'(u^2 + y^2 + z^2) \times (2u + 2zp) \quad \text{--- (i)}$$

diff. w.r.t y on both sides,

$$\frac{\delta(lu)}{\delta y} + \frac{\delta(my)}{\delta y} + \frac{\delta nz}{\delta y} = \frac{\delta \phi(u^2 + y^2 + z^2)}{\delta(y^2 + z^2)} \times \frac{\delta(y^2 + z^2)}{\delta y}$$

$$\text{or } 0 + m + nq = \phi'(u^2 + y^2 + z^2) \times (2y + 2zq)$$

$$\therefore \phi'(u^2 + y^2 + z^2) = \frac{m + nq}{2y + 2zp} \quad \text{--- (ii)}$$

$$\therefore \phi'(u^2 + y^2 + z^2) = \frac{l + np}{2u + 2zp} \quad \text{--- from eq (i)}$$

Equating eq (i) and (ii), we get,

$$\frac{m + nq}{2y + 2zp} = \frac{l + np}{2u + 2zp}$$

$$a. mx + nqz = ly + npz$$

~~∴ P.D.E.~~

$$(m+zq)(m+nq) = (l+np)(y+zp)$$

Linear P.D.E of first order:

④

$$Pp + Qq = R$$

Langrange Auxilliary equation:

$$\frac{du}{P} = \frac{dy}{Q} = \frac{dz}{R}$$

$$\left[P = \frac{\partial z}{\partial u}, Q = \frac{\partial z}{\partial y} \right]$$

solution is $\phi(u, y, v) = 0$ [P, Q, are the function of u, y, z]

$$u(u, y, z) = C_1$$

$$v(u, y, z) = C_2$$

★ Solve : $y^2pz + u^2qz = ny^2$

→ Solution,

Comparing with $Pp + Qq = R$

$$\therefore P = y^2z \quad Q = u^2z \quad R = ny^2$$

Now,

Langrange Auxilliary eqⁿ.

$$\frac{du}{y^2z} = \frac{dy}{u^2z} = \frac{dz}{ny^2}$$

taking only two variable

$$\frac{du}{y^2 z} = \frac{dy}{u^2 z}$$

Integrating both sides,

$$\int \frac{u^2 du}{z} = \int y^2 dy$$

$$\text{a, } \frac{u^3}{3} = \frac{y^3}{3} + C_1$$

$$\text{a, } \frac{C_1}{3} = \frac{u^3 - y^3}{3} \quad \therefore C_1 = u^3 - y^3$$

Now,

again taking two variables,
taking first and 3rd

$$\frac{du}{y^2 z} = \frac{dz}{u y^2}$$

taking Integrating both sides,

$$\int u du = \int z dz$$

$$\text{a, } \frac{u^2}{2} = \frac{z^2}{2} + C_2$$

$$\text{a, } \frac{C_2}{2} = \frac{u^2 - z^2}{2} \quad \therefore C_2 = u^2 - z^2$$

Solution is,

$$\phi(u^3 - y^3, u^2 - z^2) = 0$$

$$* P + q = u$$

→ Solution,

Comparing question with $Pp + Qq = R$
 $\therefore P = 1 \quad Q = 1 \quad R = u$

Now,

Lagrangian Auxiliary eqⁿ is

$$\frac{dy}{P} + \frac{dy}{\phi} - \frac{dz}{R}$$

$$\text{a, } \frac{du}{1} = \frac{dy}{1} = \frac{dz}{u}$$

taking first and second,

Integrating
 $\int du = \int dy$

$$\text{a, } u - y = C_1$$

taking first and 3rd

$$du = \frac{dz}{u}$$

$$\text{a, } u du = dz$$

Integrating both sides,

$$\int u du = \int dz$$

$$\text{a, } \frac{u^2}{2} = z + C_2$$

$$\text{a, } \frac{u^2}{2} - \frac{2z}{2} = \frac{C_2}{2}$$

$$\text{a, } C_2 = u^2 - 2z$$

Solution is $\phi(z_1, z_2) = 0$

$$\phi(u-y, u^2-2z) = 0 \#$$

by $(y-z) \frac{\partial z}{\partial u} + (u-y) \frac{\partial z}{\partial y} = z-u$

→ Solution,

$$(y-z)p + (u-y)q = z-u \quad \left(\because \frac{\partial z}{\partial u} = p, \frac{\partial z}{\partial y} = q \right)$$

Comparing with $Pp + Qq = R$

$$\therefore P = (y-z), Q = (u-y), R = (z-u)$$

Lagrange Auxilliary eqⁿ is

$$\frac{du}{P} - \frac{dy}{Q} = \frac{dz}{R}$$

$$\text{or, } \frac{du}{(y-z)} - \frac{dy}{(u-y)} = \frac{dz}{(z-u)}$$

Using multiplier 1, 1, 1, we get,

$$\frac{du+dy+dz}{y-z+u-y+z-u} = k$$

$$\text{or, } du+dy+dz = k$$

Integrating,

$$C_1 + u + y + z = 0$$

$$u + y + z = C_1$$

Using multiplier u, y and z

$$\begin{aligned} \underline{ndu + zdy + ydz} &= k \\ \underline{ny - nz + zu - zy + yz - ny} \end{aligned}$$

$$\text{a, } \int ndu + \int zdy + \int ydz = 0$$

$$\text{3, } \frac{u^2}{2} + zy + yz = C_2$$

\therefore solution is $\phi(c_1, c_2) = 0$

$$\Rightarrow \phi(u+y+z, \frac{u^2}{2} + zy + yz) = 0 \quad \#.$$

$$\text{c) } (mz - ny) p + (nu - lz) q = ly - mu$$

\rightarrow solution,

Comparing question with $Pp + Qq = R$

$$\therefore P = (mz - ny), Q = (nu - lz), R = ly - mu$$

then,

Auxiliary eqⁿ is

$$\frac{du}{(mz - ny)} = \frac{dy}{(nu - lz)} = \frac{dz}{(ly - mu)}$$

Using multiplier u, y, z , we get,

$$\begin{aligned} \underline{ndu + ydy + zdz} &= k \\ mz^2 - ny^2 + nuy - lyz + lyz - muyz \end{aligned}$$

c, Integrating both sides

$$\frac{u^2}{2} + \frac{y^2}{2} + \frac{z^2}{2} = \frac{C_1}{2}$$

$$\therefore C_1 = u^2 + y^2 + z^2$$

Using multiplier l, m, n

$$\frac{ldu + mdy + ndz}{mzl - myl + mny - mz + nlz - nny} = k$$

a. $\int l du + \int mdy + \int ndz$
 $\therefore C_2 = l + m + n$

\therefore Solution is $\Phi (u^2 + y^2 + z^2, l + m + n)$

$$d) u^2 \frac{\partial z}{\partial u} + y^2 \frac{\partial z}{\partial y} = (u+y)z$$

\rightarrow Solution,

$$u^2 p + y^2 q = (u+y)z \quad \dots \text{(i)}$$

Comparing eq (i) with $Pp + Qq = R$
 $\therefore P = u^2 \quad Q = y^2 \quad R = (u+y)z$

then Langrange auxilliary eq is,

$$\frac{du}{P} = \frac{dy}{Q} = \frac{dz}{R}$$

$$\text{a. } \frac{du}{u^2} = \frac{dy}{y^2} = \frac{dz}{(u+z+y)}$$

taking 1st and 2nd

$$\frac{du}{u^2} = \frac{dy}{y^2}$$

Integrating, $\int \frac{du}{u^2} = \int \frac{dy}{y^2}$

$$\text{a. } \int u^2 du = \int y^{-2} dy$$

$$\text{a. } -\frac{u^{-2+1}}{-2+1} = -\frac{y^{-2-1+1}}{-2+1}$$

$$\text{a. } \rightarrow \frac{1}{u} = \frac{1}{y} \neq C_1$$

$$\text{a. } \frac{1}{u} - \frac{1}{y} = C_1$$

$$\text{a. } \frac{y-u}{uy} = C_1$$

taking 1st and 3rd variables,

$$\int \frac{du}{u^2} = \int \frac{dz}{uz + yz}$$

$$\text{c. } \int u^{-2} du = \int \frac{dz}{uz} + \int \frac{dz}{yz}$$

$$\text{a. } \frac{u^{-2+1}}{-2+1} = \frac{\ln z}{u} + \frac{\ln z}{y}$$

$$\text{a. } -\frac{1}{u} = \frac{\ln z}{u} + \frac{\ln z}{y} + C_2$$

Taking 1st and 2nd

$$\int \frac{du}{u^2} = \int \frac{dy}{y^2}$$

$$\Rightarrow u^{-2} \int du = \int y^{-2} dy$$

$$\Rightarrow +\frac{1}{u} - \frac{1}{y} = C_1$$

$$\therefore C_1 = \frac{1}{u} - \frac{1}{y}$$

Again,

$$\frac{du - dy}{(u^2 - y^2)} = \frac{dz}{z(u+y)}$$

$$\frac{d(u-y)}{(u-y)(u+y)} = \frac{dz}{z(u+y)}$$

$$\int d = \int z^{-1} dz$$

$$C_2 = \log(1-z) \Rightarrow \log 2 + \log u$$

$$(u^{-2}) = \frac{C_1}{C_2 \left(\frac{u-1}{u}\right)^2}$$

$$\textcircled{e} \quad \frac{y^2 z}{u} \frac{\partial z}{\partial u} + u^2 z \frac{\partial z}{\partial y} = u y^2$$

→ Comparing $y^2 z p + u^2 z q = u y^2$ with $Pp + Qq = R$

$$\therefore P = y^2 z, Q = u^2 z, R = u y^2$$

Now

Lagrange Auxillary eq? is

$$\frac{du}{P} = \frac{dy}{Q} = \frac{dz}{R}$$

$$\text{a. } \frac{du}{y^2 z} = \frac{dy}{u^2 z} = \frac{dz}{u y^2}$$

Using multipliers $-u, y^2, z$.

$$\text{a. } \frac{-u dx + y dy + z dz}{-u y^2 z - y u z^2} \quad \text{taking 1st and 2nd}$$

$$\int z u^2 du = \int y^2 dy$$

$$\text{a. } \frac{z u^3}{3} = \frac{y^3}{3} + C_1$$

$$\therefore C_1 = z(u^3 - y^3)$$

$$\text{taking 1st and 3rd } \frac{du}{y^2 z} = \frac{dz}{u y^2}$$

$$\text{a. } \int u y^2 du = \int y^2 z dz$$

$$\text{a. } \frac{y^2 u^2}{2} = \frac{y^2 z^2}{2} + \frac{C_2}{2}$$

$$\text{a. } C_2 = y^2(u^2 - z^2)$$

\therefore solution is $\Phi(z(u^3 - y^3), y^2(u^2 - z^2))$

$$f) uP - yq = y^2 - u^2$$

→ solution,

Comparing $uP - yq = y^2 - u^2$ with $Pp + Qq = R$
 $\therefore P = u \quad Q = y, \quad R = y^2 - u^2$
 Langrang Auxilliary Eqⁿ is,

$$\frac{du}{P} = \frac{dy}{Q} = \frac{dz}{R}$$

$$\text{a. } \frac{du}{u} = \frac{dy}{y} = \frac{dz}{y^2 - u^2}$$

Using multiplier i, -y, 1

$$\text{a. } \frac{u du + (-y) dy + dz}{u^2 - y^2 + y^2 - u^2} = k$$

$$\text{a. } u du - y dy + dz = 0$$

Integrating we get

$$\text{a. } \int u du - \int y dy + \int dz = 0$$

$$\text{a. } \frac{u^2}{2} - \frac{y^2}{2} + z = \frac{C_1}{2}$$

$$\therefore C_1 = (u^2 - y^2 + 2z)$$

taking 1st and 3rd

$$\int \frac{du}{u} = \int \frac{dz}{y^2 - u^2}$$

taking 1st and 2nd

$$\int \frac{du}{u} = \int \frac{dy}{y}$$

$$\text{a. } \ln u =$$

$$\text{a. } \ln u = \ln y + C_2$$

$$\text{a. } C_2 = \ln u - \ln y = \ln \left(\frac{u}{y} \right)$$

$$8) u^2 P + Q = z^2$$

→ Solution,

Comparing eqⁿ with $Pp + Qq = R$

$$\because P = u^2 \quad Q = 1 \quad R = z^2$$

Now,

longrang eqⁿ of Auxillary is,

$$\frac{du}{P} = \frac{dy}{Q} = \frac{dz}{R}$$

$$\text{or } \frac{du}{u^2} = \frac{dy}{1} = \frac{dz}{z^2}$$

Integrating 1st and 2nd Integrating 1st and 3rd

$$\text{a. } \int \frac{du}{u^2} = \int dy$$

$$\int \frac{du}{u^2} = \int \frac{dz}{z^2}$$

$$\text{or } \frac{u^{-2+1}}{-2+1} = y + C_1$$

$$\text{or } \frac{u^{-2+1}}{-2+1} = \frac{z^{-2+1}}{-2+1} + C_2$$

$$\therefore C_1 = -\frac{1}{u} - y$$

$$\text{or } \frac{1}{u} = -\frac{1}{z} + C_2$$

$$\therefore C_2 = \frac{1}{z} - \frac{1}{u}$$

∴ Solution is $\phi(C_1, C_2)$

$$\phi = \left(-\frac{1}{u} - y, \frac{1}{z} - \frac{1}{u} \right) \#$$

PDE of Second Order:

PDE of the form:

$$F(D, D')z = f(u, y)$$

(Partial diff. Eq.)

is PDE of second order.

where,

$$D = \frac{\partial}{\partial u} \quad \text{and} \quad D' = \frac{\partial}{\partial y}$$

$$r = D^2 = \frac{\partial^2 z}{\partial u^2} \quad S = DD' = \frac{\partial^2 z}{\partial u \cdot \partial y} \quad t = D'^2 = \frac{\partial^2 z}{\partial y^2}$$

* Complementary Function: (C.F)

Solution of $F(D, D')z = 0$ is called complementary function.

→ Solution:-

1) When roots are real and distinct, then

$$C.F = \Phi_1(y + m_1 u) + \Phi_2(y + m_2 u) + \Phi_3(y + m_3 u) + \dots$$

2) When roots are real and same (equal)

$$C.F = \Phi_1(y + m_1 u) + u \Phi_2(y + m_2 u) + u^2 \Phi_3(y + m_3 u)$$

* Particular Integral (P.I)

$$z = \underline{f(u, y)} \quad \text{is} \quad \underline{F(D, D')}$$

Called particular integral.

Complete Solution:

$$z = C.F + P.I$$

Note: For P.I

Binomial Expansion

$$(1+D)^{-1} = 1 - D + D^2 - D^3 + \dots$$

$$(1-D)^{-1} = 1 + D + D^2 + D^3 + \dots$$

$$(1+D)^{-2} = 1 - 2D + 3D^2 - 4D^3 + \dots$$

$$(1-D)^{-2} = 1 + 2D + 3D^2 + 4D^3 + \dots$$

Particular Integral:

$$1) f(u, y) = u^m \cdot y^n$$

$$Z = F(D, D')^{-1} u^m \cdot y^n$$

Use binomial expansion.

$$2) f(u, y) = \cos(au+by) \quad \text{or} \quad f(u, y) = \sin(au+by)$$

$$Z = \frac{\cos(au+by)}{F(+D^2, +DD', +D^2)} = \frac{\cos(au+by)}{F(-a^2, -ab, -b^2)}$$

$$3) f(u, y) = \frac{e^{au+by}}{F(D, D')} = \frac{e^{au+by}}{F(a, b)}$$

{provided that $F(a, b) \neq 0$ }

if $F(a, b) = 0$, differentiate denominator with respect to D and multiply by u .

Q. Solve :-

$$(D^2 - D'^2) Z = u - y$$

→ Solution :-

for complementary function,

$$F(D, D') = 0$$

$$D = m, D' = 1$$

Auxiliary equation is put always

$$m^2 - 1 = 0$$

$$m = \pm 1$$

$$\therefore m_1 = 1, m_2 = -1$$

$$D = m$$

$$\text{and } D' = 1$$

$$\therefore C.F = \Phi_1(y+u) + \Phi_2(y-u)$$

Now,

Particular Integral (P.I),

$$Z = \frac{f(u, y)}{F(D, D')} \text{ is}$$

$$Z = \frac{(u-y)}{(D^2 - D'^2)}$$

doing binomial expansion

take D^2 common

$$Z = \frac{u-y}{D^2 \left(1 - \frac{D'^2}{D^2} \right)}$$

$$S \frac{\cos(\ln+ny)}{S(\ln+ny)} \times \frac{S(\ln+ny)}{Sy} = -\sin(\ln+ny) \times m$$

CLASSE MATE
Date _____
Page _____

$$= -A \frac{[l^2 \cos(\ln+ny) - D' \cos(\ln+ny)]}{l^4 + m^2}$$

$$P. = - \frac{A(l^2 \cos(\ln+ny) + \frac{\sin(\ln+ny) \times m}{l^4 + m^2})}{l^4 + m^2}$$

$$P. I = - \frac{1}{l^4 + m^2} (A(l^2 \cos(\ln+ny) - \sin(\ln+ny) \cdot m))$$

$$Z = P \cdot I + C \cdot I \quad u^m y^n = f(ny)$$

$$(1 - D)^{-1} = 1 + D + D^2 + D^3 + \dots$$

$$(1 + D)^{-1} = 1 - D + D^2 - D^3 + \dots$$

$$(1 - D)^{-2} = 1 + 2D + 3D^2 + 4D^3 + \dots$$

$$(1 + D)^{-2} = 1 - 2D + 3D^2 - 4D^3 + \dots$$

$$f(ny) = \frac{\cos(\ln+ny)}{\sin(\ln+ny)}$$

$$F(\cos(\ln+ny)) = \cos(\ln+ny)$$

$$F(D^2, DD', D'^2) = (-a^2, -ab, -b^2)$$

$D^2 \times 2$

* Important long question:-

* Solve $(D^2 - D'^2)z = u - y$

→ Solution:-

for complementary function
 $F(D, D') = 0$

where $D = m$ and $D' = 1$

$\therefore m^2 - 1 = 0$ (Auxiliary eqn)
 $\therefore m^2 = 1$

$\therefore m = \pm 1 \quad m_1 = +1 \quad m_2 = -1$

\therefore complementary function (C.F) =

$$\Phi_1(y+u) + \Phi_2(y-u)$$

Now,

Particular integral (P.I) =

$$Z = F(u, y)$$

$$Z = \frac{(u-y)}{(D^2 - D'^2)}$$

$$= \frac{(u-y)}{D^2 \left(1 - \frac{D'^2}{D^2}\right)} = \frac{1}{D^2} \left(1 - \frac{D'^2}{D^2}\right)^{-1} (u-y)$$

$$= \frac{1}{D^2} \left(1 + \frac{D'^2}{D^2} + \frac{D'^4}{D^4} + \dots\right) (u-y)$$

$$= \frac{1}{D^2} \left[(u-y) + (u-y) \frac{D'^2}{D^2} \right]$$

$$= \frac{1}{D^2} [(u-y) + 0]$$

$$\begin{aligned}
 &= D^{-2}(u-y) \\
 &\text{Integrating w.r.t. } u \text{ because } D = \frac{\partial}{\partial u}, D' = \frac{\partial}{\partial y} \\
 &= \frac{\partial}{\partial u}(D^{-2}(u-y)) \\
 &= D^{-1}\left(\frac{u^2}{2} - uy\right) \\
 &= \frac{\partial}{\partial u}\left(D^{-1}\left(\frac{u^2}{2} - uy\right)\right) \\
 &= \frac{u^3}{6} - \frac{u^2y}{2} \\
 &\therefore \text{Particular integral is } \frac{u^3}{6} - \frac{u^2y}{2}
 \end{aligned}$$

$$\therefore \text{Complete soln } (z) = \Phi_1(y+u) + \Phi_2(y-u) + \frac{u^3}{6} - \frac{u^2y}{2}$$

* Solve:

$$(D^2 - D')Z = A \cos(du + my)$$

→ Solution,

for complementary function,

$$F(D, D') = 0$$

$$D = m, D' = 1$$

∴ Auxiliary eqn is

$$m^2 - 1 = 0$$

$$m^2 = 1$$

$$m = \sqrt{1}$$

$$m = \pm 1$$

$$\therefore m_1 = 1, m_2 = -1$$

Since roots are real and distinct.
so,

Complementary function (C.F) = $\Phi_1(y+u) + \Phi_2(y-u)$
Now,

for particular integral

$$Z = \frac{f(u, y)}{F(D, D')}$$

$$Z = \frac{A \cos(lu+my)}{(D^2 - D')}$$

$f(u, y)$ is \cos , so

$$\frac{\cos(lu+my)}{F(D^2, DD', D')}$$

$$\therefore D^2 = -l^2, D'^2 = -m^2$$

we can't put $D' = -m$
so make D' to D'^2

$$\text{to put } -m^2$$

$$\text{or } Z = \frac{A \cos(lu+my)}{(-l^2 - D'^2)}$$

$$\text{or } Z = \frac{A \cos(lu+my) \times (l^2 - D')}{(-l^2 + D^2) \quad (l^2 - D^2)}$$

$$\text{or } Z = \frac{A \cos(lu+my) \times (l^2 - D')}{(-l^4 - D'^2) \quad D'^2 = -m^2}$$

$$\text{or } Z = \frac{A \cos(lu+my) \times (l^2 - D')}{(-l^4 - m^2)}$$

$$\text{or, } Z = - \frac{A \cos(lu+my) l^2 - D' A \cos(lu+my)}{(l^4 + m^2)}$$

$$\text{or, } Z = - \frac{A l^2 \cos(lu+my) + S(D' A \cos(lu+my))}{(l^4 + m^2)} \quad [8y]$$

$$a, Z = -\frac{Al^2}{(l^4+m^2)} \cos(lu+my) + AS \frac{\cos(lu+my)}{S(lu+my)} \times \frac{S(lu+my)}{Sy}$$

$$a, Z = -\frac{Al^2}{(l^4+m^2)} \cos(lu+my) + (-) \sin(lu+my) \times m \times A$$

$$\therefore Z = -\frac{Al^2}{(l^4+m^2)} \cos(lu+my) - \frac{Am \sin(lu+my)}{(l^4+m^2)}$$

∴ Complete Solution.

$$Z = C.F + P.I$$

$$= \Phi_1(y+u) + \Phi_2(y-u) - \frac{Al^2}{(l^4+m^2)} \cos(lu+my) - m \sin(lu+my)$$

* Find particular integral of $eq^n (D^2 - D') = Z = 2y - u^2$

$$D = \frac{8}{8u}, D' = \frac{8}{8y}$$

$$u^2 y - \frac{u^4}{4} + \frac{u^3}{3}$$

→ Solution,

Particular integral is given as

$$Z = \frac{F(u, y)}{F(D, D')} = f \frac{(2y - u^2)}{(D^2 - D')}$$

Using binomial expansion,

$$Z = \frac{1}{D^2} \left(1 - \frac{D'}{D^2} \right)^{(2y - u^2)}$$

Let $\left(\frac{D'}{D^2}\right)$ be D $(1-D) = 1+D+D^2+\dots$

classmate

Date _____

Page _____

$$Z = \frac{1}{D^2} \left(1 - \frac{D'}{D^2}\right)^{-1} (2y - u^2)$$

$$= \frac{1}{D^2} \left(1 + \frac{D'}{D^2} + \frac{D'^2}{D^4} + \dots\right) (2y - u^2)$$

$$= \frac{1}{D^2} \left[(2y - u^2) + \frac{1}{D^2} (D' (2y - u^2)) \right]$$

$$= \frac{1}{D^2} \left[(2y - u^2) + \frac{1}{D^2} \frac{8(2y - u^2)}{8y} \right]$$

$$= \frac{1}{D^2} \left[(2y - u^2) + \frac{1}{D^2} \times 2 \right]$$

$$= \frac{1}{D^2} (2y - u^2) + \frac{1}{D^4} 2$$

$$= \frac{8(2y - u^2)}{8u^2} D + \frac{1}{D^3} 2u$$

$$= \frac{8(2uy - \frac{u^3}{3})}{8u} + \frac{1}{D^2} \frac{2u^2}{2}$$

$$= \frac{2u^2 y - \frac{u^4}{12}}{2} + \frac{2u^3}{2 \times 3} \frac{1}{D}$$

$$= u^2 y - \frac{u^4}{12} + \frac{u^4}{3 \times 4}$$

$$= u^2 y \quad \#.$$

$$(D^2 - D) Z = 2y - u^2$$

$$\frac{8^2 Z}{8u^2} - \frac{8^2 Z}{8y^2}$$

$$\therefore (D^2 - D^2) Z = 2y - u^2$$

$$D \cdot Z = \frac{1}{D - D^2}$$

$$(2y - u^2) \times \frac{1}{D - D^2}$$

* Solve:

$$\text{D} \frac{S^2 z}{S u^2} + 2 \frac{S^2 z}{S u S y} + \frac{S^2 z}{S y^2} = \sin(2u+3y)$$

$$\Rightarrow (D^2 - 2DD' + D'^2)z = \sin(2u+3y)$$

for complementary function

$$D = m \quad D' = 1$$

The auxilliary Eqⁿ is,

$$m^2 - 2m + 1 = 0$$

$$\text{a, } m^2 - m - m + 1 = 0 \quad \text{a, } m(m-1) - 1(m-1) = 0$$

$$\text{Either } m_1 = 1 \quad m_2 = +1$$

The roots are equal and real.

$$\therefore C.F = \Phi_1(y+u) + \Phi_2(y+u).u$$

for particular Integration.

$$Z = \frac{f(u, y)}{F(D, D')} = \frac{\sin(2u+3y)}{D^2 - 2DD' + D'^2}$$

$$= \frac{\sin(2u+3y)}{-2^2 + 2 \times 2 \times 3 - 3^2} = -1 \sin(2u+3y)$$

$$\therefore Z = -\sin(2u+3y)$$

$$\therefore \text{Complete Soln is } Z = \Phi_1(y+u) + u \Phi_2(y+u) - \sin(2u+3y)$$

$$2.7 \quad r + 3s + 2t = u + y$$

$$\left(\frac{S^2 z}{S u^2} + 3 \frac{S^2 z}{S u S y} + 2 \frac{S^2 z}{S y^2} \right) = u + y$$

$$\text{or } D^2 + 3DD' + 2D^2 = u + y$$


for particular integral.

$$Z = \frac{f(u, y)}{F(D, D')} = \frac{u + y}{D^2 + 3DD' + 2D^2}$$

$$= \frac{1}{D^2} \left(1 + \frac{3D'}{D} \right)^{-1} (u + y) = \frac{1}{D^2} \left(1 - \frac{3D'}{D} + \frac{3^2 D'^2}{D^2} + \dots \right) (u + y)$$

$$= \frac{1}{D^2} (u + y - (u + y) \frac{3D'}{D} + 0)$$

$D^2 \rightarrow$ Integ. w.r.t. to u 2 times
 $D' =$ deriv. w.r.t. to y .



$$= \frac{1}{D^2} \left((u+y) - \frac{3}{D} (0+1) \right)$$

$$= \frac{1}{D^2} [(u+y) - 3u]$$

$$= \frac{1}{D^2} (y-2u)$$

$$= ny - \frac{2u^2}{3} D^{-1}$$

$$= \frac{u^2 y}{2} - \frac{u^3}{3}$$

for Complementary function,

$$D' = I \quad D = m$$

\therefore auxiliary eq is, $m^2 + 3m + 2 = 0$

$$\therefore m^2(m+2) + 1(m+2) = 0$$

$$m_1 = -2 \quad m_2 = -1$$

\therefore Roots are real and distinct

$$\therefore C.F = \Phi_1(y-2u) + \Phi_2(y-u)$$

$$\therefore Z = \frac{u^2 y}{2} - \frac{u^3}{3} + \Phi_1(y-2u) + \Phi_2(y-u) \quad \#$$

$$* \frac{S^2 Z}{S u^2} - 2 \frac{S^2 Z}{S u y} + \frac{S^2 Z}{S y^2} = e^{u+2y}$$

$$\Rightarrow (D^2 - 2D D' + D'^2) Z = e^{u+2y}$$

\rightarrow for particular integral.

$$P.I = \frac{e^{au+by}}{(D^2 - 2D D' + D'^2)}$$

$$= \frac{e^{au+by}}{1^2 - 2 \times 1 \times 2 + 2^2} = \frac{e^{u+2y}}{1}$$

$$\therefore \frac{e^{au+by}}{F(D, D')} = \frac{e^{au+by}}{(a, b)}$$

for complementary function:

$$D = m \quad D' = 1$$

Auxiliary eqⁿ is, $m^2 - 2m + 1 = 0$
 $\therefore m^2 - m - m + 1 = 0$

$$\therefore m_1 = 1 \quad m_2 = 1$$

$$\therefore C.F = \Phi_1(y+u) + \Phi_2 u(y+u)$$

\therefore Complete Sol^r,

$$Z = \Phi_1(y+u) + \Phi_2 u(y+u) + e^{u+2y} \#$$

$$4) (r - z^2 + = u^2)$$

$$= \underline{S^2} - \underline{S^2} \underline{S^2} Z$$

$$\underline{S^2}$$

$$\Rightarrow [D^2 + (a+b)DP' + abP'^2] z = ny$$

$$P-I = \frac{f(u, y)}{f(D, D')}$$

$$P^2 \left(1 + \frac{(a+b)D'}{D} \right)$$

$$= \frac{ny}{D^2 \left(1 + (a+b) \frac{D'}{D} + ab \frac{D'^2}{D^2} \right)}$$

$$\Rightarrow \frac{1}{D^2} \left(1 + (a+b) \frac{D'}{D} \right)^{-1} ny$$

$$\Rightarrow \frac{1}{D^2} \left(ny - (a+b) \frac{D'}{D} + (a+b) ny \frac{D'^2}{D^2} \right)$$

$$= \frac{1}{D^2} \left(ny - \frac{(a+b) ny}{D} + 0 \right)$$

$$\Rightarrow \frac{1}{D^2} \left(ny - \frac{(a+b) u^2}{2} \right)$$

$$= \frac{u^3 y}{3 \times 2} - \frac{(a+b) u^4}{2 \times 3 \times 4}$$

$$P_i = \frac{u^3 y}{6} - \frac{u^4 (a+b)}{24} \checkmark$$

$$m^2 + (a+b)m + ab = 0$$

$$m^2 + am + bm - ab = 0$$

$$m(m+a) + b(m+a) = 0$$


$$(m+a)(m+b) = 0$$

$$\therefore m = -a$$

$$m = -b$$

$$\therefore C.F = \Phi_1(y-a) + \Phi_2(y-b)$$

D'Alembert's Solution of Wave Equation



→ The one dimensional wave equation is,

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$$

where $c^2 = \frac{T}{S}$

Then $u(x, t)$ is a function of v and ω , $v = x + ct$

$\frac{\partial v}{\partial t} = c$ and $\frac{\partial v}{\partial x} = 1$

The solution can be obtained by introducing the two independent variable v and ω defined by

$$v = x + ct \quad \text{and} \quad \omega = x - ct$$

$$w = x - ct$$

Differentiating v w.r.t. x

$$\frac{\partial w}{\partial t} = -c \quad \frac{\partial w}{\partial x} = 1$$

$$\begin{aligned} \frac{\partial v}{\partial x} - \frac{\partial v}{\partial t} \cdot \frac{\partial v}{\partial x} + \frac{\partial v}{\partial t} \cdot \frac{\partial w}{\partial x} \cdot \frac{\partial w}{\partial t} \\ = \frac{\partial v}{\partial x} + \frac{\partial v}{\partial t} \end{aligned}$$

Again diff. w.r. to x

$$\frac{\partial^2 v}{\partial x^2}$$

D'Alembert's solution of wave equation

The wave eqⁿ is $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$ --- (i)
 $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$

let $v = x + ct$ and $w = x - ct$

then $u(v, t)$ is a function of v and w

$$v = x + ct$$

$$\frac{\partial v}{\partial t} = c \quad \text{and} \quad \frac{\partial v}{\partial x} = 1$$

$$w = x - ct$$

$$\frac{\partial w}{\partial t} = -c \quad \frac{\partial w}{\partial x} = 1$$

by chain rule,

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{\partial u}{\partial v} \cdot \frac{\partial v}{\partial t} + \frac{\partial u}{\partial w} \cdot \frac{\partial w}{\partial t} \\ &= \frac{\partial u}{\partial v} \times 1 + \frac{\partial u}{\partial w} \times -1 \\ &= \frac{\partial u}{\partial v} - \frac{\partial u}{\partial w} \end{aligned}$$

Assuming that all the partial derivatives are continuous.

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial}{\partial v} \left(\frac{\partial u}{\partial v} \right) + \frac{\partial}{\partial w} \left(\frac{\partial u}{\partial w} \right)$$

$$\text{or} \quad \frac{\partial^2 u}{\partial t^2} = \frac{\partial}{\partial v} \left(\frac{\partial u}{\partial v} + \frac{\partial u}{\partial w} \right) \cdot \frac{\partial v}{\partial t} + \frac{\partial}{\partial w} \left(\frac{\partial u}{\partial v} + \frac{\partial u}{\partial w} \right) \cdot \frac{\partial w}{\partial t}$$

$$\text{or} \quad \frac{\partial^2 u}{\partial t^2} = \left(\frac{\partial^2 u}{\partial v^2} + \frac{\partial^2 u}{\partial v \cdot \partial w} \right) \cdot 1 + \left(\frac{\partial^2 u}{\partial w \cdot \partial v} + \frac{\partial^2 u}{\partial w^2} \right) \cdot 1$$

$$\text{or} \quad \frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial v^2} + 2 \frac{\partial^2 u}{\partial v \cdot \partial w} + \frac{\partial^2 u}{\partial w^2}$$

$$\therefore u_{ttt} = u_{vvv} + 2u_{vuw} + u_{www} \quad \text{--- (2)}$$

Again for $U_{tt} \left(\frac{\delta^2 u}{\delta t^2} \right)$

$$\begin{aligned} \frac{\delta u}{\delta t} &= \frac{\delta u}{\delta v} \cdot \frac{\delta v}{\delta t} + \frac{\delta u}{\delta w} \cdot \frac{\delta w}{\delta t} \\ &= \frac{\delta u}{\delta v} \cdot C + \frac{\delta u}{\delta w} (-C) \\ &= C \left(\frac{\delta u}{\delta v} - \frac{\delta u}{\delta w} \right) \end{aligned}$$

$$\begin{aligned} \frac{\delta^2 u}{\delta t^2} &= C \frac{\delta}{\delta t} \left[\frac{\delta u}{\delta v} - \frac{\delta u}{\delta w} \right] \\ &= C \frac{\delta}{\delta v} \left(\frac{\delta u}{\delta v} - \frac{\delta u}{\delta w} \right) \frac{\delta v}{\delta t} + C \frac{\delta}{\delta w} \left(\frac{\delta u}{\delta v} - \frac{\delta u}{\delta w} \right) \cdot \frac{\delta w}{\delta t} \\ &= C \left[\frac{\delta^2 u}{\delta v^2} - \frac{\delta^2 u}{\delta v \cdot \delta w} \right] \delta v \cdot C + C \left[\frac{\delta^2 u}{\delta v \cdot \delta w} - \frac{\delta^2 u}{\delta w^2} \right] \cdot (-C) \\ &= C^2 \left[U_{vv} - 2U_{vw} + U_{ww} \right] \end{aligned}$$

$$U_{tt} = C^2 \left[U_{vv} - 2U_{vw} + U_{ww} \right] \quad \text{--- (3)}$$

putting eqⁿ (2) and eqⁿ (3) in eqⁿ (1)

$$U_{tt} = C^2 U_{uu}$$

$$C^2 \left[U_{vv} - 2U_{vw} + U_{ww} \right] = C \left[U_{vv} + 2U_{uv} + U_{ww} \right]$$

$$\begin{aligned} \text{on } U_{vv} - 2U_{vw} + U_{ww} - U_{vv} - 2U_{uv} + U_{ww} &= 0 \\ \text{or } -4U_{vw} &= 0 \end{aligned}$$

$$\text{or } U_{v \cdot w} = 0$$

$$\frac{\delta^2 u}{\delta v \cdot \delta w} = 0$$

Integrating we get,

$$u = \int f(w) dw + \Phi(v)$$

$$\text{or } u = \Psi(w) + \Phi(v)$$

where,

$$\Psi(w) = \int f(w) dw$$

and $\psi(u)$ and $\phi(v)$ are arbitrary function.

$$u = \phi(u+ct) + \psi(u-ct)$$

$$\therefore u(v, t) = \phi(v+ct) + \psi(v-ct)$$

is the required ~~eqn~~ D'Alembert's solⁿ of wave equation.

D'Alembert's Solⁿ of wave Eqⁿ when initial condition are given.

→ we have,

$$\text{wave Eq}^n \text{ is } u_{tt} = c^2 u_{vv} \quad \dots \dots \text{①}$$

initial conditions

$$u(v, 0) = f(v)$$

$$u(v, 0) = g(v)$$

$$u(v, t) = \phi(v+ct) + \psi(v-ct) \quad \dots \dots \text{②}$$

solⁿ of wave equation.

we impose given integration constants in order to remove arbitrary function ϕ and ψ .

Diff. Eqⁿ ② w.r.t t

we get,

$$\frac{\partial u}{\partial t} = \frac{\partial \phi(v+ct)}{\partial t} \times \frac{\partial (v+ct)}{\partial t} + \frac{\partial \psi(v-ct)}{\partial t} \times \frac{\partial (v-ct)}{\partial t}$$

$$\therefore u_t = \phi'(v+ct) \cdot c + \psi'(v-ct) \times (-c)$$

$$u_t = c \phi'(v+ct) - c \psi'(v-ct) \quad \dots \dots \text{③}$$

Using initial condition in Eqⁿ ③, we get,

$$u(v, 0) = \phi(v+0) + \psi(v-0)$$

$$u(v, 0) = \phi(v) + \psi(v)$$

$$\phi(v) + \psi(v) = f(v) \quad \dots \dots \text{④}$$

Using initial condition:

$$u_t(u, 0) = g(u)$$

from eqⁿ ③

$$u_t(u, 0) = c \phi'(u+0) - c \psi'(u-0)$$

$$\therefore g(u) = c \phi'(u) - c \psi'(u) - \dots \quad \textcircled{5}$$

$$\phi'(u) - \psi'(u) = \frac{1}{c} g(u)$$

Integrating both sides,

$$s=u$$

$$\phi(u) - \psi(u) = \frac{1}{c} \int_{s=a}^u g(s) ds \dots \textcircled{6}$$

Adding eqⁿ ④ and ⑥

$$2\phi(u) = f(u) + \frac{1}{c} \int_{s=a}^u g(s) ds$$

$$\textcircled{7} \quad \phi(u) = \frac{1}{2} f(u) + \frac{1}{2c} \int_a^u g(s) ds$$

$$\textcircled{8} \quad \phi(u+ct) = \frac{1}{2} f(u+ct) + \frac{1}{2c} \int_a^{u+ct} g(s) ds \leftarrow \textcircled{7}$$

(Subtracting eqⁿ ④ and ⑥)

$$2\psi(u) = f(u) - \frac{1}{c} \int_{s=0}^u g(s) ds$$

$$\textcircled{9} \quad \psi(u) = \frac{1}{2} f(u) - \frac{1}{2c} \int_{s=0}^u g(s) ds$$

$$\textcircled{10} \quad \psi(u-ct) = \frac{1}{2} f(u-ct) - \frac{1}{2c} \int_{s=a}^u g(s) ds \leftarrow \textcircled{9}$$

Putting $\Phi(u+ct)$ and $\Psi(u-ct)$ from eqⁿ ⑦ and ⑧ into eqⁿ ②

we get

$$s = u + ct$$

$$u(u, t) = \frac{1}{2} f(u+ct) + \frac{1}{2c} \int_{s=0}^u g(s) ds + \frac{1}{2} f(u-ct) - \frac{1}{2c}$$

$$s = u - ct$$

$$\int_{s=a}^u g(s) ds \quad \left[\because - \int_a^b f(u) du = \int_b^a f(u) du \right]$$

$$u(u, t) = \frac{1}{2} f(u+ct) + \frac{1}{2c} \int_a^{u+ct} g(s) ds + \frac{1}{2} f(u-ct) +$$

$$\frac{1}{2c} \int_{s=u-ct}^a g(s) ds \quad \left[\because \int_a^b f(u) du + \int_b^c f(u) du = \int_a^c f(u) du \right]$$

$$u(u, t) = \frac{1}{2} [f(u+ct) + f(u-ct)] + \frac{1}{2c} \int_{s=u-ct}^{u+ct} g(s) ds \dots \text{---} ⑨$$

when initial Velocity is zero.

$$u_t(u, 0) = g(u) = 0$$

eqⁿ ⑨ becomes,

$$u(u, t) = \frac{1}{2} [f(u+ct) + f(u-ct)] \#$$

which is the required initial condition of D'Almeida's solⁿ of wave equation.

Solution of heat eqⁿ by Fourier Series :-

$$\text{PDE} = \frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$$

→ The heat equation is,

$$u_t = c^2 u_{xx} \dots \text{--- (i)}$$

Boundary Condition:

$$u(0, t) = 0 \quad y > 0$$

$$u(l, t) = 0$$

Initial Condition:

$$u(x, 0) = f(x)$$

$$\text{Let } u(x, t) = XT \dots \text{--- (2)}$$

where,

$X(x)$ = function of x only.

$T(t)$ = function of t only.

Diff. w.r. to x and t , we get

from eqⁿ (2)

$$\frac{\partial u}{\partial t} = XT', \frac{\partial u}{\partial x} = X'T \text{ and } \frac{\partial^2 u}{\partial x^2} = X''T$$

Using these values in eqⁿ (1)

$$XT' = c^2 X''T$$

Separating Variables,

$$\frac{X''}{X} = \frac{T'}{c^2 T} = k \text{ (say)}$$

From above

$$\frac{X''}{X} = k$$

$$X'' - kX = 0$$

$$\left(\frac{d^2}{dx^2} - k \right) X = 0 \dots \text{--- (3)}$$

$$\frac{T'}{c^2 T} = k$$

$$T' = k c^2 T$$

$$T - k c^2 T = 0$$

$$\left(\frac{d}{dt} - k c^2 \right) T = 0 \dots \text{--- (4)}$$

which are ODE

solving eqⁿ of ODECase 1:where $k = \lambda^2 > 0$ from eqⁿ ③

Auxiliary equation is,

$$m^2 - \lambda^2 = 0$$

$$m = \pm \lambda$$

$$\therefore y = C_1 e^{m_1 n} + C_2 e^{m_2 n}$$

from eqⁿ ④

$$m = c^2 \lambda^2$$

$$m_1 = c^2 \lambda^2$$

$$\therefore T(t) = C_3 e^{c^2 \lambda^2 t}$$

$$\therefore X(n) = C_4 e^{\lambda n} + C_5 e^{-\lambda n}$$

Case 2:when $k = 0$ from eqⁿ ③

$$m^2 = 0$$

$$\therefore X(n) = C_4 n + C_5$$

from eqⁿ ④

$$m + \lambda^2 c^2 = 0$$

$$\therefore m = -\lambda^2 c^2 < 0$$

$$\therefore X(n) = n(C_4 + C_5)$$

$$\therefore T(t) = C_6 e^{\lambda^2 c^2 t}$$

$$\therefore T(t) = C_6$$

Case 3:when $k = -\lambda^2 < 0$ from eqⁿ ③

$$m^2 + \lambda^2 = 0$$

$$m = \pm \lambda i$$

$$= a + ib$$

$$y = (e^{an} (C_1 \cos bn + C_2 \sin bn))$$

from eqⁿ ④

$$m + \lambda^2 c^2 = 0$$

$$m = -\lambda^2 c^2$$

$$\therefore T(t) = C_7 e^{-\lambda^2 c^2 t}$$

$$\therefore X(n) = C_8 \cos \lambda n + C_9 \sin \lambda n$$

among those three cases, solution of Case 3 is consistent. It is because temperature is a periodic function of u and it must contain trigonometry function.

The feasible solⁿ is,


$$u_{(n,t)} = C_9 e^{-\alpha^2 C^2 t} + C_7 \cos \alpha u + (C_8 \sin \alpha u)$$

$$\therefore u_{(n,t)} = e^{-\alpha^2 C^2 t} (A \cos \alpha u + B \sin \alpha u)$$

$$\text{where, } A = C_7 C_9$$

$$B = C_8 C_9$$

Mathematical Modeling of one dimensional Heat flow



Let us consider flow of heat by conduction in an uniform bar with sides insulated so that the loss of heat from sides by conduction is negligible.

Physical Assumption

- i) The bar is made up of heat conducting material.
- ii) The bar is uniform and thin so that the temp at all points of cross section is constant.

Let one end of the bar is placed at origin which is at higher temp. and u denotes temperature function. Then u at any point of the bar depends upon distance (u) and time (t). i.e. $u = u(u, t)$

The amount of heat flowing inside the section of bar depends on

- a) Cross sectional area of bar. i.e. A
- b) Thermal Conductivity of bar i.e. K
- c) Rate of temp w.r.t to position i.e. $\frac{\partial u}{\partial u}$ (temp. gradient)

Let P and Q be two nearly points on bar at u and $u+\Delta u$ from origin. The amount of heat flowing into Section PQ from P is given by

$$Q_{in} = -KA \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_u \quad \text{--- (1)}$$

The amount of heat flowing out from Q is given by

$$Q_{out} = -KA \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_{u+\Delta u} \quad [(-) \text{ indicate that } x \text{ increases and } u \text{ decreases}]$$

from eq (1) and eq (11)

$$Q_{in} - Q_{out} = -KA \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_u + KA \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_{u+\Delta u}$$

Heat gained in Section PQ is given by,

$$Q_{in} - Q_{out} = KA \left[\left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_{u+\Delta u} - \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_u \right] \quad \text{--- (11)}$$

Let s be the specific heat capacity and ρ be the density of the material.

The Volume is given by $\pi r \Delta u A$ and mass = $\rho \cdot \Delta u A$. Then,

amount of heat gained by section PQ is,

$$\begin{aligned} &= m \times s \times \text{rate of change of temp per unit time} \\ &= \rho \cdot \Delta u \cdot A \times \frac{\frac{\delta u}{\delta u}}{\delta t} \quad \text{--- (iv)} \end{aligned}$$

Since eq (11) and (iv) are equal, so,

$$\frac{\frac{\delta u}{\delta u}}{\delta t} \rho \Delta u A = KA \left[\left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_{u+\Delta u} - \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_u \right]$$

$$\therefore \frac{\frac{\delta u}{\delta u}}{\delta t} = \frac{KA}{\rho s} \left[\left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_{u+\Delta u} - \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_u \right] \frac{\Delta u}{\Delta u}$$

taking limit $\Delta u \rightarrow 0$ on both sides.

$$\lim_{\Delta u \rightarrow 0} \frac{\frac{\delta u}{\delta u}}{\delta t} = \lim_{\Delta u \rightarrow 0} \frac{KA}{\rho s} \left[\left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_{u+\Delta u} - \left(\frac{\frac{\delta u}{\delta u}}{\frac{\delta u}{\delta u}} \right)_u \right] \frac{\Delta u}{\Delta u}$$

$$a. \frac{S_4}{S_1} = \frac{k}{S_1} \cdot \frac{S^2 4}{S_1 u^2}$$

$$a. \frac{S_4}{S_1} = \frac{c^2}{S_1} \frac{S^2 4}{S_1 u^2}$$

where $c^2 = \frac{k}{S_1}$ is diffusivity of material.

$\therefore u_t = c^2 u_{xx}$ is required P.D.E of One dimensional heat flow eqⁿ.

Mathematical modeling of wave eqⁿ:

Physical assumption:

i) The motion takes place on the vertical plane only and each particle of string execute transvers vibration only.

ii) The string is perfectly elastic and it transmit tension only but not bending and shearing force.

iii) The motion of string is subject to only a constant tension (T) and no other external forces.

Consider an elastic string of length 'L' (when it is stretched).

Fix two ends of string at $x=0$ and $x=L$ along x-axis.


Let $u(x,t)$ denotes deflection of string at any position 'x' & time 't'.

Consider a position PQ of the string. Since there is no resistance to bending the tension is traditional to the string at each point.

Let T_1 and T_2 be the tension at P and Q respectively.

Since the points of string moved vertically, there is no motion in the horizontal direction, so that the tension produced at horizontal component must be constant. i.e.

$$T_1 \cos \alpha = T_2 \cos \beta = T = \text{constant} \quad \text{--- (1)}$$



The vertical component T_1 and T_2 are $-T_1 \sin \alpha$ and $T_2 \sin \beta$ at P and Q respectively.

Resultant force acting on the portion PQ of the string is given by

$$T_2 \sin \beta - T_1 \sin \alpha \quad \text{--- (2)}$$

Let s be the mass of undeflected string per unit length.

$$\text{i.e. } s = \frac{m}{\Delta u}, \quad m = s \cdot \Delta u$$

By newton second law of motion, the resultant force is equal to ma . i.e. $F = ma$, $a = \frac{s^2 u}{s t^2}$

$$= s \cdot \Delta u \cdot \frac{s^2 u}{s t^2} \quad \text{--- (3)}$$

Since eqⁿ (2) and (3) are equal so,

$$T_2 \sin \beta - T_1 \sin \alpha = s \cdot \Delta u \cdot \frac{s^2 u}{s t^2} \quad \text{--- (4)}$$

Dividing eqⁿ (4) by eqⁿ (1)

$$\frac{T_2 \sin \beta}{T_1 \cos \beta} - \frac{T_2 \sin \alpha}{T_1 \cos \alpha} = \frac{s \Delta u}{T} \cdot \frac{s^2 u}{s t^2}$$

$$\therefore \frac{s^2 u}{s t^2} = \frac{T}{s \Delta u} (\tan \beta - \tan \alpha)$$

Since $\tan \beta$ and $\tan \alpha$ are slopes of string at $u + \Delta u$ and u .

$$\tan \alpha = \left(\frac{s u}{s u} \right)_u, \quad \tan \beta = \left(\frac{s u}{s u} \right)_{u + \Delta u}$$


then eqⁿ (4) becomes,

$$\frac{s^2 u}{s t^2} = \frac{T}{s \Delta u} \left(\frac{s u}{s u} \right)_{u + \Delta u} - \left(\frac{s u}{s u} \right)_u$$

taking limit $\Delta u \rightarrow 0$ on both sides,

$$\lim_{\Delta u \rightarrow 0} \frac{s^2 u}{s t^2} = \lim_{\Delta u \rightarrow 0} \frac{T}{s} \left(\frac{s u}{s u} \right)_{u + \Delta u} - \left(\frac{s u}{s u} \right)_u$$

Δu



$$\therefore \frac{S^2 u}{S t^2} = \frac{T}{S} \frac{S^2 u}{S v^2}$$

$$c^2 = \frac{T}{v^2}$$

$\therefore u_{tt} = c^2 u_{vv}$ is the p.d.e of wave eqⁿ